




THAI  
GEOSCIENCE  
JOURNAL

Special Issue

# THAI GEOSCIENCE JOURNAL

Vol. 2 No. 1 January 2021



Dr. Sangad Bunopas

Published By  
Department of Mineral Resources  
Geological Society of Thailand

## Editorial Committee

### Honorary Editors

Dr. Sommai Techawan

Department of Mineral Resources and  
Geological Society of Thailand

Mr. Niwat Maneekut

Department of Mineral Resources, Thailand

Mr. Montri Luengingkasoot

Department of Mineral Resources, Thailand

### Advisory Editors

Prof. Dr. Clive Burrett

Palaeontological Research and Education Centre,  
Mahasarakham University, Thailand

Prof. Dr. Katsuo Sashida

Mahidol University, Kanchanaburi Campus, Thailand

Prof. Dr. Nigel C. Hughes

University of California, Riverside, USA

Prof. Dr. Punya Charusiri

Department of Mineral Resources and  
Geological Society of Thailand

### Editor In Chief

Dr. Apsorn Sardud

Department of Mineral Resources, Thailand

### Associate Editors

Prof. Dr. Che Aziz bin Ali

University Kebangsaan Malaysia, Malaysia

Prof. Dr. Clive Burrett

Palaeontological Research and Education  
Centre, Mahasarakham University, Thailand

Assoc. Prof. Dr. Kieren Howard

Kingsborough Community College, CUNY, USA

The Graduate Center, CUNY, USA

American Museum of Natural History, USA

Dr. Pol Chaodumrong

Department of Mineral Resources

and Geological Society of Thailand

Prof. Dr. Punya Charusiri

Department of Mineral Resources

and Geological Society of Thailand

### Editorial Secretary

Department of Mineral Resources, Thailand

Ms. Cherdchan Pothichaiya

Ms. Paveena Kitbutrawat

Mr. Inthath Chanpheng

Ms. Peeraporn Nikhomchaiprasert

Ms. Jeerawan Mermana

Dr. Puangtong Puangkaew

Mr. Kitti Khaowiset

Ms. Roongrawee Kingsawat

Mr. Kittichai Tongtherm

Mrs. Sasithon Saelee

Ms. Kunlawadee Nirattisai

Ms. Thapanee Pengtha

### On the cover



1. Dr. Sangad Bunopas (1937-2020) was born and raised in Chaiyaphum near the edge of the Khorat Plateau in north eastern Thailand. After graduating from Chulalongkorn University in 1960, Sangad joined the Department of Mineral Resources (DMR) in July 1961. Within a few years, He was sponsored to study abroad by the DMR and supported initially by a Colombo Plan scholarship. He completed the rigours of a BSc (Honours) degree before progressing to an MSc programme (1965-6) in New Zealand. He returned to regional mapping in Thailand and then from 1977 to 1981 completed a PhD at the Victoria University of Wellington supervised by Professor Paul Vella. He was the first established the word Shan-Thai microcontinent in 1981.



2. Ancient cratonic areas; I, Indochina (including eastern Thailand); SC, South China and ST, Shan-Thai (eastern Burma, Western Thailand and Northwestern Malay Peninsula). Adjacent fold-belts are formed of thick mainly marine Paleozoic to Triassic sediments and tholeiitic volcanic rocks that accumulated along the margins of the cratons. Ophiolites lie between contiguous fold belts. Sinistral faulting and orocline bending occurred mainly during the Jurassic and Cretaceous. K, Khorat Basin; CM, Chiangmai; V, Vientiane; WM, West Malay Peninsula; EM, East Malay Peninsula; B, Bentong ophiolite line.



3. Stratigraphic sections through the Chum Phuang sandpits in Nakhon Ratchasima (Khorat) Province showing Normal (N = Brunhes) and Reversed (R = Matuyama) magnetostratigraphy. Unabraded, hollow sphere, tektite in Chum Phuang section shown as Tn and reworked abraded tektites as Ta. From Haines et al. (2004).



# THAI GEOSCIENCE JOURNAL

---

Vol. 2 No. 1 (Special Issue)  
January 2021



**Published By**  
Department of Mineral Resources and Geological Society of Thailand

Copyright © 2021 by the Department of Mineral Resources of Thailand  
Thai Geoscience Journal website at <http://www.dmr.go.th>



## Preface

January 2021, is the 129<sup>th</sup> anniversary of Department of Mineral Resources that was first established as the Royal Department of Mines and Geology in 1891 and recently changed to Department of Mineral Resources. On this occasion we proudly present this second volume of Thai Geoscience Journal (TGJ) as a special issue. This issue is presented and dedicated to the memory of the late Dr. Sangad Bunopas, an important geologist of Department of Mineral Resources and Thailand who initiated research in Thai geological sciences and named “Shan-Thai and Indochina micro-continents” in an early tectonic study. Moreover, he was the crucial person that drove Thai geology and the Department of Mineral Resources to internationalization. His work inspired stratigraphic and tectonic research in Thailand from the 1960's until now. He will be also remembered as an inspiring, pioneering and influential geologist and as the father of Thai stratigraphy. Thank you very much for our reviewers, editors, Dr. Bunopas's friends and family for their valuable help in this issue.

The New Year has just started, on behalf of the Department of Mineral Resources and the Geological Society of Thailand, I would like to express my sincere care to TGJ authors, reviewers, editors, members, researchers and your families. Wishing you a happy new year and good health. Even though we are in the hardest time for people around the world with the COVID-19 outbreak, we will embrace to hope and dream to get through this time together.

Thank you very much.

(Dr. Sommai Techawan)  
Director-General of Department of Mineral Resources  
President of Geological Society of Thailand



## From the editors

It is our great pleasure that the Thai Geoscience Journal was successful in the first edition of the journal and we distributed more than 200 hard copies and made it also available on the web to every authors, reviewers, editors, members and researchers. Welcome to the Volume 2 No.1 (Special Issue) of Thai Geoscience Journal (TGJ). This is a special issue to the memory of the late Dr. Sangad Bunopas, who is one of the most famous geologists in Thailand and the SE Asia region. This special issue will not happened without the loves, faiths and helps from many people who have ever been Dr. Sangad Bunopas's friends and colleagues and also his family.

This issue aims to review his valuable work for inspiring the next generation of researchers to dare to show their hypothesis, scientific imagination, and ideas. We have the honor from Prof. Dr. Clive Burrett who was Dr. Bunopas's colleague and friend to revise his research as in "The man who made an impact-the scientific work of Dr. Sangad Bunopas". We also received the honor and permission from Dr. Bunopas's family to give us his huge and final article about his interests, ideas and research on the geology of Thailand and cometary collisions for planetary science in "Australasian Pleistocene cometary impact and its effects in Thailand". This article is edited by Prof. Dr. Clive Burrett with contributions from Prof. Dr. Kieren Howard (New York) and his many colleagues.

Thank you very much for your interest in TGJ. Finally, we hope to receive more valuable articles in the next issue in July 2021 and wishing you a happy new year.

Thank you very much.

(Dr. Apsorn Sardud)  
Editor-in-Chief

Thai Geoscience Journal



Thai Geoscience Journal

Vol. 2 No. 1

(Special Issue for Dr. Sangad Bunopas)

January 2021

ISSN 2730-2695

# LIST OF CONTENTS

---

	Page
The man who made an impact – the scientific work of Dr Sangad Bunopas <b>Clive Burrett</b>	1-15
Australasian Pleistocene cometary impact and its effects in Thailand <b>Sangad Bunopas</b>	16-109

Any opinions expressed in the articles published in this journal are considered the author's academic autonomy and responsibility about which the editorial committee has no comment, and upon which the editorial committee takes no responsibility

ข้อคิดเห็นของบทความทุกเรื่องที่ดีพิมพ์ลงในวารสารฯ ฉบับนี้ถือว่าเป็นความคิดอิสระของผู้เขียน กองบรรณาธิการไม่มีส่วนรับผิดชอบ หรือไม่จำเป็นต้องเห็นด้วยกับข้อคิดเห็นนั้น ๆ แต่อย่างไร

## The Man Who Made Impact – The Scientific Work of Dr Sangad Bunopas

Clive Burrett<sup>1,\*</sup>

<sup>1</sup>*Palaeontological Research and Education Centre, Mahasarakham University,  
Kantarawichai, Maha Sarakham, 44150 Thailand*

*\*Corresponding author: cfburrett@gmail.com*

Received 29 December 2020; Accepted 15 January 2021

### Abstract

Sangad Bunopas published extensively on the geology of Thailand and left a legacy of maps, reports, stratigraphic nomenclature and insightful papers covering most of the kingdom. His 1981 PhD thesis published in 1982 laid the foundation of modern Thai stratigraphy. His highly cited papers on the tectonics of South East Asia, often with Professor Paul Vella, were based on Sangad's extraordinary first-hand knowledge of Thai geology which allowed him to draw the first reliable tectonic subdivisions of the whole country. Although controversial at the time, these tectonic subdivisions and hypotheses have since been accepted as generally correct in that the Shan-Thai Terrane (as part of Sibumasu) rifted from Australian crust in the early Permian, drifted and warmed as it crossed the Palaeotethyan Ocean and collided with the Indochina Terrane in the late Triassic. His stimulating palaeomagnetically constrained, combined palinspastic-palaeogeographic maps for the Permian to Cenozoic were well in advance at the time and have not been attempted again in the subsequent 40 years.

His controversial later work, which attempted to explain the widespread, anomalous loess in northeast Thailand, the Australasian tektite strewn field, the fluvial deposits in Nakhon Ratchasima (Khorat) Province and their middle Pleistocene faunas and flora as the result of a cometary impact at 788 ka has received some support and some criticism but has nevertheless stimulated a considerable research and a lot of discussion.

Over more than 50 years, Sangad made many major contributions to Thai geology, South East Asian tectonics and meteoritic research. His wit, wisdom and iconoclastic approach to earth and planetary science will be sorely missed by his numerous friends and colleagues.

**Keywords:** biography, cometary impact, loess, Quaternary stratigraphy, Shan-Thai Terrane, tectonics, Thailand

### 1. INTRODUCTION

Dr Sangad Bunopas (1937-2020) was born and raised in Chaiyaphum near the edge of the Khorat Plateau in northeastern Thailand, surrounded by areas of scenic geology and geomorphology. To the east lay the Mesozoic sandstones of the Khorat Plateau and to the west the Central Plains with Cenozoic cover of marine Palaeozoic rocks. After graduating from Chulalongkorn University in 1960, Sangad joined the Department of Mineral Resources (DMR) in July 1961. Within a few years, he was sponsored to study abroad by the DMR and supported initially by a Colombo Plan scholarship. He was encouraged and helped in his overseas

studies by his wife and by senior DMR staff Pisoot Sudasna Na Ayudhaya (who became Deputy Permanent Secretary of the Minister of Industry), Saman Buravas, Manus Veraburras and significantly, in the light of his pioneering work on tektites, by Dr Kaset Pitakpaivan who later became Director-General of the Department of Industrial Promotion.

Sangad completed the rigours of a BSc (Honours) degree before progressing to an MSc programme (1965-6) in New Zealand. He returned to regional mapping in Thailand and then from 1977 to 1981 completed a PhD at the Victoria University of Wellington supervised by Professor Paul Vella. Sangad was lucky to study at



Fig. 1 Sangad Bunopas. Photos kindly supplied by the Bunopas family

Victoria University as the geology of New Zealand is diverse, rugged, spectacular and active (Figs. 2-3) and the local geologists are renowned for their excellence, enthusiasm for fieldwork and all-round abilities. This was excellent training which he later put to good use working on Thai geology. He graduated with an MSc in 1966 at a time when the theory of plate tectonics was being developed and then applied to orogenic belts around the world. Sangad's regional mapping and his PhD work on palaeomagnetism would soon lay the foundations for his new ideas on the tectonics of South East Asia

## 2. REGIONAL GEOLOGY OF THAILAND

His early work for the DMR, based in Bangkok, was in regional mapping and special projects. Unlike today, there was little infrastructure in Thailand in the late 1960's to 1970's. Roads were very bad, vehicles old and poorly maintained, road cuts were limited and thick jungle still

covered large areas of the country. Communist insurgency made some areas dangerous for government personnel. "Traverses of the Western Mountains and some parts of the Sukhothai Foldbelt often require transportation by elephant..." (Raksaskulwong and Bunopas, 1985, p.9). With some understatement they reported that progress on foot and by elephant made mapping very slow.

His early assignments were in several provinces on a wide variety of rocks and economic deposits. He investigated the geology and mineral resources of Amphoe Mae Thaeng in Chiang Mai Province in 1961, lignite and oil shale in Li, Lamphun Province (1962 and 1963), the geology of Song in Phrae Province in 1967, the geology of the Nam Phrom Dam in Chaiyaphum Province (Bunopas, 1971) and tin deposits in Rayong Bay in eastern Thailand in 1973. With DMR colleagues, Sangad produced 1: 250,000 maps and reports for geologically significant quadrangles in central Thailand such as Phitsanalok (Bunopas, 1974b and 1976a), Nakhon Sawan (Bunopas, 1976c, 1980) and Suphanburi (Bunopas, 1976d). Part of the Suphanburi mapping project led to an important paper in the first volume of the Journal of the Geological Society of Thailand on the fascinating geology of Bo Phloi and Uthai Thani in western Thailand (Bunopas and Bunjitradulya, 1975). They mapped a variety of Palaeozoic rocks, metamorphic and igneous rocks with comments on the economic resources including the famous Bo Phloi sapphires. They discussed the confusing lithostratigraphic nomenclature at the time concerning the Kanchanaburi Series, Phuket Group and Tanaosri Group and most significantly recognised the several thrust faults such as the large, low-angle Chong Insi Thrust where the Siluro-Devonian rocks are overthrust on gneisses. The inclusion of structural data, and especially the recognition of thrust faults, was very unusual for Thai geology at that time and was a feature of many of Sangad's (with DMR colleagues) regional papers including the still very interesting structural geology chapter 7 of his 1981 thesis. An example of his emphasis on structural geology is shown in Fig. 4.

An important study on the famous (and as a prison, infamous) island of Ko Tarutao in southern Thailand followed in 1978 with DMR colleagues Sahat Muenlek and Veerapong Tansuwan (Bunopas et al., 1978). Cambrian



Fig. 2 Gladstone East Coast Road. Cold and wet South Island, New Zealand – a long way from tropical Thailand. (From Bunopas, 1966, MSc thesis, Victoria University, New Zealand).



Fig. 3 Eringa Cyclothem, near Eringa Station, New Zealand. The area was studied in Wairarapa, a little north of Wellington. In the Wainuiru River Valley in Gladstone near Masterton. Miocene and Pliocene mudstones younging west (left hand), are overlain by limestone cyclothem with coquina limestones and the terrestrial Hawera Series (From Bunopas, 1966, Victoria University, Wellington, MSc thesis).




Fig. 4 From Bunopas (1982, fig.42) showing an example of the structural cross-sections in his thesis. Section from north of Lampang to west of Phrae, northern Thailand.

fossils had previously been found there by DMR Director Saman Buravas in 1958 and described by Teichii Kobayashi but little work had been done on these important and fossiliferous Lower Palaeozoic sections in the intervening 20 years. They mapped and described the sandstones and limestones and recognised important faults thereby laying the foundations of scores of palaeontological papers, reports and theses in subsequent years. Work on Ko Tarutao and in mainland Satun was strongly encouraged and supported by Sangad. These foundational studies and discoveries eventually led to Satun being recognised as the first UNESCO Geopark in Thailand.

Another regional study of note was written-up by Lertsin Raksakulwang and Sangad in 1985 on the complex geology of 3000 square kilometres of Mae Hong Son Province in northwest Thailand. They included structural, lithostratigraphic and sedimentological details including many cross sections, stratigraphic columns and Landsat maps of lineaments with much of the work carried out on elephants and on foot across mountainous terrain. Within a few years this area became another well-studied and palaeontologically and tectonically significant area.

Sangad also produced several important and influential syntheses of stratigraphic correlations across Thailand (Bunopas 1974a, 1976, 1983, 1992b, 1994). Some of these papers were written prior to his massive encyclopedia of Thai geology (his PhD thesis) and others subsequent to Bunopas (1981, 1982). All were


influential and there is hardly a chapter of recent books on the geology of Thailand that do not quote these studies extensively.

A paper with Sirot Salypongse, Father Henri Fontaine and Daniel Vachard (Bunopas et al., 1983) summarised the Permian stratigraphy of Thailand and included important new palaeontological data. Sangad also worked with Father Fontaine on the fossiliferous Devonian limestone of Loei Province, in northeastern Thailand. This was followed by many significant papers and monographs by Fontaine, Vachard and DMR colleague Dr Voravudh Suteethorn, mainly on the Devonian to Permian palaeontology of Thailand.

Throughout his career, Sangad emphasised the importance of biostratigraphy to Thai geology. From the 1970's he cooperated with palaeontologists in many countries and had at least three species named after him. These are the Triassic bivalved arthropod *Euestheria bunopasi* (Kobayashi, 1975), the Permian brachiopod *Rhamneria bunopasi* Waterhouse 1981, and the Ordovician trilobite *Rossaspis bunopasi* Stait, Burrett and Wongwanich 1984.

### 3. TECTONICS OF SOUTH EAST ASIA

Armed with an enormous data-base of Thai geology acquired after detailed studies in every corner of the kingdom, Sangad developed a new tectonic model for the whole of South East Asia with his mentor and supervisor Professor Paul Vella. This model was first published in Bunopas and Vella (1978) and enlarged on in



Silurian-Devonian lithofacies (excepting peninsular Thailand). *cs*, carbonate shelf facies; *cc*, continent derived clastics, sandstone and shale; *L*, land; *bab*, back-arc basin (chert and shale); *va*, metavolcanics (metatuff and agglomerate); *fab*, fore-arc basin (tuff, greywacke, marble and chert); *t*, trench deposits, pelitic greenschist. *PSO*, Pha Som Ophiolite and *SKO*, Sra Kaeo Ophiolite thought to have been emplaced in the Triassic; *css*, shelf sandstone, shale and limestone. Note: No Silurian-Devonian thick marginal deposits are known east of the ophiolite belt. *MP*, Mae Ping Fault Zone; *TP*, Three Pagodas Fault Zone.

Fig. 5 From Bunopas (1982, fig 141b) showing one of several diagrams outlining Sangad's major tectonic subdivisions of Thailand (for the Siluro-Devonian) offset by major transcurrent faults. Note the recognition, mapping and offset of the Sra Kaeo ophiolite in eastern Thailand and the proposed back-arc basin in northern Thailand in what we would now term the Inthanon Zone (or terrane).

subsequent years (Bunopas and Vella 1983a, Bunopas et al., 1990, 2003, 2004). The model was developed fully in chapter 9 of his 801 page tour de force PhD from Wellington which was then published as special paper number 5 of the Geological Survey of the DMR (Bunopas, 1981, 1982). Unusually for a PhD thesis, the first 443 pages is a comprehensive, country-wide summary and it may well be considered the first book on the geology of Thailand. It has been cited over 400 times. Any geologist working in Thailand can still dip into these chapters and find something significant that they either did not know or had forgotten.


The first 443 pages of Sangad's thesis contains numerous maps and useful lithostrati-

graphic and correlation charts with the text full of new paleontological lists and sedimentological insights. Sangad included thin section descriptions and numerous chemical analyses of igneous rocks. He redefined and clarified old, loosely defined lithostratigraphic units and on almost every page named and defined numerous new formations and groups, many of which are still in every-day use. Bunopas (1982) may be regarded as the foundation of modern Thai stratigraphy.

The 100 pages of Chapter 8 of his thesis concern Sangad's palaeomagnetic studies on both Palaeozoic and Mesozoic rocks. His preliminary palaeomagnetic results were published with Dr Kaset Pitakpaivan and J. Sukroo in 1978 (Bunopas et al., 1978). His conclusions on the rotation of Indochina and his Mesozoic results have been confirmed by later work. Unfortunately, many of his Permian samples had high alpha 95's ('statistical uncertainty' or scatter) which were unacceptably large but even his low inclination results with moderate alpha 95's on Permian limestones appear to have been validated by recent studies. Sangad fully realised the limitations of his Palaeozoic palaeomagnetic studies and on returning from Wellington collected a large number of orientated cores on Cambrian and younger Palaeozoic rocks. Sadly, hardware and software problems meant that no further reliable results could be obtained.

Generally Sangad's basic tectonic concepts have stood the test of time. He drew his Shan-Thai Terrane to include western Thailand and Malaysia and eastern Myanmar and regarded what is now termed the Inthanon Zone or Terrane as a back-arc basin of his Sukhothai Arc. He identified the ophiolitic Nan Suture and for the first time suggested its extension to the ophiolite of Sra Kaeo in eastern Thailand which he briefly described - also for the first time. He emphasised the contrast between the late Carboniferous and early Permian faunas of the Shan-Thai (cool Gondwana) and Indochina (tropical Cathaysian) terranes which were separated by the Nan-Sra Kaeo suture. He subdivided Thailand into distinctive tectonic belts, offset by major transcurrent faults and showed their palaeogeographic evolution through time (Fig. 5).

Sangad suggested Shan-Thai rifted from the Australian sector of Gondwana, with the



Suggested early Carboniferous reconstruction showing Western Thailand and Burma (Shan-Thai) adjacent to north-west Australia. Malay Peninsula is left off the map because its position and orientation are uncertain. FAB, fore-arc basin; BAB, back-arc basin; VA, volcanic arc. M.N.P.F., Mae Ping Fault Zone; T.P.F., Three Pagodas Fault Zone; MF, Mandalay Fault.

Fig. 6 A novel and iconoclastic palinspastic reconstruction of Thailand adjacent to northwest Australia in the early Carboniferous, figure 144 in Bunopas (1981, 1982). This is one of many stimulating and provocative palinspastic reconstructions in his thesis.

deposition of the glacial pebbly mudstones of the Kaeng Krachan Group (Bunopas and Vella, 1984, 1989) in the early Permian, drifted into lower and warmer palaeolatitudes during the middle Permian to Early Triassic and collided with Indochina in the Norian (Late Triassic).

Sangad did not confine himself to the Palaeozoic-Triassic but wrote an interesting paper on Cenozoic rifting and the development of the Gulf of Thailand (Bunopas and Vella, 1983b). Unusually, he attempted to remove Mesozoic deformation and Cenozoic faults and their movements and incorporated his palaeomagnetic results, showing block rotations, in order to draw combined palinspastic palaeogeographic restorations for South East Asia (Bunopas 1981, Figs. 151-155b).

These restorations (Fig. 6) have not been

attempted to the same level of detail since and contrast markedly with the inchoate 'blobs' drawn on many recent palaeogeographic maps. A feature here is that Sangad bravely extended his initial ideas where most conventional geologists would have finished. This pushing of hypotheses to their limits - and sometimes beyond was also characteristic of his later work on impacts.

Apart from the excision of the Upper Palaeozoic limestones and Devonian to Triassic cherts of the Inthanon Zone from the Shan-Thai Terrane, arguments on subduction directions and the inclusion of Shan-Thai in an enlarged Sibumasu Terrane, his initially controversial proposals, have been confirmed by most subsequent studies. Today few would disagree with his major conclusion that Shan-Thai (as a part of Sibumasu) was part of northwest Australian crust that

rifted off in the early Permian, warmed-up in its drift across Palaeo-Tethys and collided with Indochina in the Late Triassic.

#### 4. IMPACTS, TEKTITES, LOESS AND COMETS

DMR Director General Dr Kaset Pitakpaivan was one of the first to study tektites in Thailand and it was not surprising that he influenced the young Sangad to eventually study them. Dr Kaset worked with Dr Virgil Barnes from Texas and made the first regional scientific study of these fascinating black glasses at about the same time that Sangad joined the DMR (Barnes and Pitakpaivan, 1962). At the time, many prominent and brilliant planetary scientists argued that tektites were extraterrestrial with the most prominent theory being that they were blasted from volcanoes on the moon (Chapman and Larsen, 1963; O'Keefe, 1976). Barnes and Pitakpaivan collected tektites throughout northeast Thailand and emphasised the importance of layered or Muong Nong tektites including the large 12.8 kg specimen donated by Dr Kaset to the DMR museum (Barnes, 1971).

Huge samples of Muong Nong tektites which were offered for sale by Mr Boomun Poonyathiro, owner of the House of Gems, Bangkok weigh up to 24 kg (Fig. 7). Detailed field studies in Ubon Ratchathani by P. Fiske, Prinya Phutthapiban and John Wasson (1996) confirmed the abundance and stratigraphic position of Muong Nong tektites on a lateritic surface overlain by loess. All subsequent workers showed conclusively that Australasian strewn field tektites are not extraterrestrial but the result of a bolide impact or impacts at 788 ka, centred in the Indochina region that blasted tektites all the way to Tasmania and as microtektites across the Indian Ocean, western Pacific Ocean and all the way to Antarctica. Tektites are so abundant in South East Asia that in some areas almost every villager has a collection and jewellery shops in Charoen Krung, Bangkok literally have buckets full of them.

In 1987 Dr Ramsay Ford (with the writer and Sangad) toured northeast Thailand (Isarn) and collected tektites from almost every province. We found a large disk tektite horizontally embedded in the middle of a 3- thick section of loess near Khon Kaen and found that the majority of tektites were restricted to the junction of a distinctive laterite and an overlying section of loess or rarely within the loess itself (Ford,



Fig. 7 Probably the largest Muong Nong tektite yet found and was on display at the House of Gems in Charoen Krung, Bangkok. This and related specimens may be from Ubon Ratchathani on the Thai-Lao border or possibly from Laos. This specimen was shown to Sangad Bunopas, Ramsay Ford, Clive Burrett and Kieren Howard at various times by Mr Boomun Poonyathiro. Weight is 24 kg. 1999 photograph kindly supplied by Prof. Kieren Howard.

1988). The loess covers NE Thailand extends to the Central Plain and varies between 4 m and 1 m in thickness and compares with the famous yellow earth of China but differs in detail (Sibava, 1993; Nichol and Nichol, 2015, see Bunopas, 2021 this volume). An interesting study by Paul Vella, Sangad and Vorakul Kaewyana that does not appear to have been followed-up, showed that loess extends to northern Thailand and covers river terrace deposits. This potentially provides constraints on dating river terraces and also provides clues to uplift rates (Vella et al., 1987).

Loess in Europe and China is related to windy environments close to glaciated areas and its ubiquity in northeastern Thailand is anomalous and difficult to explain on current models of loess formation (Nichol and Nichol, 2015). Sangad was already interested in the tektite problem and he soon wrote papers on the catastrophic effects of a large impact in Indochina close to the area of the large Muong Nong tektites eastern Isarn, Laos and Vietnam (Bunopas, 1989, 1990, 1992a). He specifically included the loess as a result of an impact and therefore suggested a c.800 ka age on the basis of finding pristine tektites within and just below this enigmatic deposit. Sirot Salyapongse and Kaset Pitakpaivan studied the Thai loess and compared it to Chinese loess. They found glass beads in the Thai loess and also, interestingly, in Pleistocene trees (Salyapongse and Pitakpaivan, 1999).

Sangad coined the term *catastroloess* for the Thailand loess. He included fluvial deposits from along the palaeo-Mun River valley, which clearly are not loess and it is unlikely that this term will survive.

He soon extended the catastrophic impact model to include the fluvial sands along the Mun River Valley of Nakhon Ratchasima Province (Khorat) containing large charcoalified and partially burnt logs, small plant material and pollen (some burnt) and numerous vertebrate fossils (Bunopas and Khositantont, 1999c, Bunopas et al., 1997, 1998, 1999 a, b, 2001, 2002a, b, 2003, 2004, 2005 and 2007; Charusiri et al., 2002). When Sangad showed me the Khorat sandpits and their abundant apparently burnt logs, I was immediately struck by the resemblance to the multitude of trees felled by the huge air-blast of the Tunguska, Siberia explosion in 1908. Sangad's correlation of the Khorat sands with the 788 ka impact was a step too far for many as it was already apparent that many but not all of the vertebrates in the sandpits are Miocene in age and not Pleistocene (Chaimanee, 2004, 2007; Duangkrayom et al., 2017). The situation is complicated by the sandpit operators who use high-powered hoses to sort and collect the poorly consolidated sands. As a result, vertebrate fossils are rarely found in situ but are found either much lower in the stratigraphy or at the bottom of the pits. The biostratigraphic situation was clarified by Sato (2002) and it appears that definite in situ though of course, fluvially transported, Miocene fossils are generally found in the basal parts of the pits and middle Pleistocene such as *Stegodon*, crocodiles and turtles in the upper parts (e.g. Lauprasert et al., 2019; Naksri et al., 2019). Of course, in a flood-prone, fluvial environment older vertebrates may be reworked into younger sands so the biostratigraphic and taphonomic context becomes complicated. The situation in northeast Thailand and beyond is also complicated by the difficulties dating Quaternary deposits with all available methods being subject to considerable problems, caveats and uncertainties.

Sangad encouraged detailed work on the Khorat sand-pits by Kieren Howard (1999) and then with Howard collaborated with Dr Peter Haines (Australia) and palaeomagnetist Professor Jason Ali (Hong Kong) (helped by the late Dr Somboon Khositantont and the writer) in a magnetostratigraphic study of the pits. The

advantage of magnetostratigraphy is that if consecutive units are present then it is unlikely that they have not been reworked or reset. The disadvantage is that short geomagnetic excursions, such as within the Brunhes Epoch, may be mistaken for epoch boundaries or vice versa. Haines et al. (2004) revealed a transition from present-day Normal magnetization to Reversed at 14 m depth at Ban Tha Chang and at 9 m depth 40 km away at Chum Phuang (Fig. 8). This was interpreted as the transition from the Matuyama Epoch (R) to the Brunhes Chron (N) which polarity reversal is now dated at 773 +/- 5 ka (Haneda et al., 2020). The magnetism of the older samples from the pits below the Reversed level was equivocal and so the reversal could be the result of short-lived magnetic excursions during the late Pleistocene as interpreted by magnetostratigraphic studies at Khok Sung also in Nakhon Ratchasima (Khorat) Province (Suraprasit et al., 2015). However, the discovery of an in situ, rare, un-abraded, delicate and hollow sphere, and definitely unreworked Australasian tektite, at the reversal level (Fig. 8) strongly supports Sangad's and Peter Haines et al's (2004) interpretation. The stratigraphic situation in the Khorat sandpits is further complicated by the detailed palynological work of Yang and Grote (2018 a, b) who reported thermoluminescence dating on 7 samples and two C14 dates ranging from 172,739 +/- 22,400 a BP to 27,332 +/- 3000 a BP in a 3.25 m section from about 13.5 m to 10.0 m depths which covers the same depth interval as the study by Haines et al. (2004). Yang and Grote (2018 b) suggest major vegetational and palaeoclimatic changes from cool to warm to cool, and correlation to parts of two glacial periods (Riss and Wurm) during the deposition of this 3.25m of sediments.

Whether or not Sangad (and Howard, 1999, 2011a, b; Howard et al., 2003 and Haines et al., 2004) were correct in ascribing some parts of the upper sections of the sandpits and their contained biota to the 788 ka Australasian impact event, the importance of the studies in the Khorat sandpits to a wide variety of earth, planetary and biological sciences suggest that more efforts should be made on dating studies along with palaeontological, sedimentological and palynological studies in the sandpits. The pits are ephemeral and most, if not all, will end up filled with water. More careful interdisciplinary studies such as at Khok Sung by Dr. Jaroon




Fig. 8 Stratigraphic sections through the 22 m of the Ban Tha Chang and Chum Phuang sandpits in Nakhon Ratchasima Khorat Province showing Normal (N=Brunhes) and Reversed (R= Matuyama) magnetostratigraphy. Unabraded, hollowsphere, tektite in Chum Phuang section shown as T<sub>n</sub> and reworked abraded tektites as T<sub>a</sub>. From Haines et al., (2004). Colourslide kindly supplied by Dr Peter Haines (Australia).

Duangkrayom et al. (2014) are needed. If Sangad was correct, then northeastern Thailand contains globally unique evidence of the catastrophic environmental effects of a major impact event. This is, of course, of global scientific interest but is also important as a possible tourist attraction that will bring employment to the local community.

With characteristic intellectual exuberance Sangad then extended his hypothesis to include events far from Khorat. He included the demise of Miocene vertebrates throughout Thailand (including Khorat) in his model which is highly unlikely. However, his speculation should prompt all who work in Khorat to carefully demarcate the Miocene biota from the Pliocene and Pleistocene and, if possible, the early from the later Pleistocene faunas and floras. He further suggested that the well-known, almost certainly, Miocene, fish beds in Phetchabun Province and shell beds near Bangkok, most likely to be Holocene, were also the result of the impact. These speculations are possible but unlikely on present evidence.

He further extended his hypothesis to the extinction of Miocene faunas in Australia. Although the widespread occurrence of unworked tektites is a very useful horizon in Australian Quaternary stratigraphy and the 788 ka event may well have had environmental and biotic consequences on that huge continent, any correlation with Quaternary megafaunal extinctions in Australia is very unlikely.

However, his suggestion that our hominid ancestor *Homo erectus* witnessed and probably suffered from the tektite fall is well supported by studies in Java and in Yunnan (Hou et al., 2000). The effects throughout SE Asia of the 788 ka 'cataclysm' on *Homo erectus* are recently well documented by Li et al. (2021) and provide considerable support for another of Sangad's 'wild' ideas.

Sangad suggested that the eroded pillars known as Sao Din (earth pillars) (Fig. 9) found from northern Thailand to Sra Kaeo include deposits of loess caused by the 788 ka impact. These deposits remain undated directly but




Fig.9 "Sao Din" earth pillars at Nanoi, south of Nan city northern Thailand.

archaeological excavations in the Nanoi Basin, south of Nan in northern Thailand have found stone tools very similar to those in the Chinese Bose Basin closely associated with tektites as discussed by Li et al. (2021) and Hou et al. (2000). Zeitoun et al. (2012) suggest an age of 1 to 0.5 Ma for the Sao Din stone tools so perhaps this is another of Sangad's 'wild ideas' that will soon be proven correct.

Sangad also claimed that the 788 ka impact in Indochina led to epeirogenesis in northern Thailand which he named the Inthanon Uplift. However, the uplifted deposits in northern Thailand (see Bunopas, 2021, this volume) are not well dated and this idea must, on present knowledge, be labelled as speculative. However, the huge Indochina impact or impacts must have caused severe ground shaking and the writer and others have observed faults in the Pleistocene sections of northeastern Thailand that may be due to this event. The tectonic effects of this cataclysmic impact event should certainly be considered.

Almost a hundred years ago the famous Australian geologist Sir Edgeworth David and colleagues noted that several tektite-like glasses plotted on a great circle band around the earth and suggested an extraterrestrial origin (David

et al., 1927). On the basis of extensive geochemical studies all meteoritic workers now accept that tektites are molten target rocks due to impacts by either asteroids or comets. Ramsay Ford (1988) suggested a similar path from SE Asia to Tasmania for the Australasian tektites. A date of 816 +/- 7 ka for the impact glass known as Darwin Glass scattered around Darwin Crater in western Tasmania (Howard, 2004) almost overlaps (to within 18 ka) with the best date for Australasian tektites (788 +/- 3 ka) and, with dates for the Zhamanshin crater in Kazakhstan (about 920-650 ka) led to the great circle idea being resurrected by several authors. Influenced by the work of Professor John Wasson (UCLA), Sangad postulated a cometary impact. In 1994, Sangad was strongly impressed by the well documented impact of huge fragments of the Shoemaker-Levy comet on Jupiter. Sangad envisaged a similar mechanism for the possibly simultaneous production of the Zhamanshin, Indochina and Darwin craters, their impact glasses and strewn fields. He suggested an initial impact of a comet fragment in Kazakhstan, followed by an impact in Indochina and then at Mount Darwin, Tasmania (Bunopas, 2021, this volume). This scenario is possible and the Shoemaker-Levy comet break-up and massive

impacts into Jupiter's atmosphere substantially increases the plausibility of Sangad's hypothesis. However, general acceptance of the idea is delayed because of the at least 18,000 year difference between the Darwin Crater and Australasian tektites and the, as yet, poorly dated glasses and crater at Zhamanshin.

The recent announcement of the discovery of a possible Australasian tektite source crater beneath Quaternary basalt of the Bolaven Plateau in Laos would, if correct, be the final crater to be identified and associated with a major tektite strewn field (Sieh et al., 2020). The crater is in the most likely position within the strewn field based on tektite morphology, microtektite and tektite geochemistry and sandstone target rock age and chemistry (Ford, 1988; Howard, 2011a, b). Our work on the same outcrops as Professor Sieh and colleagues does not support their conclusions but if Sieh et al. (2019) are found to be correct then their small 15 km diameter crater would probably support the late Professor John Wasson's hypothesis of multiple small impact craters rather than one major Indochina crater (Wasson, 1991). If correct, then the Bolaven Crater impact was of sufficient size to account for the events summarised by Sangad in northeast Thailand but possibly not large enough to account for the whole Australasian strewn field.

## 5. MENTORING

One of Sangad's greatest contributions was to encourage and support both the study and research of Thai geologists at overseas universities and foreign geologists both senior and junior to carry on research in Thailand. He mentored, helped and encouraged DMR geologists Drs Thanis Wongwanich and Pol Chaodumrong to pursue PhD studies at the University of Tasmania in Australia and Somkiet Maranate and Worakul (Vorakul) Kaewayna to complete MSc degrees respectively on palaeomagnetism and Quaternary sedimentology at Victoria University in New Zealand.

He helped and encouraged Clive Burrett, Steve Carey, Ramsay Ford, Peter Haines, John Long, John Shergold and Kieren Howard (from Australia), John Wasson and Gudron Wasson (from USA) and Shigeki Hada (from Japan) and initially Father Henri Fontaine (from France) on a wide variety of fruitful projects all of which led

to many publications in the international scientific literature.

Sangad helped graduate students from Tasmania who studied the Permian glacial deposits in Phuket (John Hills), the Lower Palaeozoic on Ko Tarutao (Tim Akerman and Roger Mason) and Kieren Howard who studied the fluvial deposits in Khorat.

## 6. CONCLUSIONS

Had Sangad retired without publishing his work on tectonics and impacts he would be remembered as a helpful, inspiring, pioneering, widely read, highly productive and influential geologist and as the father of Thai stratigraphy. By using his unrivalled and hard-won knowledge of Thai geology to construct, tectonic subdivisions of Thailand and an original and provocative tectonic model of South East Asia he placed Thai geology on a global stage, mentored generations of young Thai geologists and induced many foreign scientists to carry-out collaborative research in the kingdom.

Like the pioneer of continental drift theory, Alfred Wegener, Sangad extended his impact theory too far, making outrageous claims that could be easily criticized often leading to the dismissal of all of his ideas. Although some of Sangad's extreme ideas on the effects of the 788 ka impact will be consigned to the waste basket of disproved hypotheses (joining some of Wegener's) many of his ideas on the Quaternary have gained support and are being currently cited and favourably discussed (e.g. Li et al., 2021; Zak et al., 2019).

Sangad had and continues to have a huge influence on Thai geoscience and with his seminal work on tectonics and cometary collisions continues to have a big impact on international earth and planetary science.

## 7. ACKNOWLEDGEMENTS

Dr Sangad welcomed me to Thailand and inducted me into Thai geology. He generously imparted his vast knowledge of Thai geology and later meteoritics and encouraged me with characteristic quiet humour and patience. His DMR teams particularly Dr Thanis Wongwanich, Dr Pol Chaodumrong and the late Dr Somboon Khositantont and Mr Charoen Butrmuang supported me in numerous ways. Professor Che Aziz Ali, Professor Kieren Howard, Dr Apsorn

Sardsud and Dr Pol Chaodumrong kindly read the manuscript and made many useful comments. Dr Peter Haines, Professor Kieren Howard, Dr Pol Chaodumrong and the Bunopas family helped with figures and photographs

## 8. REFERENCES

- Barnes, V. E. 1971. Description and origin of 12.8-kg layered tektite from Thailand (abstract). *Meteoritics*. 6 (4), 249.
- Barnes, V.E. & Pitakpaivan, K. 1962. Origin of Indochinite tektites. *Proceedings of the National Academy of Sciences, USA*. 48, 947-955.
- Bunopas, S. 1961. Report on the geology and mineral deposits at Huai Tad-Bang Kud, Chiangmai. DMR Bangkok.
- Bunopas, S. 1962. Report on investigations of lignite and oil shale deposits, Li, Lamphun. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1967. Report on the investigation of geology of Amphoe Song – Rong Khwang, Phrae Province, Department of Mineral Resources, Bangkok.
- Bunopas, S. 1969. Report on the geology of Amphoe Tha Pla-Nam Pat and Pha Som Dam. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1966. Upper Miocene – Pleistocene Stratigraphy and the Paleontology of Gladstone, Wararapa New Zealand. Unpublished M.Sc.Thesis, Geology Department, Victoria University of Wellington, New Zealand.
- Bunopas, S. 1971a. On the geology and stratigraphy of the Nam Phrom Dam and its vicinities. *Geological Society of Thailand Newsletter*, 4, 17-40.
- Bunopas, S. 1971b. Jurassic beach sandstone. *Mineral Resources Gazette*. 16 (1), 40-48.
- Bunopas, S. 1972a. Preliminary report on the geological survey of Phitsanalok Quadrangle, 1:250,000 map. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1972b. Brief note on the Phitsanalok Sheet. *Geological Society of Thailand Newsletter*. 4, (1-3), 17-40.
- Bunopas, S. 1973. On the tin deposit in Rayong Bay, eastern gulf of Thailand. *Geological Society of Thailand Newsletter*. 6 (5-6), 120-128.
- Bunopas, S. 1974a. On the Lower Palaeozoic rocks in Thailand. *Geological Society of Thailand Newsletter*. 7 (3), 61-83.
- Bunopas, S. 1974b. Geological map of Changwat Phitsanalok, 1:250,000. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1975a. On the Carboniferous volcanic sequence and red beds at Khao Luang, Sukhothai. Department of Geological Sciences, Chiang Mai University. Special Paper 1. *Proceedings of the Conference on the Geology of Thailand, 1973*. 2, 39-40.
- Bunopas, S. 1975b. Permo-Triassic rocks in the Lansang Fault Zone. Department of Geological Sciences, Chiang Mai University. Special Paper 1. *Proceedings of the Conference on the Geology of Thailand, 1973*. 1, 39-40.
- Bunopas, S. 1976a. On the stratigraphic successions in Thailand- A preliminary summary. *Journal of the Geological Society of Thailand*. 2, 31-58.
- Bunopas, S. 1976b. Geology and Mineral Resources of Phitsanalok. Department of Mineral Resources, Bangkok. Report on Investigations. 16 Parts 1 and 2, 1-217, 30 plates.
- Bunopas, S. 1976c. Geological map of Nakhon Sawan, 1:250,000 series. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1976d. Geological map of Suphanburi, 1:250,000 series. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1980. Geology and mineral resources of Changwat Nakhon Sawan. Department of Mineral Resources, Geological Survey Report 1. Bangkok.
- Bunopas, S. 1985. Review on tectonic events of north-eastern Thailand (abstract). Conference on geology and mineral resources development of the northeast, Thailand, Khon Kaen University.
- Bunopas, S. 1980. Geology and Mineral Resources of Changwat Nakhon Sawan Quadrangle, Department of Mineral Resources, Geological Survey Report 1.
- Bunopas, S. 1981. Paleogeographic history of western Thailand and adjacent parts of Southeast Asia - a plate tectonics interpretation, Ph.D Thesis, 810 pp., University of Wellington, New Zealand, 1981. 810 pp., 164 figs., 21 tables.
- Bunopas, S. 1982. Palaeogeographic history of Western Thailand and adjacent parts of Southeast Asia. Geological Survey of Thailand Paper 5, Royal Thai Department of Mineral Resources, Thailand. 1-810 pp 164 figs., 21 tables.
- Bunopas, S. 1983. Paleozoic succession in Thailand. In: Nutalaya, P., ed., *Proceedings of the Workshop on Stratigraphic Correlation of Thailand and Malaysia (Vol. 1)*. Haad Yai, Thailand. 39–76.
- Bunopas, S. 1989. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Abstract. Pacific Science Congress, Seoul, Korea.
- Bunopas, S. 1990. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Proc. of the 16th Conf. on Sciences and Technology of Thailand, 25-27 October, 1990, Bangkok. p. 512-513.
- Bunopas, S. 1992a. Tektites - Their origin and the Early Quaternary catastrophic destruction in Northeastern Thailand and south Indochina. proceedings 29th international geological congress. 24 Aug.- 3 Sept. 1992. Kyoto, Japan, Abstracts Vol. 3 (3), II-12-4 0-8 2114, p. 650.
- Bunopas, S. 1992b. Regional Stratigraphic Correlation in Thailand. In: Piancharoen, C., ed., *Proceedings of the National Conference on Geological Resources of Thailand: Potential for Future Development*. Department of Mineral Resources. Bangkok, Thailand: 189–208.
- Bunopas, S. 1994. The Regional Stratigraphy, Paleogeographic and tectonic events of Thailand and Continental Southeast Asia. In: Angsuwathana, P., Wongwanich, T., Tansathien, W., et al., eds., *Proceeding of the International Symposium on Stratigraphic Correlation of Southeast Asia*, 2–24
- Bunopas, S. 2021. Australasian Pleistocene cometary impact and its effects in Thailand. *Thai Geoscience Journal*, 2(1).
- Bunopas, S., & Bunjitradulya, S. 1975. Geology of Amphoe

- Bo Phloi, north Kanchanaburi, with special notes on the 'Kanchanaburi Series'. *Journal of the Geological Society of Thailand* 1, 51-67.
- Bunopas, S., Burrett, C. & Hada, S. 1990. Stratigraphy and paleogeographic evolution of the upper Paleozoic-early Mesozoic continental collision in Thailand (abstract). *Proc. Dev. Geol. Thailand year 2000*, Chulalongkorn University, Bangkok.
- Bunopas, S., Fontaine, H., Salyapongse, S. & Vachard, D. 1983. Permian paleogeography in southeast Thailand evidenced by new discoveries. *Journal Geological Society of Thailand*. 6, 17-21.
- Bunopas, S., & Khositantont, S. 1999. Geology and National Development in accordance with the King's new theory for self-sufficient agriculture as viewed from the Buntharik event (in Thai). In: *The Special Memorial Book for the King's 72 Birthday in 2542 (1999), contribution made to the King, "The Geographer of The Nation" by The Rajabhat Institute Buriram: 3 pp.*
- Bunopas, S., Khositantont, S., & Wasson, J. T. 1997. Evidence of the Early Global Disaster and Destruction from Extraterrestrial Impacts Comet in Northeast Thailand and South Indochina within the Australian Tektites Field: The last Mass Extinction, (extended abstract). *Proc. of the national conf. on stratigraphy and tectonics evolution of Southeast Asia and the South Pacific*. 19-24 August. 1997, Bangkok, Thailand, 434-435.
- Bunopas, S., Muenlek, S., & Tansuwan, V. 1978. Geology of Tarutao Island. *Journal of the Geological Society of Thailand*. 6, 121-138.
- Bunopas, S., Pitakpaivan, K., & Sukroo, J. 1978. Preliminary palaeomagnetic results from Thailand sedimentary rocks. In: *Proceedings of the 3rd Regional Conference on Geology and Mineral Resources of Southeast Asia*, Nutalaya, P. (ed.). Bangkok. 24-32.
- Bunopas, S., & Vella, P. 1978. Late Palaeozoic and Mesozoic structural evolution of northern Thailand: a plate tectonic model. In: *Proceedings of the 3rd Regional Conference on Geology and Mineral Resources of Southeast Asia*, Nutalaya, P. (ed.). Bangkok, 133-140.
- Bunopas, S., & Vella, P. 1983a. Tectonic and geologic evolution of Thailand. In: Nutalaya, P. (ed.) *Proceedings Of the Workshop on stratigraphic correlation of Thailand and Malaysia*, Haad Yai, Thailand, 8-10 September 1983, Tech. Paper 1, 307-323.
- Bunopas, S., & Vella, P. 1983b. Opening of the Gulf of Thailand - rifting of continental Southeast Asia, and late Cenozoic Tectonics, *Journal of the Geological Society of Thailand*, 6, 1-12.
- Bunopas, S., & Vella, P. 1984. Phuket-Kaeng Krachan Group, A rifted continental margin deposit with effect of ice-rafting from Gondwana. *Abstr. paper presented at GEOSEA V.*, 9-13 April, Kuala Lumpur.
- Bunopas, S., & Vella, P. 1989. "Pebbly mudstones", A composited assemblage of ice-rafted sediments on the western rifted marginal basin of Shan-Thai while drifting away from Gondwana during the Upper Paleozoic. *Proceedings of the 4th Intern. Symp. On Pre-Jurassic East Asia*, IGCP Project 224. Hat Yai and Langkawi Island, Thailand and Malaysia, 24 April-2 May 1989.
- Bunopas, S., Vella, P., Fontaine, H., Hada, S., Burrett, C., Haines, P., Howard, K. T. & Khositantont, S. 2003. The Pleistocene epeirogenesis, rapid vertical uplifting of the western mountains of Thailand in the late active himalayan extrusional continuum at 0.8 Ma impact, *Journal of Geological Society of Thailand*.
- Bunopas, S., Vella, P., Hada, S., Fontaine, H., Burrett, C., Haines, P., Khositantont, S. & Howard, K.T., 2001. Australasian tektites and catastrophic products enclosed in impact ejecta horizon from the Buntharik Impact Event in Thailand. *Symposium on Rodinia, Gondwana and Asia*, Osaka, Japan, *Gondwana Research* 4 (4), 586.
- Bunopas, S., Vella, P., Hada, S., Fontaine, H., Burrett, C., Haines, P., Khositantont, S. & Howard, K. T. 2002. Australasian tektites and catastrophic products enclosed in impact ejecta horizon from the Buntharik Impact Event in Thailand on old surface gravel beds marked real-time Asian Quaternary correlation, In: N. Mantajit (ed.), *The Symposium on Geology of Thailand (GEOTHA 2002)*, Bangkok, Thailand, 181-187.
- Bunopas, S., Vella, P., Hada, S., Fontaine, H., Haines, P.W., Burrett, C., Potisat, S., Wongwanich, T., Chaodamrong, P., Khositantont, S. & Howard, K.T. 2003. The Khorat Plateau early Quaternary vertically uplifting after long Cenozoic cratonization of the Mesozoic Khorat Group, its northeast tilting diverted Mae Khong River before the 0.8 Ma cometary impact catastroloess and the overflowing Mid-Pleistocene Buriram Basalt. *1st International Conference on Paleontology of Southeast Asia*, Mahasarakham University Journal 22, Special Issue.
- Bunopas, S., Vella, P., Fontaine, H., Hada, S., Burrett, C., Haines P., Howard K. T. & Khositantont. S., 2004. Did Shan-Thai Twice Mary with Indochina and India? *Discovery Quaternary Pushed-Pulled Himalayan Extensional Continuum Heritage to Major vertical Crustal Uplifting Inthanon Epeirogenesis, That Also Tilted Khorat Plateau Northeastwards Diverting Mae Khong River From Tonle Sap Prior to Buntharik Event, 0.8 MA Cometary Impact, Biriram Basalt And Asian Glacial Age*. *International Conference on the Geologic Evolution of East and Southeast Asia*. Department of Geology, Faculty of Science, Chulalongkorn University, 8-14 February 2004, Asia Hotel, Bangkok. (Full paper in the print in 2005)
- Bunopas, S., Vella, P., Fontaine H., Hada S., Burrett C., Haines P., Howard K. T. & Khositantont S. 2005. Latest Pliocene to early M. Pleistocene Inthanon Epeirogenesis, 2.0-0.65 Ma, in north Thailand connecting to late Himalayan extrusion continuum: An application of extraterrestrial / terrestrial real-time layer of catastroloess from Buntharik Cometary Impact Event (new). *Proceeding of the International Conference on Geology, Geotechnology and Mineral Resources of Indochina*. GEOINDO 2005, 28-30 Nov. 2005, Kosa Hotel, Khon Kaen. pp. 468-499.
- Bunopas, S., Vella, P., Fontaine, H., Hada, S., Burrett, C., Haines, P., Khositantont, S., Chintasakul, P., Chaodamrong, P., Charusiri, P. & Howard, K. T. 2007. Australasian Cometary Impact, 0.8ma3 Catastroloess Buried Alive Miocene1, Pliocene2 Faunas in Thailand And Central Australia, And Tektite-Bearing Flood Deposits In Ne. Thailand, New World's Discovery Extinction Age In Thailand (Indochina), and Australia Impact Fields4. (GEOTHA'07). *International Conference on Geology*

- of Thailand: Towards Sustainable Development and Sufficiency Economy, 21-22 November, Imeral Hotel, Bangkok, Thailand. pp. 30-43.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999a. Catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. *Journal Geological Society of Thailand*. 1999 (1), 142-151.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999c. Ancient analogs of burial alive extinction of the mastodons in catastroloess in Thailand, and of the last dinosaurs (in eggs) in Gobi Desert: further on tektites. *Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000*, October 28-29 1999, Bangkok, Thailand: 168-177.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C. & Khositantont, S., 1998. The early extraterrestrial collision with a whole comet in the Australasian tektite field, apparent evidences and discoveries from Thailand and East Asia (abstract). *Geosea '98. Ninth Regional Congress on Geology, Mineral and Energy Resources of Southeast Asia*. 17-19 August 1998, Shangri-La Hotel, Kuala Lumpur, Malaysia. Programme and Abstracts. 191-192.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H.; Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999a. Catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. *Journ. Geol. Soc. of Thailand* 1999 (1): 1-17.
- Bunopas, S., Yaemniyom, Ng. & Khositantont, S. 1999b: Catastroloess and its derivatives, the life time sustainable construction sands was originally high atmospheric settling in the Quaternary cometary impact in Thailand and Asia. *Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000*.
- Chaimanee, Yaovaluk, 2004. Fossils from Khok Soong and Ban Tha Chang, Khorat. In: *Numerous fossils in Paleontology of Thailand*, in 114 years of the Department of Mineral Resources. (in Thai)
- Chaimanee, Yaovaluk, 2007. Diversity of Cenozoic Mammals in Thailand: Paleoenvironment and Age Updated. (GEOTHAI'07) *International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy*. Geothai 2007, pp. 73-79. 21-22 November. Imeral Hotel, Bangkok, Thailand.
- Chapman, D. R., & Larson, H. K. 1963. On the Lunar origin of tektites. *Journal of Geophysical Research*. 68, 4305 - 4358.
- Charusiri P, Daorerk V, Krowchan V, Klongsara N, Kosuwan S, Srirattachatchawan V, & Santatiwongchai U. 2002. Quaternary tektites and their sediment hosts at Ban Tachang Sand Pit, Chaloeam Prakiat, Nakhon Ratchasima, NE Thailand: Stratigraphy and TL Ages. In: Mantajit N, et al. editors. *The Symposium on Geology of Thailand*. 2002 August 26-31; Bangkok, Thailand: Department of Mineral Resources, Bangkok; p. 178- 180.
- David T. W. E., Summers H. S. & Ampt G. A. 1927. The Tasmanian tektite - Darwin glass. *Proceedings of the Royal Society of Victoria*. 39, 167- 190.
- Duangkrayom J, Ratanasthien B, Jintasakul P, & Carling P.A. 2014. Sedimentary facies and paleoenvironment of a Pleistocene fossil site in Nakhon Ratchasima province, northeastern Thailand. *Quaternary International*. 325, 220-238. doi: 10.1016/j.quaint. 2013.07.048
- Duangkrayom, J., Wang, S.-Q., Deng, T., & Jintasakul, P. 2017. The first Neogene record of Zygodontodon (Mammalia, Proboscidea) in Thailand: implications for the mammutid evolution and dispersal in Southeast Asia. *Journal of Paleontology*. 91 (1), 179-193.
- Fiske P.S., Putthapiban P., & Wasson J. T. 1996. Excavation and analysis of layered tektites from northeast Thailand: Results of 1994 field expedition. *Meteoritics & Planetary Science*. 31, 36-41
- Ford, R., J. 1988: An empirical model for the Australasian tektite field. *Australian Journal of Earth Science*. 35, 483-490.
- Haines, P.W., Howard, K.T., Ali, J.R., Burrett, C., & Bunopas, S. 2004. Flood deposits penecontemporaneous with ~0.8 Ma tektite fall in NE Thailand: impact-induced environmental effects? *Earth and Planetary Science Letters*. 225, 19-28.
- Haneda, Y., Okada, M., & Kitamura, T. 2020. A full sequence of the Matuyama-Brunhes reversal in the Chiba composite section, Central Japan. *Progress in Earth and Planetary Sciences*. 7 no.44, doi.org/10.1186/s40645-020-00354-y
- Hou, Y., Potts, R., Baoyin, Y., Zhengtang, G., Deino, A., Wang, W., Clark, J., Guangmao, X., & Weiwen, H., 2000. Mid-Pleistocene Acheulean-Like Stone Technology of the Bose Basin, South China. *Science*, 287, 1622-1626.
- Howard, K. T. 1999. Distal environmental effects of the 770ka impact event. Unpublished Hons. Thesis at the University of Tasmania, Australia. 175 pp.
- Howard, K. T. 2004. Origin of Darwin Glass. Unpublished Ph. D. Thesis at the University of Tasmania, Australia. 362 pp.
- Howard K. T. 2011a. Volatile enhanced dispersal of high velocity impact melts and the origin of tektites. *Proceedings of the Geologists Association*. 122, 363-382.
- Howard, K.T. 2011b. Tektites in: Ridd, M.F., Barber, A.J., Crow, M.J. (Eds.), *The Geology of Thailand*, The Geological Society of London, pp. 574-591. doi. org/10.1144/GOTH.21
- Howard, K. T., Haines, P.W., Burrett, C., Ali, J., & Bunopas, S. 2003. Sedimentology of 0.8 Ma log-bearing flood deposits in north-east Thailand and mechanisms for pre-flood deforestation, *Proceedings, 8th International Congress on Pacific Neogene Stratigraphy*, Chiang Mai, Thailand, 2003, pp. 49-67.
- Kobayashi, T. 1975. Upper Triassic estheriids in Thailand and the conchostracan development in Asia in the Mesozoic era. *Geology and Palaeontology of Southeast Asia* 16, 57-90.
- Lauprasert, K., Watchajittaphan, P., Juanggam, S., & Bhuttarach, S. 2019. Freshwater crocodile, *Crocodylus siamensis* Schneider, 1801, from the Middle Pleistocene deposits in Chaloeam Phrakiat District, Nakhon Ratchasima, Thailand. *Annales de Paléontologie*. 105 (4), 269-274. doi:10.1016/j.annpal.2019.05.002
- Li, H., Lotter, M., Kuman, K., Lei, L., & Wang, W. 2021. Population dynamics during the Acheulian at 0.8 Ma

- in East and SE Asia: Considering the influence of two geological cataclysms. *Palaeogeography, Palaeoclimatology, Palaeoecology* 562, <https://doi.org/10.1016/j.palaeo.2020.109927>. (online-in press)
- Naksri, W., Tong, H., Lauprasert, K., Jintasakul, P., Suteethorn, V., Vidthayanon, C., & Claude, J. 2019. Plio-Pleistocene giant tortoises from Tha Chang sandpits, Nakhon Ratchasima Province, Thailand. *Annales de Paléontologie*. 105 (4), 257-267. doi:10.1016/j.annpal.2019.06.001
- Nutalaya, P., Vella, P., Bunopas, S., & Kaewyana, V., 1986. Quaternary processes in Thailand. In: Thiramongkol (ed.), *Proceeding of the Workshop on Economic, Geology, Tectonics Sedimentary Processes and Environmental of the Quaternary of Southeast Asia*. p.23-33, Bangkok.
- Nichol, J.E., & Nichol, D. W. 2015. Character and provenance of aeolian sediments in northeast Thailand. *Aeolian Research*. 19 (A), 5-14. <https://doi.org/10.1016/j.aeolia.2015.09.001>
- O'Keefe, J. A., 1976. *Tektites and Their Origin*. Elsevier. Amsterdam. 354 pp.
- Raksaskulwang, L., & Bunopas, S. 1985. Stratigraphy and tectonic evolution in the Mae Hong Son Province, north western Thailand. Unpublished report of the Royal Thai Department of Mineral Resources. 1-39.
- Salyapongse, S., & Pitakpaivan, K., 1999. Older eolian evidences on the Khorat Plateau. *Journal of the Geological Society of Thailand*, 1, 18-26
- Sato, Y. 2002. Preliminary report on the occurrence of fossil mammals in Nakhon Ratchasima, northeast Thailand, in: N. Mantajit (Ed.). *Proceedings of the Symposium on Geology of Thailand*, Department of Mineral Resources. Bangkok, 2002, pp. 230-232.
- Sibava, V. 1993. Quaternary Sequences in Southeast Asia and their regional correlation. In: ESCAP Atlas of Stratigraphy XII., *Quaternary Stratigraphy of Asia and the Pacific*, IGCP 296. Mineral Resources Development Series, No 62. United Nations. pp. 1 -29.
- Sieh, S., Herrin, J., Jicha, B., Schonwalder, D., Moore, J., Banerjee, P., Wiwegwin, W., Sihavong, V., Singer, B., Chualaowanich T., & Charusiri, P. 2020. Australasian impact crater buried under the Bolaven volcanic field, Southern Laos. *Proceedings of the National Academy of Science*. 117 (3), 1346-1353. <https://doi.org/10.1073/pnas.1904368116>
- Suraprasit, K., Jaeger, J.-J., Chaimanee, Y., Benammi, M., Chavasseau, O., Yamee, C., Tian, P., Panha, S., 2015. A complete skull of *Crocota crocota ultima* indicates a late Middle Pleistocene age for the Khok Sung (northeastern Thailand) vertebrate fauna. *Quat. Int.* 374, 34-45. <https://doi.org/10.1016/j.quaint.2014.12.062>.
- Vella, P., Bunopas, S., & Kaewyana, W. 1987. A loess covered alluvial gravel terrace near Chiangmai, Northern Thailand. *Geological Society of Thailand Newsletter*. 1987 (3), 29-31.
- Wasson J. T. 1991. Layered tektites: A multiple impact origin for the Australasian tektites. *Earth and Planetary Science Letters* 102, 95-109.
- Yang, F.-C., & Grote, P.J. 2018 a. Variation of fluorescence intensity in Pleistocene pollen and the correlation of fluvial deposits in the Khorat Plateau, Thailand, *Palynology*. 42(1), 28-35, DOI: 10.1080/01916122.2017.1303796
- Yang, F.-C., & Grote, P.J. 2018 b. Riverine vegetation and environments of a Late Pleistocene river terrace, Khorat Plateau, Southeast Asia. *Palynology*. 42 (2) 158-167. doi: 10.1080/01916122.2017.1296044
- Žák, K., Skála, R., Pack, A., Ackerman, L., & Křížová, Š. 2019. Triple oxygen isotope composition of Australasian tektites. *Meteoritics & Planetary Science*. 54 (5), 1167-1181. doi:10.1111/maps.13251
- Zeitoun, V., Forestier, H., Auetrakulvit, P., Khaokhiew, C., Rasse, M., Davtian, G., Winayalai, C., Tiamtinkrit, C. 2012. Discovery of a prehistoric site at Sao Din (Nanoi, Nan province, Northern Thailand): Stone tools and new geological insights. *Découverte d'un site préhistorique à Sao Din (Nanoi, province de Nan, Nord de la Thaïlande): études lithique et géologique préliminaires*. *Comptes Rendus Palevol*. 11, 575-580. doi:10.1016/j.crpv.2012.06.007

## Australasian Pleistocene cometary impact and its effects in Thailand

Sangad Bunopas<sup>1</sup>

with contributions from Kieren Howard<sup>2,3,4,\*</sup>

<sup>1</sup>*Deceased. Department of Mineral Resources of Thailand,  
 Rama VI Rd., Ratchathewi, Bangkok 10400, Thailand*

<sup>2</sup>*Physical Sciences, Kingsborough Community College, CUNY, New York, USA*

<sup>3</sup>*Earth and Environmental Sciences, the Graduate Center, CUNY, New York, USA*

<sup>4</sup>*Earth and Planetary Sciences, American Museum of Natural History, New York, USA*

*\*Corresponding author: khoward@gc.cuny.edu*

Received 18 January 2021 ; Accepted 28 January 2021

### Abstract

Tektites are silica-rich glasses distributed in 4 strewn fields around 3 definite impact craters in the USA, Ghana and Germany and the Australasian strewn field around possibly one or more craters in Indochina. The distribution, morphology, size, mineralogy, geochemistry and contained gases of Australasian tektites confirms their terrestrial origin from a sandstone target rock of approximately Jurassic age and an impact site or sites within or offshore Indochina. The Indochina impact is dated at about 788 ka. Layered tektites are known from a more limited area and probably are proximal to the impact crater or craters. The largest layered tektites weigh up to 24 kg and come from Ubon Ratchathani Province in NE Thailand or perhaps from neighbouring Laos. Dumbbell, spherical, teardrop and disc shaped splash-form tektites are found across SE Asia and southern China. Aero-ablated tektites (australite buttons) extend to the Indian Ocean and Australia and microtektites are found in Antarctica and in deep-sea cores from the Indian Ocean and the marginal seas of the western Pacific. In many deep-sea cores, microtektites occur just below the Matuyama-Brunhes magnetic reversal dated at about  $772.9 \pm 5.4$  ka. and occur with high pressure minerals such as coesite. Coesite and reidite along with silt-size detrital minerals within layered tektites are conclusive evidence of an impact origin. The abundance and distribution of iridium and microtektites in ocean cores has led several authors to calculations of a single crater size of between 15 km and >120 km with a most likely diameter of about 40 + km.

Both layered tektites and splash-form tektites occur at the junction of laterite and a loess-like sand or rarely within the loess, across much of SE Asia and this stratigraphic relationship is especially clear on the Khorat Plateau in NE Thailand. The widespread loess is different to typical Chinese loess, is palaeoclimatically anomalous and is considered herein as 'catastroloess' – the result of the 788 ka impact event.

Unreworked, tektites, including one very fragile hollow sphere, are also found in fluvial sands dated by thermoluminescence as 748 ka in Nakhon Ratchasima (Khorat) Province along with large, unusually very abundant charcoalified and partially charcoalified trees and smaller plant matter such as bamboo. This section of the sandpits along the palaeo-Mun River contains Pleistocene vertebrates in the upper sections and Miocene vertebrates below the tektite horizon. The tektite horizon occurs close to the Matuyama-Brunhes magnetic reversal. The sandpits are argued to contain globally unique evidence of a large extraterrestrial impact event.

Impact craters and abundant impact glasses at the Zhamanshin crater in Kazakhstan (c. 920-650 ka) and at the Darwin Crater in Tasmania, Australia ( $816 \pm 7$  ka) are argued to be contemporaneous with the Australasian strewn field and the result of a comet travelling southeastwards and disintegrating, creating catastrophic impacts first in Kazakhstan, then in Indochina and finally in Tasmania. This hypothetical trail of impacts is comparable to the disintegration and multiple impacts of the Shoemaker-Levy comet fragments into Jupiter's atmosphere in 1994.

These multiple impacts on Earth would have had profound effects on faunas, climate and sedimentation along the comet's path including on our immediate ancestor *Homo erectus* in

China, in Java and possibly in Lampang and Nan in Thailand. More evidence of impact in Thailand may be present near Nan and Phrae in northern Thailand and at Sra Kaeo in eastern Thailand where sao din or earth pillars are eroded 'catastroloess' and at Bo Phloi in western Thailand where sand-pits contain abundant charcolified logs. Large trees within river terrace gravels at Khorat (with a tektite) and at Tak in NW Thailand may also be due to the impact.

Other proposed impact effects on probable Miocene fish mortality in Phetchabun, Miocene-Pleistocene mammal mortality in Australia, probably Holocene shell-beds around Bangkok and the Inthanon Uplift in northern Thailand should be considered but are more speculative.

**Key words:** burnt trees, cometary impact, loess, megaflood sediments, Quaternary, Shan-Thai Terrane, tektites, wildfire

### Editorial Preface

The late Dr Sangad Bunopas was the first to relate the Australasian tektite strewn field, the loess covering most of NE Thailand and parts of Laos and Vietnam and the burnt logs and Pleistocene fauna in the Mun River sand deposits of Khorat to an extraterrestrial collision event at about 800 ka. With typical intellectual exuberance, he explained the demise of Miocene fish in Lomsak, Phetchabun, Miocene mammals in Khorat, shell-beds near Bangkok, Miocene and Pliocene marsupials in Australia, epeirogenic uplift in northern Thailand, glaciation in the southern hemisphere and definite glass-producing impacts in Kazakhstan and Tasmania as all caused by a cometary collision. To do this he travelled widely, communicated with experts worldwide and read voraciously in the scientific literature including, palaeontology, sedimentology, geophysics, meteoritics and astronomy. The first list has been partially validated by subsequent work; the second list has not been validated and would be regarded rightly or wrongly as unlikely or very unlikely. His main argument, though, that a major extraterrestrial impact or impacts at about 800 ka had widespread environmental effects throughout SE Asia can hardly be denied and has received considerable support. It remains to be seen whether or not his hypothesis relating the Zhamanshin impact in Kazakhstan to the Indochina impact and to the Tasmanian Darwin Crater impact is correct. A recent paper argues that both the K/T impact and the Zhamanshin Crater were caused by comet impacts thereby providing support for Dr Sangad's hypothesis ((Siraj and Loeb, 2021). His new and controversial ideas have provoked much research and brought together researchers from usually isolated scientific communities. Hopefully this

partial publication of his magnum opus will encourage interest from a younger generation of scientists and provoke considerably more research on tektites, impacts and on the Quaternary sediments and biota of Thailand and beyond. The editor has removed much of Dr Sangad's speculations due to copyright issues but anyone interested in his numerous provocative ideas may obtain a 250 MB copy of his original manuscript from members of the TGJ editorial committee.

Dr Sangad's monumental work was submitted for editing in 2020 but, unfortunately, the manuscript contained too much copyright material which had to be deleted for publication. Professor Kieren Howard (New York) has kindly edited and given permission to include diagrams, photographs and substantial text from his unpublished theses on the Ban Tha Chang sandpits in Nakhon Ratchasima (Khorat) and on the Darwin Crater in Tasmania (Australia). Dr Peter Haines (Perth, Australia) has also generously provided unpublished diagrams and photographs of the Nakhon Ratchasima (Khorat) sandpits studied in Haines et al. (2004). Other deleted figures have been replaced with public domain 'creative commons' figures. The editor has corrected Dr Sangad's English expression where necessary but much of Sangad's inimitable writing style has been retained where his meaning is clear. Where the pronoun 'we' is kept it refers solely to Dr Sangad. Several editorial comments are included in italics, for clarification and in order to bring the manuscript up-to-date.

Dr Sangad introduced several new terms. These are catastroloess which includes all sediments produced as a result of the 0.788 impact event. It includes loess-like sands that cover much of NE Thailand Khorat Plateau and

surrounding areas and also includes sands and gravels in the valleys that cross the plateau. He uses the term DCIM - Destructive and Constructive Impact Materials in order to emphasise the important role of the impact, not only in destruction, but also in supplying building material and fertile farmland throughout the region. Dr Sangad refers to the 0.788 impact event as the Buntharik Event named after Buntharik in Ubon Ratchathani Province following his belief that a crater of that age is present there and that most of the features of the impact features, such as large layered tektites, may also be examined there. The term Buntharik Event is retained until such time as a definite crater or craters can be identified.

For non-Thailand based readers the place Khorat is the short-form name of both a provincial capital city and the largest province in Thailand - Nakhon Ratchasima.. Khorat also gives its name to the extensive Khorat Plateau that encompasses most of NE Thailand and to the Mesozoic Khorat Group sandstones and correlates that may have been the tektite target rocks.

Clive Burrett, Palaeontological Research and Education Centre, Mahasarakham University, Kamriang, Maha Sarakham 44150 Thailand, January 2021. cfburrett@gmail.com

### 1. Introduction

The main purpose of this paper is to suggest that impacts from a fragmenting comet at about 0.78 Ma caused the Australasian tektites and microtektites, the impact craters and their impact glasses from Kazakhstan to Indochina to Tasmania and Antarctica along with widespread catastroloess, fire, mega-floods and mass mortality in NE Thailand and very possibly across much of the region. A cometary impact on earth was difficult for scientists, particularly geoscientists, to believe until the 1994 Shoemaker-Levy 9 orbital break-up into 21 fragments and their subsequent impacts on Jupiter was both predicted and well documented (Figs. 1-3). In this paper it is argued that cometary impacts formed craters, tektites, impactites and catastroloess. Tektites are well known in Thailand but, on the contrary, catastroloess was known only very recently. It has long been known that yellowish, apparently wind-blown sands and silts cover much of northeast Thailand and are very similar in appearance to the yellow-earth of

northern China and Europe which were deposited in arid environments close to Pleistocene ice sheets. The loess ranges from about 3 m thickness around Khon Kaen to less than 1 m in Laos and on the Khorat Plateau, in Laos and in Vietnam overlies a laterite horizon the upper surface of which contains tektites. This loess has a somewhat different grain-size distribution to that of 'normal' Chinese and European loess and extends across the northern half of Thailand.



Fig. 1: Artist's impression of asteroid hitting Earth. Author is Fredrik. Public domain NASA image. The discovery of high pressure/high temperature minerals such as coesite and reidite in Muong Nong (layered tektites) from Thailand and Laos (Cavosie et al., 2018; Glass et al., 2020) indicate Australasian impact temperatures of >1673°C and pressures >30 GPa.




Fig. 2: Impacts of Shoemaker-Levy comet fragments (labelled) on Jupiter . The large mass of Jupiter attracts both comets and asteroids and acts as a sentinel that saves earth from many impact events. Black circle in upper part is the moon Io. From Hubble Space telescope. Open source NASA website.



Fig. 3: Impact of fragments of comet Shoemaker-Levy on Jupiter in 1994. The oval red spot is 16,350 km wide – larger than the earth (12,742 km). Dark spots are explosions several hundred kilometres across. Image from Hubble telescope. NASA open source. <http://hubblesite.org/newscenter/archive/releases/1994/1994/>

In a tropical setting it is considered rather anomalous (Nichol and Nichol, 2015) and overlies fluvial sands along the palaeo-Mun River in Nakhon Ratchasima (Khorat) Province. These fluvial gravels, sands and muds are included here in the new term 'catastroloess' in order to emphasise the catastrophic origin of both sediments. [Ed: It may have been better to keep the genetic 'catastroloess' for the loess-like sediments and the term 'catastrophic fluvial sediments' for the sediments found in the palaeo-valleys, but the term 'catastroloess' is, nevertheless, retained here].

Catastroloess was known from the work of Howard, (1999) and Haines et al. (2004) at Ban Tha Chang and Chum Phuang east Khorat. Later Bunopas et al. (2007) proposed the name catastroloess and is newly studied in Thailand. It was formerly called unmelted ejecta and received little interest. Catastroloess is the most destructive to lives in either breathing and covered to death and also block sun light many years so the climate became cool. The proposed comet entered at a low angle, relatively west of North Pole in N-NW direction travelled southeast, produced craters, induced tektites, impact glasses and catastroloess and other DCIMs, (Destructive and Constructive Impact Materials) with moderate angle from Kazakhstan across Asia and ended at Mt. Darwin, Tasmania, Australia. Microtektites reached the South Pole (Fig. 4).

The comets in every period enter a little west of North Pole (in N-NW direction) leading to South Pole on either the Planet-X Theory or

the Nemesis Theory on low angle route from the impact to the end (could be in the sea or on land). The Planet X (or Planet 9) theory proposes a hypothetical planet beyond the orbit of Pluto that periodically disturbs the asteroidal belts of the solar system and dislodges asteroids or comets towards the inner solar system. Similarly the Nemesis 'Theory' hypothesizes a dwarf star dislodging comets or asteroids towards the inner solar system. [Ed: There has been some support for the existence of Planet 9 recently (e.g. Batygin and Brown, 2016) but neither the Planet X or Nemesis hypotheses are needed to explain a cometary impact on Earth. Dr Sangad has emphasised the importance of Jupiter as protecting earth from cometary impacts due to its large mass and called Jupiter a castle or sentinel protecting our planet from most impacts].

Planet X follows a sharply inclined elliptical orbit that continuously shifts because of gravitational tug of the other planets. As Planet X passes through the disk, it dislodges comets and sends the comets towards earth. Australasian tektite event or Cenozoic's Australasian cometary impact was the most complete journey, when linked those tektite fields together, starting from Kazakhstan, via Indochina (and Thailand) to Australia and the Darwin crater. Comet entered N-NW the continued clockwise S-SW to impact Darwin crater.

In Australasian cometary impact most living die because of dense dusty sands, dusty ejecta or dusty catastroloess, burying or being in the contemporaneous flood, as at Khorat. Thailand.

Cometary impact induced firstly melted ejecta that were tektites, Irghizites and Darwin Glasses, with also equivalent micro-melting. The second is unmelted ejecta that was catastroloess which was different from glacial-related loess or weathering loess.

The destruction could be the following effects of the Cometary Impact:

- Temperature-cooling, leading to frigid conditions for months to years.
- Dust cloud-extensive. Covering surface with choking loess- killing vegetation. No food sources for fauna for months to years.
- Darkness- no photosynthesis, no light for weeks-months.
- Acid rain-poisonous sulphurous fumes for weeks-months. No food sources.

- Wild fires- caused by initial thermal pulse and then by a rain of hot tektites- immediately killing flora and fauna – fires continuing for weeks to months.
- Floods- megafloods- immediate and blocking streams with detritus- killing aquatic life.
- Mass flow and landslide deposits killing trees and blocking rivers.
- Blast wave removes surface vegetation leading to mass faunal mortality.

The story of the Australasian cometary impact primarily began with the work [Ed: studies instigated, inspired and supported by Sangad Bunopas] of Howard, (1999) and later of Haines et al. (2004) on the ancient flood sands on the Mun River's bank at Ban Tha Chang east of Nakhon Ratchasima (or Khorat), Thailand, which were later included by Bunopas et al. (2007) in catastroloess. Catastroloess on land is widespread in Thailand, except in the south, under different local names, and is also newly recognized in SW China. The beginning of the Australasian cometary impact account started here. The comet trail was from the North Pole to Kazakhstan, South China, South-East Asia (Indochina Peninsula) to the Darwin Crater in Tasmania, SE Australia and then to Antarctica (Fig. 4).

The type areas where the cometary impact has been in mind since Barnes and Pitakpaivan (1962) discovered a 12.5 kg. layered tektite, in Ubon Ratchathani Province and donated it to the Department of Mineral Resources (DMR). Tektites are common shiny glasses that look very different from rare meteorites that are dull and rock-like. These must have different in origin and tektites melting condition (Bunopas, 1990, 1992) from a catastrophic event. After exploration on tektites over all upper Thailand (Ford, 1988) and for crater and layered tektites (Wasson, 1991; Wasson et al., 1995; Fiske, Putthapiban and Wasson 1996, and Xu et al., 1989) had an extensive meeting in China, we still could not find any evidence of a cometary collision, until we visited the sand quarry at Ban Tha Chang, east of Nakhon Ratchasima (Khorat) in 1997 (Bunopas, et al., 1999). Other later known localities are at Chum Phuang, Sa-Auong, Bo Phloi, and Kamphaengpet.



Fig. 4: Showing the positions of the Zhamanshin Crater in Kazakhstan, Indochina and Australia. [Ed: The Darwin Crater in western Tasmania is south of the xx'x marked on the map].

## 2. Tektites

### 2.1 Generalizations about tektites

Tektites are small (generally less than 5 cm across, Wasson, 1985), glassy objects that commonly show heat altered zones characteristic of passage through the atmosphere. They appear to be melted ejecta produced by major impacts on to silica-rich sedimentary terrains. The absence of silica-rich lunar rocks rules out a suggested lunar origin for these objects. The explosions apparently were so large that the expanding vapour cloud “blew off” the overlying atmosphere and launched tektites into arching ballistic trajectories and transported them hundreds of kilometers away from the impact site. In 3 cases and possibly 4 cases, the parent crater is known. The tektites consist almost entirely of terrestrial material, though in some case a bit of “meteoric spice” is present.

Tektites, as described by Wasson and Heins (1993) are silica-rich impact glasses (1) devoid of crystallines (2) having very low contents of relict minerals and (3) very low H<sub>2</sub>O contents. The majority of tektites have shapes indicating melt solidification while spinning; such spin form (or splash form) tektites are commonly found at location ten or more crater radii away from suspected source crater, thus long-distance transport is required. Morphological evidence

for atmospheric ablation following solidification is common only tektites found in Australia, though probable ablation features are occasionally found in tektites from Indonesia.

Tektites are found scattered over large areas of the Earth's surface in areas called strewn-fields. The tektites from a given strewn-field are believed to be the result of a single event or tektite shower; and tektites from each strewn-field are characterised by distinctive compositions and age.

Australasian microtektites also extend to Antarctica. This is the youngest of the 4 strewn fields. Part of the Ivory Coast strewn field (1.07 Ma) derived from the Bosumtwi Crater in Ghana is shown. The Central European strewn field (shown as Czechoslovakian) is composed of moldavites (dated as 15.1 Ma) derived from the Ries Crater in Germany. The North American strewn field (not shown) consists of bediasite tektites derived from the Chesapeake Bay Crater dated at 35.5 Ma.

The youngest of these fields, the Australasian strewn-field (Figs. 5,6), has an age of about 0.788 Ma. (Zahringer, 1963, Jourdan et al., 2019). Tektites belonging to the Australasian strewn-field are found in Australia, Indonesia (Borneo, Java and Belitung Island), Malaysia, Vietnam, Cambodia, Laos, Thailand, southern China and the Philippines. Microtektites belonging to the Australasian strewn field have been found in deep-sea sediments from the Indian Ocean, the western Pacific Ocean, and the Philippine Sea (Glass et al., 1979).

Older Ivory Coast tektites have a radiometric age of 1.07 Ma. Tektites from this strewn - field are rare, though Ivory Coast microtektites are sometimes found in the eastern equatorial Atlantic Ocean. Tektites found in Czechoslovakia, called moldavites, are 15.1 Ma old (Gentner et al., 1967) and constitute the Central European Strewn Field. These tektites are distinguished by their transparency and green colour.

The oldest strewn-field is the North American, which has an age of ~ 35.5 Ma (Zahringer, 1963). Tektites belonging to this strewn-field have been found in Texas (bediasites) and Georgia (georgianites). Microtektites belonging to the North American strewn-field have been found in the Gulf of Mexico, the Caribbean Sea, Barbados, and on the continental slope off New Jersey




Fig. 5: Showing the distribution of the Australasian tektite strewn field dated as 788 ka.




Fig. 6: Australasian tektite strewn field. This strewn-field also extends to Antarctica (Folco et al., 2008, 2009, 2010) Only one ablated tektite has been found in oceanic sediments – in the Indian Ocean. Areas with microtektites in DSDP and ODP cores are shown. By synmedia Creative Commons licence CC0- 1.0 Wikipedia.

(Glass and Zwart, 1979; Sanfilippo et al., 1985; Keller et al., 1987; Thein, 1987).

Some kinds of terrestrial impacts for the origin of tektites were suggested since Barnes and Pitakpaivan (1962), Ford (1988), Wasson (1991), Wasson and Heins (1993), Bunopas (1992), McHone et al. (1994), Fiske et al. (1996), Bunopas et al. (1999) in Thailand; Yuan (1981),

Xu et al. (1989) in Hainan, south China; and many others' studies of Australasian tektites of the Australasian tektite strewn field, but Izokh and An (1988) stressed an extraterrestrial impact. Glass (1990), Glass and Pizzutto (1994), and Glass and Wu (1993) introduced an interesting coverage field of these Australasian tektites and made further studies from Asia, Australia and from South China Sea and Indian Ocean deep-sea cores.

Since the first scientific description of an Australasian tektite by Charles Darwin (1844) there has been considerable speculation on their origin. The argument was initially clouded by the fortuitous occurrence of moldavites in Europe and glass at Mt Darwin, Tasmania, along approximately the same great circle, leading David et al. (1927) to propose an extra-terrestrial source to account for this distribution. This led ultimately to the name 'glassy meteorites' for tektites and speculations about a lunar source. Tektites are found in southern Australia, and in the Indian Ocean (Glass, 1990; Glass and Wu, 1993), and microtektites to as far as Antarctica (Folco et al., 2006, 2008, 2009, 2010) The Australasian strewn-field, which is by far the largest identified so far, is about 0.709 Ma old (Gentner et al., 1969) to  $0.770 \pm 0.020$  Ma (in Wasson and Heins, 1993) and is represented by a wide variety of tektite morphologies. The geochemical data indicates that the major components are consistent from China to Antarctica and suggest a common terrestrial source. [Editor's note: Jourdan et al. (2019) have recently provided a reliable 'high precision' date of  $788 \pm 3$  ka for Australasian tektites from Thailand.] Darwin glass from Darwin crater in Tasmania gave an Ar/Ar date of  $816 \pm 7$  ka. A large Muong Nong Tektite (lodged at the Department of Mineral Resources, Bangkok, Thailand) has a diameter of about 0.3 m and weighs 12.5 kilograms.

Tektites are generally small, black, rounded glass objects that superficially resemble obsidian (Fig. 7B). They occur on four widely separated regions of the Earth's surface called strewn fields which range in age from 0.788 to 35 Ma from 4 strewn-fields (Glass, 1990). The presence in tektites of lechatelierite, coesite, baddeleyite and reidite proves a high pressure impact origin.

The Australasian strewn-field contains a radial sequence of tektite shapes ranging from unmodified impactite (layered or Muong Nong type, Fig 7A), through dumbbells and discs (thailandites, indochinites), and spheres (phillipinites, billitonites, and javanites) to ablated button shapes (australites) (Figs. 8-12). This sequence extends from a suspected impact area in Laos or Cambodia or offshore Vietnam following an approximate southeasterly bearing to southeastern Australia and Tasmania. Australasian microtektites are found in a huge area covering much of the Indian ocean and extends across the South China Sea to the Philippines and to Antarctica.

The Australasian strewn-field and the Kazakhstan Zhamanshin impact structure (Florensky and Dabizha, 1980) and tektites to Darwin glasses (Howard, 2004) suggest a cometary impact. In the case of the Australasian Comet, it ended up with the collision of the Earth at the Darwin Crater dated at  $0.816 \pm 7$  Ma.




Fig. 7A: Sites of large Muong Nong or layered tektites in Indochina.



Fig. 7B: Types of tektites. A. Splash-form tektites. B. Small layered tektites. Largest layered tektites known (about 24.5 kg. each). Photographed at the House of Gems in Bangkok, Thailand by Kieren Howard. These specimens are either from Ubon Ratchathani Province or possibly from Laos.

For more than half a century, scientists were mystified by the origin of tektites - were they formed by cometary impact, or meteor or asteroid impact, through terrestrial impact or were they formed by a volcanic explosion on the moon? Recently dozens, of undoubtedly terrestrial impact origin of tektites, and many others DCIMs (Destructive and Constructive Impact Materials) were discovered in Thailand, SE Asia, China, Australia, and many others places, from Kazakhstan to Antarctica. Tektite-like, Irghizites from Kazakhstan, Australasian tektites, Darwin glass (Howard, 2004), were the products on the trail of that comet, ~0.8 Ma ago. Dating of the event is provided by radiometric ages of tektites, real-time correlation is provided by catastrophes.

Layered tektites in NE Thailand (Barnes and Pitakpaivan, 1962, Fudalli and Wasson, 1993, and Wasson et al., 1994, Fiske et al., 1995, 1999), excavated layered tektites at Buntharik in Ubon Ratchathani Province. Tektite strewn fields have been associated with known impact craters, and tektites from one strewn field have been found associated with impact ejecta (here-catastrophes). Some of the tektite events may be associated with other terrestrial events, such as geomagnetic reversals, climatic changes and changes in marine biota (Glass, 1990; Wasson and Heins, 1993), or terrestrial biota (this discovery in Thailand). Tektites defined the date of the impact event and of the impact structure, of the Australasian Cometary Impact, as also other 3 strewn-field tektites.

Irghizites, Australasian tektites (Glass, 1990), and Darwin Glass are silica rich with close dates, their out looked differences could be the same from impact rocks. We link the sites of Irghizites, southern Kazakhstan Zhamanshin impact structure (Florensky and Dabizha, 1980) in Kazakhstan; Indochina tektite field in tektites (Ford, 1988; Glass, 1990; Wasson, 1991, Wasson and Heins, 1993), and the Australasian impact field; to Mt. Darwin glass, now first thoroughly investigated (Howard, 2004).

Microtektites belonging to the Australasian strewn field have been found in deep-sea sediments from the Indian Ocean, the western Pacific Ocean, and the Philippine Sea (Glass et al., 1979). There is now also tentative evidence of Australasian microtektites being found in Chinese Cenozoic loess (Heins et al., 1991; Li et al., 1991). Australasian microtektites have a stratigraphic and formation age of 0.7 Ma (Gentner et al., 1970). Age dating of the Australian tektites by  $^{40}\text{Ar}/^{39}\text{Ar}$  laser fusion gives a date of  $0.770 \pm 0.020$  Ma. (Izett and Obradovich, 1992). There is no evidence of a 0.8-0.9 Ma age for a micro-australite layer in deep-sea cores (Glass, 1976).

Based on the number (mass) of microtektites per unit area at each site, Glass et al. (1979) calculated that there are approximately 100 million, 20 million and 1,000 million tons in the Australasian, Ivory Coast and North American strewn-fields, respectively.

Other glasses that have been associated with tektites include Darwin Glass, irghizites and Libyan Desert Glass. Darwin Glass in Tasmania

is found in a small, radially arranged strewn-field centred on a small crater – Darwin Crater (Fudali and Ford, 1979). Darwin glasses were investigated and studied by Howard (2004), and may have been formed by the Australasian cometary coma that collided on Lower Paleozoic quartzite and limestone in Tasmania. Irghezites are found in the Zhamanshin impact structure in Kazakhstan. They have compositions and ages similar to the Australasian tektites, but many have a composite form consisting of numerous droplets welded together. Libya Desert Glass refers to silica-rich glass bodies found in the Egyptian desert (Week et al., 1984). They have been likened to tektites, but they do not have typical splash shapes and they are composed primarily of silica (>97 %). These glasses have an age of 28 Ma (Storzer and Wagner, 1971). Microtektite-like bodies, some with glass preserved in their interiors, have been found in sediments associated with the K-T boundary and some other impact events (e.g., Izett et al. 1991; Sigurdsson et al., 1991).

Tektites could be easily reworked to a higher level at present surface. This is true from the pits at Khorat sandpits where tektites can be found at any level up to the present surface. With regarding the time of the event, we must count the lowest stratigraphic position of tektite.

For many years the occurrence of tektites dating to about 800-700 ka within much older sedimentary rocks in Australia or apparently unabraded within much younger sedimentary rocks as in Vietnam was known as the tektite age-paradox. The Australian age-paradox was dismissed by the NASA geologist, crater-study pioneer and astronomer co-discoverer of the Shoemaker-Levy Comet using an old fashioned, but effective, plane-table method of surveying critical outcrops (Shoemaker and Uhlherr, 1999). Tektites are easily reworked and great care should be taken in assessing tektites and their abraded surfaces for possible reworking. Downslope reworking and redeposition is more likely in hilly or mountainous terrains and less likely on plateaux such as the Khorat Plateau of northeast Thailand.

Tektites show a number of common features in that they are: (1) devoid of crystallites, (2) have very low content of relict minerals, and (3) very low H<sub>2</sub>O content (Wasson and Heins, 1993). Mineral inclusions range from 20-150 micron and include zircon, chromite, rutile, quartz and

corundum (Glass and Barlow, 1979). Inclusions in Muong Nong-type tektites (indochinites) always show evidence of various degrees of shock metamorphism (e.g. fracturing, droplet formation, x-ray asterism). It appears that indochinites were formed by shock melting of a well sorted, silt-size, sedimentary material (Glass and Barlow, 1979). Trapped gases are similar in composition to the atmosphere or a little lighter (Hennecke et al., 1975).

The best known tektite field, the Australasian strewn-field consists of two kinds of tektites in Thailand:

Layered tektites (Barnes, 1971; Barnes and Pitakpaivan, 1962 and Wasson, 1991), alternatively known as stratified or Muong Nong tektites (Bunopas et al., 1997) have distinctive banding and relicts of incomplete melted parent materials. These tektites were solidified on the ground shortly after impact and were distributed close to the source crater(s). Layered tektites in or near the Hainan-Indochina-Thailand impact center(s) can be up to 12.8 kg (Barnes, 1971) or even 24 kg (Wasson, 1991). In China, Yuan et al. (1981); and Futrell and Wasson (1993) reported the occurrence of a layered tektite weighing 10.8 kg from Wenchang, Hainan Island, south China.

Splash-form tektites (Wasson, 1971) or formed tektites (Bunopas et al., 1997) are found in various forms and shapes, distributed further away from craters. Many splashed tektites have shapes that indicate melt solidification while spinning. Such spinning form (or splashed form) tektites are commonly found at locations ten or more crater radii away from suspected source craters; thereby requiring long distance transport. Aero-ablated forms, common in Australia, were ejected into space and fell down at sites more than 3,000 km from the source, reaching points as distant as Tasmania. Atmospheric ablation following solidification is common only for tektites found in Australia, though probable ablation features are occasionally found in Indonesian tektites. Splashed tektites are usually small, with weights of less than 0.3-0.5 kg.

Both varieties of tektites consist of >70% SiO<sub>2</sub>, 12% Al<sub>2</sub>O<sub>3</sub>, 5% Fe<sub>2</sub>O<sub>3</sub>+FeO, with the remainder constituted by TiO<sub>2</sub>, MnO, CaO, Na<sub>2</sub>O and K<sub>2</sub>O (Glass, 1990; Glass and Koeberl, 1989; Wasson, 1991). A model for the Australasian

tektite field must incorporate explanations for Muong Nong types, discs, dumbbells, spheres, tear-drops and australite buttons. An empirical model for the Australasian tektite field of Ramsay J. Ford (1988) explains this distribution.

The Australasian strewn-field contains a radial sequence of tektite shapes ranging from unmodified impactite (layered or Muong Nong type), through dumbbells and discs (thailandites, indochinites), and spheres (phillipinites, billitonites, and javanites) to ablated button shapes (australites) (Figs. 8-12). This sequence extends from a suspected impact area in Indochina following an approximate southeasterly bearing to southeastern Australia and Tasmania.

“Field work in northeastern Thailand has confirmed that the tektites lie on the surface of a laterite layer below the normal soil profile (Barnes and Pitakpaivan 1962; Fontaine & Workman, 1978; E. Gangadaram pers. comm. 1986). Muong Nong tektites were collected from northeast Thailand near the centres of Ubon Ratchathani and Yasothon. Smaller fragments were found near Chaiyaphum to the west indicating a distribution of fragments of decreasing size in that direction. Muong Nong tektites collected from Hainan Island are also generally small (a few centimetres in diameter). When compared with the large Muong Nong types from the type locality and elsewhere in Laos (Lacroix, 1935), near Dalat in Vietnam (Fontaine, 1966), and near Mukdahan in Thailand (K. Pitakpaivan pers. comm. 1986), there is an apparent radial variation of fragment size of Muong Nong tektites about an area in northeast Cambodia near the Laotian border”.

“In summary, tektites from the mainland of South-East Asia have shapes derived from uncongealed spinning glassy fragments passing through the atmosphere. An event, comparable with that responsible for the South-East Asian strewn-field, must have a characteristic set of fragment trajectories. In the first 400 km from impact there are large Muong Nong tektites; between 400 and 1000 km there are disc and dumbbell shapes (e.g. thailandites, indochinites); between 1000 and 3000 km, spherical and tear-drop shaped phillipinites, billitonites and javanites occur; and at distances greater than 3000 km ablated tektites (australites) would be most abundant. The button form of australites (Fig.8) is unique to the Australian strewn-field

and is produced by travelling more than 3000 km from the source”. Ford (1988)

## 2.2 Impact Model for Tektite Origin

The presence in tektites of lechatelierite, coesite, baddeleyite, and shocked relict crystal fragments suggests formation in an impact event. A multiple impact origin for Australasian tektites was documented by intensive laboratory work and field observation by Wasson (1991). High-precision neutron-activation studies of Southeast Asian tektites confirm their close compositional relationship to well-mixed continental sediment. In confirmation of early studies, concentrations of the volatile metals Zn, Ga, As and Sb are 2-4 times lower in splashed-form tektites than in layered (or Muong Nong-type) tektites. The Th:U ratio is higher in splash-form than in layered tektites; which implies a loss of about 30 % of initial U from splashed-form tektites. Volatile contents of layered tektites are



Fig. 8: Aero-ablated australite button from Australia. Note typical ablated flange. These buttons usually range from 1 to 2 cm in diameter. From Wikipedia article on tektites. H.Raab (user: Vesta) Creative Commons licence : CC BY. SA.3.0



Fig. 9: Dumbbell (c.10 cm long) and teardrop splash-form tektites from Thailand. Photograph by Broken Inaglory, under Creative Commons licence CC BY-SA 3-0. Wikipedia article on tektites.

similar to those in continental sediments indicating minor or negligible loss during formation. A remarkably consistent relationship between Cr and Co and an associated Ni-Co correlation indicates the variable presence of a minor ultramafic component with an atypically low Cr/Co ratio.

“The chief compositional difference between SE Asian layered and splash tektites is that the former have higher contents of volatile elements (Taylor and McLennan, 1979; Shaw and Wasserburg, 1982) and, possibly, higher  $Fe^{3+}/Fe^{2+}$  ratios (Fudali et al., 1987). The two classes are remarkably similar in their contents of non-volatile incompatible elements, chondrite-normalized rare-earth elements (REE) patterns for layered tektites from Laos, Thailand and Vietnam as well as for splash tektites from Australia reveal patterns that are essentially identical for layered and splash tektites”.

“The coherence of the trace-element patterns is further demonstrated by comparing key ratios between incompatible elements: Cl-normalized La/Lu, Eu/Sm, Zr/Hf, and Th/Sc ratios are plotted against Th/U in Wasson (1991). To help resolve the points, splash tektites are shown with dot symbols, layered tektites with open symbols. The difference in Th/U between layered and splash tektites was unexpected; average ratios in the splash tektites are 30% higher. This observation almost certainly indicates that a significant fraction of U was lost as a volatile from the smaller splash tektites. The Eu/Sm, Zr/Hf and Th/Sc ratios are the same in layered and splash tektites: the mean La/Lu ratio in splash tektites is about 2-3% higher than that in the layered tektites, but this difference may not be significant.” Wasson 1991.

Wasson and Heins (1993) found that tektites are derived from the impact melting of continental sediments, and based this proposal on close similarities of elemental abundance patterns (Taylor and McLennan, 1979; Koeberl, 1986; 1992; Wasson, 1991) as well as similarities in isotopic ratio (Shaw and Wasserburg, 1982; Blum et al., 1992).

Fiske et al. (1996) analysed shape, composition and spatial distribution for 6 kg of layered tektite fragments excavated from a 9 m<sup>2</sup> area near Ban Huai Sai, north of Amphoe Buntharik in northeast Thailand. Their analysis suggests that the fragments represent a single homogeneous

mass that underwent ancient fragmentation and has experienced little disturbance since its deposition. They have also studied the stratigraphic occurrence of layered tektites exposed in-situ near Ban Huai Om. Tektites were found along a disconformable paleo-erosion surface, covered by recent aeolian sand (partly reworked catastroloess) and in a manner similar to their occurrence elsewhere in Southeast Asia. This stratigraphic relationship provides little chronostratigraphic information and does not constitute a stratigraphic ‘age paradox’ for Australasian tektites. The present-day surface density of layered tektites in this area is 2 to 20 g/m<sup>2</sup>.

Iridium anomaly associated with the Australasian tektite-producing impact from masses regarded as the impactor of the Australasian tektites was restricted due to at least three orders of magnitude smaller than that impactor at the end of Cretaceous (Schmidt et al., 1993).

Basement rocks in Thailand prior to melting had originally been continental sandstones from the Jurassic-Cretaceous continental red-beds (Khorat Group). Thin decomposed sandy soil has also been identified in the region, a finding confirmed by our latest field investigation (Bunopas et al., 1997). Blum et al. (1992) found that Nd and Sr isotope data provide evidence that all Australasian tektites were derived from single sedimentary formation with an age close to 170 Ma. Moreover Blum et al. (1992) propose that such tektites are explained by a single impact event. Tektites are constituted by molten terrestrial material and in the impact process they undergo no geochemical change. The absence of geochemical disparity between tektites and their terrestrial source has been attested to by chemical analyses, rare earth studies, mineral fluid inclusions and trace elements (McLennan, 1989, Schwarcz, 1962; Glass, 1970, 1972; Glass and Barlow, 1979; Glass and Koeberl, 1989; Koeberl, 1986; Muller and Genter, 1973; Taylor, 1962; Taylor and McLennan, 1979; Shaw and Wasserberg, 1982; Wasson, 1991; Blum et al., 1992). A possible offshore impact site has been suggested (Schnetzler et al. 1988) with Hartung, (1990) nominating Tonle Sap for site candidacy. More recently a site within the Bolaven Plateau of Laos has been suggested (Sieh et al., 2020).

### 2.3 Contending Views of Tektites

Although most investigators believe that tektites are impact-produced glasses, their origin was controversial (e.g., Taylor, 1973; O'Keefe, 1976; Wasson, 1991). During the 1970's and 80's, the idea of extraterrestrial origin for the Australasian tektite strewn field was popular (Izokh and Le Dyk An, 1988). It was indicated by the dimensions of the belt itself, the lack of impact craters, an absence of known impactites, and little indication of concentric zonation relative to possible terrestrial source (O'Keefe, 1976). However, the overall chemistry of tektites is terrestrial, bubbles in tektites contain atmospheric gases and layered tektites contain silt-size clasts derived from a sedimentary source.

The number of associated impact craters is a hotly debated topic. Two models for the origin of Australasian tektites compete in recent research.

Ford (1988) used satellite imagery to define an elliptical depression 10km x 6km in Phnum Voeene, northeast Cambodia, a formation claimed to be caused by the impact of an extraterrestrial object. This single-crater hypothesis has been endorsed by Blum et al. (1992), Koeberl (1992) and Schnetzler (1992). Based on the study of trace elements in Muong Nong-type tektites from Indochina, Koeberl (1989) interpreted a single, slightly heterogeneous source, rather than different source regions.

However, Wasson (1991) argued that large layered tektites (with thickness of 2-3 cm and masses up to 24 kg), found throughout northeast Thailand and other Indochinese regions, implied a multicrater centre with minimum dimensions of 800 x 1124 km, extending from Hainan Island Vietnam (Izokh and An, 1988) and Cambodia. This represents a major centre containing numerous impact craters with a collective size thousands of times larger than that estimated by Ford (1988). Wasson did not overlook the possibility of a fragmented extraterrestrial object or a large comet as being responsible for forming such an extra-large cratering centre. The largest impacts ejected the microtektites (Yuan, 1981; Futrell and Wasson, 1993) to southern regions and some splashed from tektites above the atmosphere, while numerous smaller impacts produced a nearly continuous sheet of impact melt across much of Southeast

Asia (Wasson, 1991; Wasson and Heins, 1993; Wasson et al., 1995).

There is compelling evidence indicating that layered tektites formed as sheets or pools of melt. From a lens-shaped mass of tektites excavated near Phang Daeng, it was concluded that layered tektites represented 'puddles' of melted soil resultant from the impact of a large 'diffused body' and the consequent fireball (Barnes, 1989). Because there are difficulties with the impact transport of melt at distances >600 km, tektite distribution across a field >1140 km in length is inconsistent with their formation in a single crater. Layered tektites should therefore be deposited within a few crater radii of their parent crater, and many craters are required to account for their distribution across a field >1140 km long.

### 2.4 Investigation of Tektites in Thailand and Southeast Asia

It is 60 years since investigation of tektites began in Thailand (Barnes, 1963, 1964; Barnes and Pitakpaivan, 1962) that included the discovery of the largest known documented tektite (Barnes, 1971). Early research on tektites in Thailand progressed slowly.

Preliminary collection of tektites was undertaken by the Geological Survey Division, Department of Mineral Resources (1960) and Barnes and Pitakpaivan (1962). More recent collections, investigative of the nature and distribution of tektites, were assembled by Ford (1988) and Bunopas (1990, 1992). Tektites in Ford's study (1988) were found in both the upper lateritic horizon as well as at the modern lateritic surface in situations where it had not been disturbed by agriculture. Ford (1988) developed an earlier hypothesis of Adam and Huffaker (1964) and proposed that tektites originated in an impact site in Cambodia, with dispersal occurring as far as Tasmania. Contributions by Wasson et al. (Wasson, 1987, 1989, 1991; Wasson and Heins, 1993; McHone et al., 1994; Fiske et al., 1996; Bunopas, Kositanont and Wasson, 1997; Bunopas et al., 1998) have also excited interest in the scientific community.

Apart from Laos (Lacroix, 1935), there has been some development of this research in other Indochinese countries. In Vietnam, Fontaine (1966) made note of medium Muong



Fig. 10: Department of Mineral Resources 12.8 kg - largest described layered tektite from Ubon Ratchathani Province, Thailand. It was contributed by Mr. Kaset Pitakpaivan. From Buriram to Ubon Ratchathani and approximately from Mukdahan to Nakhon Phanom all tektites are essentially layered tektites (Muong Nong) tektites, fragmented to small size. We can differentiate fragmented layered tektite, usually with experience, from splashed tektite by air bubbles and skin of the latter.

Nong tektites near Ho Chi Minh City, though there was little analysis of the discovery. In a more recent comprehensive study of Vietnamese tektites, Izokh and Le Dyk An (1983, 1988) explored 39 localities throughout Vietnam and subdivided the area into various sub-parallel belts. Tektites were found at variable levels of Quaternary deposits consisting of loess, conglomeratic sands, and laterites; though in a few instances tektites occurred at the bottom of the basement covered by loess and burnt wood/charcoal. They proposed that their tektites represented a sample from an Australo-Asiatic zone extending as a narrow 11,000 km belt from Vietnam to Tasmania. The Australo-Asiatic Belt was characterised as showing zonation in tektite size, morphology, as well as in certain aspects of fine internal structure. Izokh and Le Dyk An (1983, 1988) concluded that the belt was the result of a single tektite shower of extraterrestrial origin. Such a view is however inconsistent with views of others (e.g. Fiske et al., 1996) in which the terrestrial impact of a meteor or comet was believed responsible. Other significant papers on Cambodian tektites include Hildebrand et al. (1994); Hartung et al. (1994); Hartung (1990); Hartung and Rivolo (1979); and Ford (1988). This paper also deals primarily with the latest investigation of tektites and associated phenomena in the Khorat Plateau and elsewhere in Thailand. Research was conducted with the participation of the University of Tasmania's

Geology Department in 1984 – 2004 (Ford, 1988; Howard, 1999, 2011; Howard et al., 2003; Haines et al., 2004) and with UCLA (University of California, Los Angeles) in 1989, 1990 and 1994 (Wasson, 1987, 1989, 1991; Wasson and Heins, 1993; Fiske et al., 1996).

*2.5 Layered tektites had a limited concentration in NE Thailand and in Laos, Hainan Island, south China and Vietnam - Layered Tektites formed in pools of melt*

Layered tektites (Figs. 11-12) have outstanding fine to very fine layering characteristics, some wavy or contorted since they solidified in situ, or were considered so, or remained nearby, or at a short distant away. The sizes range from a half kg to as heavy as 24 kg from Buntharik in Ubon Ratchathani (Fiske et al., 1996; Bunopas et al., 1999a). Ubon Ratchathani is at a similar latitude between 15°N-16°N comparing with Muong Nong in Laos and Danang in Vietnam, all are rich in layered tektites. The largest layered tektite of more than 24 kg was assumed to come from either Ubon Ratchathani Province, NE Thailand or from Laos (Fig. 11).

“Layered tektites first described by Lacroix (1935) as “tektites sang formes figurées” i.e. as unshaped tektites. The first of these were found near the village of Muong Nong in the Laotian highlands, and Barnes (1963) used the designations “layered tektites” and “Muong-Nong -type tektite” synonymously. The descriptor “layered” derives from the presence of subparallel layers defined by colour and/or bubble contents (Barnes and Pitakpaivan, 1962). Layered and splash-form tektites have very similar chemical compositions. The traditional view is that each of the tektite fields was formed as ejecta from a single crater. In fact, it appears impossible that ejecta from a single crater could form most pools stretching across an area with linear dimensions of 1200 km from Kampong Speu, Cambodia (Barnes and Pitakpaivan, 1962) to Hainan, China. There are several major problems associated with the deposition of melt at great distances: (1) the melt would be dispersed into small masses by atmospheric shear, and these would be unable to penetrate large distances through the atmosphere; (2) during the long flight time globules of melt would lose much of their heat and solidify; (3) if (by some miracle) the atmosphere were temporarily removed’ from the entire 1200-km region such that large,



Fig. 11: Two c. 24 kg layered tektites at House of Gems in Bangkok courtesy of Boonmun Poonyathiro (photo by Howard, 1999). They are either from Ubon Ratchathani Province or from Laos.

m-size globules could complete their trajectory intact, their impact velocities would be so great that they would make hypervelocity craters and be dispersively mixed with unmelted ejecta”.

“Our study of the stratigraphic occurrence of layered tektites exposed in situ near the Huai Om reservoir shows that tektites are found along a disconformable paleo-erosion surface covered by recent aeolian sand, similar to other occurrences throughout Southeast Asia. Based on our study at Huai Om the present-day surface density of layered tektites < 10 g in this area is 2 g/m<sup>2</sup> to 20 g/m<sup>2</sup> Our study of the stratigraphic occurrence of layered tektites exposed in situ near the Huai Om reservoir shows that tektites are found along a disconformable paleo-erosion surface covered by recent aeolian sand, similar to other occurrences throughout South-east Asia. Based on our study at Huai Om the present-day surface density of layered tektites < 10 g in this area is 2 g/m<sup>2</sup> to 20 g/m<sup>2</sup>.”

“Layered tektites formed as pools of SiO<sub>2</sub>-rich melt on the Earth’s surface. In Southeast Asia layered tektites having thicknesses of 3-20 cm are found throughout a region 800 x 1200 km in extent. Pools of impact melt cannot be deposited at great distances (>100 km) from a crater, instead, they are probably confined to the area (extending 2-3 crater radii) covered by continuous ejecta. The layered tektites of Southeast Asia probably formed in a myriad of small craters produced by fragments from a comet that disintegrated several months before accretion” (Wasson, 1991).




Fig. 12: Layered Tektite from Danang 0.5 kg from H. Fontaine, 1966.

“Glass (1970) described silt-size and size sorted zircon and chromite crystals, in addition to quartz and possibly coesite, in a Muong Nong-type indochinite. Glass (1970; 1972) believed that most of the mineral grains were relict detrital grains and that the mineral assemblage suggested a sedimentary parent material. Glass also pointed out that the mineral grains all showed evidence of shock damage

consistent with the presence of coesite as reported by Walter (1965)” [Ed: and Glass et al. (2020)].

“The presence of lechatelierite indicates a high temperature of formation for tektites (> 1700 °C) and the presence of cristobalite associated with quartz, could be due to conversion of coesite and/or stishovite due to high residual temperatures”. Data support a high temperature of formation for the Muong Nong-type Australasian tektites and that the minimum temperature must have been in excess of about 1,850° C.”

Punpate et al. (2005) collected five samples of bedded or layered tektite and tektite-bearing sediments from an excavated trench and were selected for thermoluminescence dating from the Buntharik area, Ubon Ratchathani in the easternmost part of the Khorat Plateau, northeast Thailand. “Five sedimentary units of semi- to un-consolidated alluvial deposits were described stratigraphically in a descending order (Fig. 13). Unit 1 (av. 0.14-m thick) is characterized by light yellowish orange, poorly compacted, sandy loam (catastroloess) with abundant roots. Unit 2 (av. 0.1-m thick) is dark brown, moderately compacted, clayey sand, with well-sorted, and well-rounded sand. Unit 3

(~av. 0.38-m thick) comprises mainly light yellowish brown, poorly- compacted, medium to fine sand, chiefly with well-sorted and well-rounded sand. Unit 4 (av. 0.3-m thick) is composed mainly of pale brown, clayey to sandy compacted gravel with moderately sorted, sub-rounded to rounded coarse-very coarse sand and bright reddish brown cloudy mottles. This unit contains sparse embedded tektites and iron concretions, the latter has variable diameters ranging from 2 to 10 mm. Unit 5 (av. 0.5-m thick) is defined by light gray, moderately compacted sandy clay, bright reddish brown cloudy mottles. This unit contains sparsely distributed iron concretions with diameters of 1-3 mm. All the tektites were observed only in Unit 4 and are regarded as “layered “tektite. The tektites ranging in size from 1 cm up to 5 cm (Fig. 14) were discovered in a pebble bed at about 0.7 meter from a topsoil surface”.

“Average TL age obtained from unit 3 is about 27.5 ka, that of Unit 4 is about 143 ka, and that of Unit 5 is about 370 ka. The average TL age of the layered tektites from Unit 4 is about 850 ka. The discrepancy in TL ages of tektites and tektite-bearing sediments are ascribed to the reworking of layered tektites from high




Fig. 13: Detailed stratigraphy of the semi to unconsolidated alluvial deposits in Buntharik area, T1, T2, and T3 are sampling points of tektite, and Q1 to Q6 are tektite bearing sediments for thermoluminescence dating (From Punpate et al., 2005)




Fig. 14: Various bedded tektite varying from 1 to 5 cm from Unit 4. (From Punpate et al., 2005).

elevation to the low land and low alluvial plain. Additionally the TL ages of tektites from this study conform quite well with earlier TL ages of splashed tektites at the Ban Tha Chang sandpit in Nakhon Ratchasima, located 400 km in the western most part of the Khorat Plateau, suggesting a widespread occurrence of approximate 0.8 Ma catastrophic event (Charusiri et al, 2002). We consider that such the event may have taken place in response to the meteorite impact in Buntharik area over the clastic beds of the Khorat Group as earlier described by Bunopas et al. (2002).” Punpate et al. (2005).

“There seems to be only one plausible explanation of the layering of Muong Nong tektites, viz., that it was formed by small amounts of differential (probably down-slope) flow in a pool or sheet of melt (Barnes and Pitakpaivan, 1962). Key supporting evidence is provided by the magnetic remanence studies of De Gasparis et al. (1975). The inclinations of the measured paleofields are inclined by about 20° relative to the layering, roughly the same as the present-day inclination of the field in Southeast Asia: since the inclination was probably similar 780 ka ago, this observation strongly supports the interpretation that the layering was produced by processes occurring after the melt was deposited on the Earth’s surface. It seems irrefutable that the layered tektites formed as pools or sheets of melt.” Wasson (1991).

“Although the layered tektites seem to require formation as melt ejected at low velocity from a multitude of local craters, the microtektites and the ablated Australian tektites were almost certainly launched into trajectories that took them above the atmosphere. The

evidence in support of this conclusion is the ablation features found on many australites and the great distance that microtektites have traveled; according to Glass (1982) the microtektite field extends from Madagascar to Taiwan, a distance > 6000 km. Because a 100-gm. microtektite will fall out of the stratosphere in days but large-scale lateral stratospheric transport requires months or years (except eastward in the jet stream), the observed wide distribution can only be achieved by ballistic transport above the atmosphere.” Wasson (1991).

#### 2.6 Multiple small (0.2-2 km) crater model of Wasson

“An important and possibly telling observation is that investigations of the tektite-bearing laterite layer in numerous locations in Ubon Ratchathani yielded many fragments of layered but no splash tektites, in contrast, most tektites found near Khon Kaen, 250 km to the northwest, are splash form. This observation is consistent with formation of craters near the Ubon Ratchathani area, and deposition of melt sheets over this region. Splash tektites are not expected under such conditions because the shapes of splash tektites that fell into the melt would have been degraded. Discovery of additional regions covered by melt >100 km from Ubon Ratchathani would confirm the necessity of multiple craters.”

“If this constraint on the distance that melt can be transported and collected into pools is correct, a multitude of craters across Southeast Asia are required to explain the layered tektites. Most of these craters must have been small perhaps 0.2-2 km in radius roughly corresponding to 20-200 m in projectile radius. During most 1-Ma time intervals the largest object accreted to the Earth has a radius of 500 m, and a mass of about 1,012 kg (Wetherill and Shoemaker, 1982, KYTE and Wasson, 1986). The formation of many craters spanning a region > 1000 km long requires that such an interplanetary object broke up at great distances from the Earth and rained down as 20-200 m projectiles. Melt ejected from the resulting craters fell out locally and formed the layered tektites. Some melt escaped these fireballs and solidified in flight to produce the splash tektites.” Wasson, (1991).

“Tektites of the early Quaternary are only distributed in the extreme south of mainland

China: the Leizhou Peninsula, the coastal area of Guangxi and Hainan Island. Xu Daoyi et al. (1989) included all of them to the Australasian tektite strewn-field. Original tektites can only be recognized in the section of (Q1-Q2) with continuous sedimentation of lacustrine facies, which can be easily traced in South China. Tektites have been discovered between the Zhanjiang Formation of the Lower Pleistocene and the Beihai Formation of the Middle Pleistocene. These two formations, whose ages may be determined by fossils, are widespread in Hainan Island and the southern part of Guangdong Province. They are characterized by mostly continental facies with interbedded layers of marine strata of various extent. The tektites occur precisely at the boundary between (Q1-Q2) which has an isotopic age of approximately 0.7 Ma; this coincides with the transition from the older Matuyama Magnetic Reversed Chron to the younger Brunhes Magnetic Normal Chron. The distribution of tektites is concentrated at the upper surface of the stratigraphic unit of the Zhanjiang Formation and the autochthonous tektites which were discovered in the boundary section at Wenchang Middle School, Hainan Island, have never been found at any other stratigraphic position" (Xu DaoYi et al., 1989).

More destructive and constructive impact materials (DCIMs) on terraces and in old rivers, were the environmental effects of the impact: The newly discovery of the extinction evidences of abundant buried, ancient felled trees (Xu et al., 1989) large tree trunks, numerously burnt or/and petrified plant material, burying under subsequently widespread mudflows, landslides, earth flows and avalanches with embedded tektites found on these previous terraces stratigraphically covered with the newly recognized 20 - 40 m thick catastroloess in Khorat in ancient river flooded deposits at Tha Chang, 300 km west of Ubon Ratchathani.

### *2.7 Tektites in Vietnam, The Problem of the Ground Transport and Scattering of Tektites in Vietnam*

We know tektites from Vietnam from Izokh and An (1988) expedition for position of them mainly for a Quaternary stratigraphy but rather limited as the following outlines, and modern concept on tektites.

This problem seems to be an artificial one for Izokh and An (1988), since it arose exclusively

because the investigators who concerned themselves with tektites tried to solve the age paradox that arose by assuming that the tektites were eroded out of the more ancient deposits into the subsequently earlier ones along the surface of the Earth. The authors of this article suggest taking this problem away from serious discussion because there are no factual bases for its establishment. More than 30 test pits in all Vietnam by Izokh and An, (1988) were proved to be similarly in Thailand with tektites immediately above the basement, or thought to be an old surface. In many pits to more than 10 m depth each contained also frequent burnt woods and charcoals. Dating of charcoals and dating by fission tracts of tektites from these pits and obtained many absolute ages confused the investigators. The age of charcoal that could be obtained was only less than 1.3 % of a maximum age of associated tektite in the same pit in similar catastroloess. There is no sense in searching for them in regions where the relief is strongly broken and where the remnants of the terraces or the ancient flat surfaces have not been preserved. This means that the tektites essentially vanish in the mass of redeposited material of the bedrocks. And they only remain in the "erosion-proofed" situation--on relatively flat leveling surfaces under a layer of loess. They remain in many practically the same place where they fell to the Earth. Thus, there also can be no discussion about the erosion of tektites from many other (later) source, or--still more about the strict selective (without the accompanying by fragments of other rocks) transport and scattering over extensive areas.

The tektites of Vietnam in most cases are devoid of signs of redeposition or transport, i.e., lie where they fell-in situ (Izokh and An, 1983). They are non-uniformly distributed in the area in the form of isolated areas or bands and are found in any planes of leveling, beginning with the 10-15 in flood terraces and ending with the highest levels (over 2.5 km). They are absent on the surface of the contemporary floodplain and on the coastal plain. This means that the tektites fell to the Earth at a time when the level of today's terraces above the floodplain, i.e., when they were the basis erosion surface. Nevertheless, they earlier concluded that the tektites fell to the Earth very recently (Izokh and An, 1983), which are now known for one age ~0.8 Ma (Glass, 1989, 1991; Wasson, 1991,

Ford, 1988, Bunopas et al., 1999; Howard, 2000). The age paradox (Izokh and An, 1983) is now no more existed. Carbon dating of coal and charcoal, associated with tektites, is only a partial age (or a minimal age) because C dating has its limitation does not exceed fifty thousand years.

Izokh and An (1988) give the characteristics of the most typical sections where tektites have been found by the authors directly in the diggings. Where it is not stated otherwise, the descriptions belong to E. P. Izokh. Many loess mentioned by them should read catastroloess, similarly but are more widespread and are more completely studied, in Thailand as catastroloess (not the wind-blown sands).

“Open pit mines 18 km north of Ho Chi Minh City near the highway running in the direction of Ben Hua, 10 separate fragmented tektites of the Muong-Nong type were found at the base of the loess layer. There were the same number of fragments of droplets and spheres, as well as one intact disc. They were all devoid of even traces of roundness. Fine details of the primary structure as well as sharp edges of fragments were preserved. Bits of woody coal which were particularly copious near the foot were found in the loess throughout the entire section of the layer.”

“Here pocket accumulations up to 10-15 cm in diameter consisting of separated bits of coal ranging in size up to 3-4 cm which preserve the structure of the ancient tubes are found. The coal undoubtedly lies in situ, and in this regard the coal, loess, and tektites are equivalent. The presence of the coal verifies namely the subaerial nature of the examined layer, i.e., the absence of subsequent washing out and redeposition”.

“Tektites were collected by Izokh and the geomorphologist, Le Duc An from more than 120 places all over the territory of Vietnam. Tektites occurred in wide belts like those in Australia (Izokh and Le Duc An, 1983). Tektites are found on various planes of denudation, from 10-15 - meter terraces to 2-2.5 km high highland plateaus. There are no tektites on the flood-plain level, on slopes and in the recent alluvium. That means that tektites fell at the time when 10-15 meter terraces served as flood plains. i.e. the lowest (basic) level of erosion. Tektites belong to the one sole stratigraphic level which can be

found throughout the Vietnam territory. It is at the bottom of the sandy loams varying in composition from place to place from clay-bearing to more sand-bearing, but ever having loess-like appearance. They are apparently subaerial sediments covering like a blanket the unevenness of the relief”.

“The underlying formations are distinguished by a great variety of age and composition. The youngest of them are alluvial pebbles of 10-15 metres high terraces dated as the nearly Pleistocene-Holocene boundary. Practically all bedrocks are lateritized, often completely. However, the tektites and loess-like layers are not affected by the process of lateritization, which provides one more evidence of their recent geological age. The loess-like layer contains abundant charcoal inclusions. The earliest <sup>14</sup>C-age (10,210 and 8,670 y.) was determined for the charcoals from the bottom of a loess-layer. The charcoals of different levels of loess-like sediments are of various ages: 3,855-1,840, 2,130, 1,435, 1,205, 1,060 and 385 years (analyst A. Panychev, Novosibirsk). It should be noted that the loess-layer is the uppermost stratum of the stratigraphic sequence of Vietnam, usually 0.2-1.5 m thick, but not more than 2- 3 m. In fact it is the main soil-forming layer, cultivated by man; charcoal's often associated with old pottery fragments. Among the Mekong-delta sediments, in the alluvial pebbles of the Saigon river, a geologist found some debris of trees dated at 27 + 0.5 thousand years at a depth of 16 m, and some others 3± 2 ka. at a depth of 63 m. Much earlier tektites were found in the same pebbles by Saurin (1935) at a depth of 40 m, i.e. on a stratigraphically higher level” Izokh and An (1988).

The valuable investigation on the position of tektites of Izokh and An (1988) in Vietnam reflect the consistent fact found in Hainan Island (Zu Dao Yi et al., 1989) in the north and recently in northeast Thailand, in the south-west (Wasson, 1995; Fiske, 1995; Bunopas et al., 1999, 2001, 2002, 2023, 2004) on the nearly NS trend of the Indochina Peninsula (Wasson, 1991; Wasson and Heins, 1993).

In Vietnam, tektites (in situ or intact tektites) are at bottom of the loess (in fact-catastroloess) on bedrock or gravel beds, as were found by Izokh and An (1988) in Vietnam in almost every

test - pits from south to north, and in Hainan (Xu Dao-Yi et al., 1989). The handicap of dating of burned wood getting ages never beyond 50,000 years which has been limited by the method. Thus it was impossible for using the age for tektites. By this concept the paradoxical age of tektites is misleading.

### *2.8 Ablated tektite in the Indian Ocean*

“A well-preserved ablated (button-shaped) tektite recovered from the surface sediments of the central Indian Ocean (Glass et al., 1996) lacks flow ridges and has apparently undergone ablation of 6.9 to 7.9 mm. The lack of flow ridges and amount of ablation indicate that, if it originated in Southeast Asia, it must have had a very shallow trajectory (only a few degrees) and a velocity on the order of 7 km/s as it re-entered the atmosphere.

“Prasad and Rao (1990) reported the discovery of a well-preserved ablated (button-shaped) tektite from the surface sediments of the central Indian Ocean (12°37’S; 78°30’E) in a water depth of 5,300 m. An ablated (button-shaped) tektite recovered from the central Indian Ocean basin is a unique specimen. It is the first tektite to be recovered from a deep ocean basin and it is the only ablated tektite recovered from the ocean floor. It is better preserved than tektite specimens found on land. There is a large (2.2 mm diameter) high-velocity impact pit on the anterior surface. The specimen lacks flow ridges on the anterior surface and appears to have undergone ablation in the range (1.9 to 7.9 mm as it came down through the Earth’s atmosphere” Glass et al. (1996).

### *2.9 Microtektites coesite, stishovite and shocked quartz in ocean-core sediments*

“Coesite and shocked quartz were found in seven out of 33 cores from the Australasian strewn field; however, no coesite or shocked quartz was found associated with the Ivory Coast microtektite layer (Glass and Wu, 1993). Stishovite was also found at three of the Australasian sites. All of the Australasian cores containing coesite and shocked quartz were taken from within 2000 km of Indochina, which is believed by most to be the source area for the Australasian tektites and microtektites. Six of the seven cores have the highest concentrations of microtektites ever determined for Australasian microtektite-bearing cores. Australasian

microtektites have been found in more than 40 cores throughout much of the Indian Ocean, the western equatorial Pacific, and the Sulu, Celebes, and Philippine seas: a total area covering ~10 % of Earth’s surface (Glass, 1990; Smit et al., 1991; Schneider et al., 1992). Shocked quartz and coesite (>125 µm size fraction) were found in seven of the 33 cores from the Australasian strewn field. All seven cores were taken within 2000 km of Indochina, and six of the cores have the highest concentration of microtektites (>125 µm in diameter) of all the Australasian microtektite bearing cores (>300/cm<sup>2</sup>). (i.e., close to the Brunhes-Matuyama reversal boundary). In three cores the shocked quartz and coesite were restricted to the microtektite-bearing layer, and their peak abundance occurred within 10cm of the peak abundance of the microtektites”.

“The presence of coesite and shocked quartz in Australasian microtektite-bearing cores closest to the Indochina area and having the highest concentration of microtektites is consistent with previous suggestions that the source crater for the Australasian tektites and microtektites is in the Indochina region. The coesite and shocked quartz found near the Brunhes-Matuyama reversal boundary in a core that did not have an Australasian microtektite layer and taken from within the Australasian strewn field indicate that the distribution of microtektites within the strewn field is patchy and may represent a ray-like pattern”. (Glass et al., 1996).

### *2.10 Iridium anomaly associated with the Australasian microtektite layer in ocean cores, the mass of the impacting body, the position and size of the Australasian crater and Antarctic microtektites*

In their introduction (Schmidt, Zhou and Wasson, 1993), of their paper concerning iridium anomaly of the Australasian tektite, essentially all Cretaceous / Tertiary (K/T) boundary sediments (deposited 65 Ma ago) show anomalously high Ir concentrations that are widely accepted to constitute evidence of a major impact event (Alvarez et al., 1980; Kyte et al., 1980) at that time.

“Data on deep-sea cores show that the microtektite fluence falls exponentially away from southeast Asia, the fluence dropping a factor of 2 in ~400 km. In southeast Asia the trend merges with a roughly estimated mass

fluence of  $\sim 1.1 \text{ g cm}^{-2}$  inferred from evidence of a melt sheet in northeast Thailand. Integration of the inferred distribution yields a total mass of Australasian tektites of  $3.2 \times 10^{16} \text{ g}$ , much higher than previous estimates. Assuming a similar fallout distribution for the impactor and a chondritic composition allows us to calculate its mass to be  $1.5 \times 10^{15} \text{ g}$ , about 3 orders of magnitude smaller than the minimum mass of the impactor responsible for the extinctions at the end of the Cretaceous. The fluence of Ir can be used to estimate the mass of the projectile. This approach yielded a mass of  $\sim 10^{18} \text{ g}$  for the end-Cretaceous projectile (Kyte and Wasson, 1982) and  $\sim 10^{14} \text{ g}$  for the late-Pliocene impact in the southern Pacific (Kyte et al., 1988). To estimate the dependence of fluence on distance from southeast Asia we calculated the great-circle distances from Buntharik, Ubon Ratchathani Province, Thailand, near the center of a region where abundant layered tektites have been recovered in the southeastern corner of northeastern Thailand.”

“Our integration of the total mass of tektitic material deposited across the field is  $3.2 \times 10^{16} \text{ g}$ , 320 x greater than the Glass et al. (1979) value of  $10^{14} \text{ g}$ . Geochemical analyses of sediments from ODP site 758B and from ODP site 769A show small Ir enhancements at the depths where microtektites from the Australasian tektite event accumulated 0.77 Ma” (in here  $\sim 0.788 \text{ Ma}$ ). The microtektite abundances in these and other deep-sea cores confirm that the fluence of tektite matter decreases radially outwards from southeast Asia. Within 5,000 km the fluence falls in a roughly exponential fashion, decreasing by a factor of 2 in  $\sim 400 \text{ km}$ . Extrapolation of the trend to southeast Asia yields a mass fluence in that region corresponding to a melt sheet 4 mm deep, consistent with the formation of layered tektites in this region. The model yields a total mass of tektitic glass of  $3.2 \times 10^{16} \text{ g}$  for the Australasian field, with an estimated uncertainty of a factor of 2. In the absence of Ir fluence data at enough sites to determine a fallout pattern, we assumed a geographical distribution identical to that inferred for tektitic matter. This yields a total Ir deposition of  $8.9 \times 10^8 \text{ g}$  corresponding to a chondritic body (Ir concentration 600 ng/g) having a mass of  $1.5 \times 10^{15} \text{ g}$  or a cometary body with a mass of  $3 \times 10^{15} \text{ g}$ . These compare with an Ir-based estimated mass of the end-

Cretaceous body of  $7 \times 10^{17} \text{ g}$ . An Australasian impactor could have excavated a crater at least having a diameter of 15-19 km.” (Schmidt et al., 1993)

M. Shyam Prasad, Vasudev P. Mahale, and V. N. Kodagali have identified 15 new Australasian microtektite sites along a roughly N-S transect in the central Indian Ocean. “These locations, in addition to the existing 46 sites, total to 61 microtektite sites in the oceans. We carried out regression analysis of a selected area between  $12\text{-}26^\circ \text{N}$  and  $98\text{-}112^\circ \text{E}$ , at  $0.50^\circ$  intervals using three microtektite data sets separately: 41 sites, 46 sites (number of sites known up to the years 1994 and 2006, respectively), and the entire data set of 61 sites. The 41 site data set defines an impact site at about  $12^\circ \text{N}$  and  $106^\circ \text{E}$ . Whereas the contours joining the highest values of the square of the correlation coefficient by using 46 site and 61 site data sets define an area located in NE Thailand-central Laos, the center of this site is at  $18^\circ \text{N}$  and  $104^\circ \text{E}$ . Therefore with progressive increase in microtektitic data the location of the impact site seems to get defined more rigorously. On the basis of the equations for the concentration of the ejecta and the distance from the source region, the calculated crater size has a diameter range between 33 and 120 km.” Prasad et al. (2007)

“On the basis of the distribution of Muong Nong-type tektites, Barnes (1964, 1989) and Wasson (1991) suspected a cometary projectile, while the former suggested in situ melting of soil by a diffuse comet over the whole region of Muong Nong-type tektites, the latter felt that there could have been multiple local sites of impact, which generated the entire strewn field. However, the multiple impact origin has been disputed by the uniform composition of Muong Nong-type tektites (e.g., Blum et al., 1992) and the target stratigraphy inferred from the  $^{10}\text{Be}$  isotope data (Ma et al., 2004). Again on the basis of the geographical variation in composition and the distribution of Muong Nong-type and splash form tektites, backed up by intensive field study, Schnetzler (1992) proposed a site located in the southern part of Thailand-Laos border ( $16^\circ \text{N}$ ;  $105^\circ \text{E}$ ).”

“On the basis of evidence derived from chemical composition and isotope analysis, Chaussidon and Koeberl (1995) narrowed down to a location off the Vietnam coast, near the

mouth of the Mekong River in the South China Sea (9°N 107°E). Ma et al. (2004) analyzed  $^{10}\text{Be}$  concentrations in tektites and on the basis of the isotope distribution pattern felt that a location in the Song Hong basin (off Vietnam, south of Gulf of Tonkin (17°N; 107°E) could be a possible impact site. Further, geochemical and Rb-Sr isotopic investigations by Lee et al. (2004) indicated the provenance of these tektites to be nonmarine: Jurassic sediments, which are found to occur in the southern part of Thailand-Laos. Including the 15 locations in this study, we find that there are 61 microtektite locations that contain the Australasian microtektite peak of abundance along with quantifiable estimates of microtektites (Prasad et al., 2007). Thus today we know 75 locations containing Australasian microtektites, which define the boundaries of this strewn field and the quantity of ejecta generated. For the total microtektite data set with 61 locations, the postulated crater location obtained from the algorithm developed to carry out rigorous analysis is 18°N and 104°E. Therefore the estimated diameter of the crater is approximately 120 km.”

“Prasad et al., 2007 mention that there may occur some uncertainty in determination of the crater diameter based on the error present in the estimates of regression analysis. Thus the uncertainty in the diameter lies between 50 km to 360 km. if we consider the 46 location data from Glass and Koeberl (2006), having slope =  $-4.3 \pm 0.27$  and intercept =  $24 \pm 0.9$ , following the same procedure we get a diameter equal to 124 km. The uncertainty in the diameter of the crater diameter  $\sim 32 (\pm 14)$  km proposed by Glass and Pizzuto (1994) using the same equation. However, Glass and Koeberl (2006) considered solely the unmelted ejecta associated with the Australasian microtektite layer for which they obtained a higher R2 value, and defined a crater diameter of  $43 \pm 9$  km by using the equations given by Stöffler et al. (1975). While the least square method shows much higher levels of uncertainty, calculation according to Stöffler et al.’s (1975) equation gives a crater size much closer to reality as ascertained by Glass and Pizzuto (1994). The estimate lies between 50 km to 350 km. However, by using Stöffler et al. (1975) equation using the concentration of ejecta and the distance from the source, the crater size considering the entire microtektite data is  $33 \pm 8$  km.” Glass and

Pizzuto (1994).

Folco, Rochette. Perchiazzi, D’Orazio, Laurenzi and Tiepolo (2008) report on the discovery of a microtektite (microscopic impact glass particles) strewn field from the Transantarctic Mountains, Antarctica. Microtektites were found trapped in the local detritus accumulated in weathering pits and in joints of several glacially eroded summits ( $\sim 2600$  m above sea level) and distributed latitudinally for 520 km. Their physical and chemical properties define a coherent population with a geochemical affinity to Australasian microtektites and a broad but compatible Quaternary  $^{40}\text{Ar}/^{39}\text{Ar}$  formation age. The margin of the Australasian strewn field is thus shifted southward by  $\sim 3,000$  km and the maximum distance from the possible parent impact site or sites in Indochina by  $\sim 2,000$  km.

### 2.11 Noble gases in Thailand tektites

Studies on gas compositions and their isotopes within bubbles in tektites show a mainly terrestrial origin and confirm their terrestrial origin (Hennecke et al., 1975; Hennecke and Manuel, 1980). “The abundance pattern of noble gases in Thailand tektites shows an anomalous spike of neon, apparently from the diffusion of atmospheric neon through the glass. The isotopic compositions of non-radiogenic Ne, Ar, Kr, and Xe are atmospheric, but the Ar /Xe and Kr/Xe ratios in the tektites are each about an order of magnitude lower than the values in air. The isotopic abundances of Ne, Ar, Kr, and Xe trapped in Thailand tektites are atmospheric in composition, except for an excess of radiogenic  $^{40}\text{Ar}$ . The  $^{36}\text{Ar}/\text{Xe}$  and Kr/Xe ratios are about an order of magnitude lower than those in the atmosphere, suggesting that most of the Xe may be from the tektite parent material.” (Hennecke et al., 1975).

## 3. Loess in SE Asia

### 3.1 Loess

Loess is a windblown sand-silt that is widespread in areas close to Pleistocene ice caps and glaciers. It constitutes very thick deposits (up to 350 m with a mean of 106 m) in northern China and widespread deposits in Europe. It is a yellowish colour and is known in China as the yellow earth and its eroded sediment has given its name to the Yellow River or Huang He. Loess is widespread in




Fig. 15: Mainland SE Asia as seen from satellite reproduced by NRCT showing the catastroloess sheet showing light colour (high albedo) when some reworked as surface windblown sands. In mountainous forestry areas and in river valleys, the loess became disintegrated to soil and obscured.

Thailand, in north Vietnam and Cambodia and the bright yellow to buff deposits have a high albedo and show-up well on satellite imagery (Fig. 15). In NE Thailand, on the Khorat Plateau, loess ranges from about 3m thickness near Khon Kaen, to about 1.5 m near the Thai-Lao border and <1m in much of Laos. Loess rests on a laterite surface that contains tektites (Ford, 1988; Wasson et al., 1995; Fiske et al., 1999). Although it a dominant surficial sediment and is very important as the basis for farming, as an engineering material and as a landfill throughout NE Thailand it has received little scientific attention except for a few publications (e.g. Nutalaya et al., 1987, Udomchoke 1989, Nichol and Nichol, 2015). The grain size distribution of Thai loess is, in detail, different to that of the typical aeolian loess (yellow earth) in North

China and Europe which were deposited close to glaciated areas in the Pleistocene. The Thai and Vietnamese loess may be regarded as anomalous in tropical Thailand (Nichol and Nichol, 2015).

Loesses in Vietnam and Thailand are substantially different, particularly with regard to their contents of sand and clay, the absence of  $\text{CaCO}_3$  or of any effect of wind abrasion. Analysis of the Thu Duc loess horizon (Hoang Ngoc Ky, 1986) shows its sand content in the northeast area to be 46.53 percent, and on islands along the Vietnam coast up to 48.45 percent. The only exception occurs on the Danang Plateau with a sand component of 11.46 percent. The clay content varies from 31.56 to 33.56 percent. Udomchoke (1989) describes up to 70 percent sand and 30 percent silt and clay in the “yellow” loess of the Khon Kaen area in northeastern Thailand. Nutalaya et al. (1986) report that “so called sand sheet deposits near Khon Kaen show characteristics of loess”. We consider this reworked surficial catastroloess, and some were previously called wind-blown sands.

“New analyses of samples taken during the IGCP 296 meeting (October 1990) in Mukdahan Province in northeastern Thailand, which were analysed in the laboratories of the Geological Survey in Bratislava and Prague, Czechoslovakia (Minariková, pers. com.) contain about 60 percent sand, 10 to 15 percent silt and 25 percent clay (Fig.16). Graphic parameters (according to Folk and Ward, 1957) indicate the following values: mean size  $M_z$  varies from 5.33 to 5.48 (e.g. 0.22-0.24 mm), inclusive graphic standard deviation 3.08 reflecting a very weakly sorted material, with graphic skewness around 0.60 showing a very positive asymmetry and graphic kurtosis  $K_o$  that varies from 0.47 to 0.71, corresponding to a very platykurtic distribution. The median  $M_d$  oscillates between 3.60 and 3.87 (0.07 to 0.08 mm) and is thus substantially coarser than the mean size  $M_z$ .” (Sibrava, 1993).

All these values differ from the grain composition as known from the loess sediments in China as well from European loess. As shown above, the main difference lies in the high content of sandy and clay components and the low content of silt in loess from Vietnam and Thailand, contrary to Chinese and European loess covers. Additionally, no substantial surface




Fig. 16: Comparison grain size of loessic sediments on the Khorat Plateau, Thailand (curves 1-3) with typical loess curve from Central European loess (4) (from Sibrava, 1993). Also see Nichol and Nichol (2015).

markings on the quartz grains caused by wind abrasion has been observed in loess of Vietnam and Thailand. Sediments there are totally decalcified.

Loess occurs in several other areas outside of the Khorat Plateau. Loess extends to northern Thailand and covers some river terrace deposits (Vella et al., 1987). As it is easily erodible, when capped by more resistant sediments it erodes into pillars known as ‘sao din’ (earth pillars) in northern Thailand and in eastern Thailand at Sra Kaeo. These widespread deposits remain undated directly but archaeological excavations in the Nanoi Basin, south of Nan in northern Thailand have found stone tools very similar to those in the Chinese Bose Basin which are closely associated with unworked tektites as discussed by Li et al. (2021) and Hou et al. (2000). Zeitoun et al. (2012) suggest an age of 1 to 0.5 Ma for the sao din stone tools based on similarity to those in Yunnan.

Catastroloess erosional feature is very common in catastroloess outcrops; as Pha Wingchoo (Cliff climbing for lover) at Hod; sao din (earth pillars) at Nanoi in the north, Pae Muang Pi (ghost town) in Phrae, Thoeng, Pai; a castle-like wall in south-west Chiangmai, Lampang, east Lom Sak, Sinking holes or land of erosion at west Ratchaburi, Burirum are outcrops of 0.8 Ma loess deposition during wet and humid and dry yearly. The outcrops are

almost readily recognized for numerous, small and even gully erosion, which is not a common feature in stream and alluvial sediments.

Catastroloess is the impact product from continental sediments and gravel beds as unmelted ejecta blown up into the atmosphere, fallen down and form layered beds (very fine stratification), and some hung up in the atmosphere for possibly years. Tektites are melted impact terrestrial rocks, the process was immediately solidified and some splashed to solidify in the atmosphere. The extraterrestrial bodies (comet) that brought in elements from far outer space including Iridium. They usually started the boundary clay that formed a layer immediately ahead of extraterrestrial body or bodies as is known from the K/T boundary. In the catastroloess, or unmelted ejecta, minute fractions contain microtektites, microglasses, PDF’s of microglasses or minute quartz crystals, usually known as spherules. These categories are not expected in ‘normal’ aeolian origin loess. Laboratory procedures are sufficient to separate normal aeolian loess from catastroloess.

Australasian tektites and North China loess have closely related compositions. Wasson and Heins plot compositions of tektites (Chapman and Scheiber, 19691) and loess data from Liu (1985b) with the latter from the classic site at Luochuan, Shaanxi Province. The loess data are corrected for  $\text{CaCO}_3$  prototektitic material and that the  $\text{CO}_2$  lost during the heating event; to minimize the effects of  $\text{CaCO}_3$ , we have excluded samples with CaO contents >38 mg/g. again choosing the cutoff to correspond to a hiatus in CaO.

The clearest trends in the tektite data are the negative correlations between Al or Fe and Si; similar trends are observed in the loess (in disagreement with the statement by Blum et al. (1992) that loess is an unsuitable source because samples of each deposit contain uniform major element compositions). Although ranges in the loess are lower by a factor of roughly 3, the fields overlap. If tektites were produced from loess deposits in SE Asia, these must have contained appreciable amounts of a high-Al and -Fe component such as laterite and of an Si-rich component such as a high- $\text{SiO}_2$  sandstones; sources of such components are widely available in SE Asia. Wasson (1991) conclude that presently available bulk

compositional data are consistent with Australasian tektites having been produced by the melting of loess, spiced by minor amounts of lateritic soils, quartz-rich materials, and even ultramafic (Wasson, 1991) materials. Loess is unmelted product or unmelted ejecta, (together the melted products-tektites), here called catastroloess.

Field evidence in the Khorat Plateau which contains catastrophic loess with more powerful middle size fraction different property from European loess (Fig. 16), such loess are found within the Australasian strewn-field area in the Gobi Desert area in China, and elsewhere in north China including the Luachuan Loess which contain common spherules (Wasson and Heins, 1993; Salyapongse and Pitakpaivan, 1999). The most important fact is they buried tektites and other DCIMs from the impact in Thailand indicating they were immediately syn- and mainly post-tektites in origin in the same event. Thus they cannot be the impact rocks/sediments, but the un-melted ejecta layer from the impact rocks which were soils, gravels and chiefly underlying Jurassic-Cretaceous redbeds in the Indochina impact field, possibly at Buntharik in Ubon Ratchatani a suspected impact crater.

### 3.2 *Catastroloess and sao din in Thailand*

Loess-like sediments (herein catastroloess) are found beyond the Khorat Plateau in hilly areas in northern Thailand, western and eastern Thailand (Figs. 17-30). They are often capped by gravels and once these gravel-bed caps are eroded the catastroloess erodes into earth pillars known as *sao din* and by other local names. The catastroloess is difficult to date definitely, though thorough studies may yield results in the future. At the *sao din* area at Nanoi south of Nan city in Nan Province, stone tools have been found associated with the catastroloess and are comparable with those found in Yunnan and may range in age from 1 Ma - 0.5 Ma, (Zeitoun et al., 2012).

### 3.3 *Catastroloess earth pillars newly recognised in Kunming (SW China)*

Catastroloess in Kunming (Fig. 31) formed among the track of the Australian Cometary Impact and should have an 0.8Ma age. Loess in North China mentioned above is normal loess associated with glaciations.

Catastroloess originated only when



Fig. 17: Mae Cho, Chiangmai above the western ranges



Fig. 18: Pha Wing Choo, Hod, Chiangmai. Typical rill effect of catastroloess covered by 2 m of medium to large, well rounded pebbly gravels, possibly river gravels raised 70 m above present river.



Fig. 19: Pha Wing Choo, Hod, Chiangmai, southern end

impacted, bursting dusting debris and fine sands including nearby ejecta, and largely, in very remote areas, the glacial outwashes were drawn in and sent to a high or very high atmosphere, many hundreds kms above the earth's surface. Catastroloess caused darkness and starting glaciation in northern Asia, China, Australia. These were soon after or almost contemporaneously with the great wildfires of massive felled trees down below were becoming the greatest killing



Fig. 20: Pha Wing Choo from south looking north




Fig. 22: Sao Din (earth pillar), Na Noi, Nan Province




Fig. 21: Pha Wing Choo from Mae Ping, south of Chiangmai



Fig. 23: Sao Din (earth pillar), Na Noi, Nan Province




Fig. 24: Sao Din (earth pillar), Na Noi, Nan Province



Fig. 25: Sao Din (earth pillar), Na Noi, Nan Province



Fig. 28: Phae Muang Pi (ghost town bush) at Phrae, catastroloess column on terrace capped by 1-2 m gravel beds (kindly supplied by Paveena Kitbutrawat)



Fig. 26: Sao Din (earth pillar), Na Noi, Nan Province



Fig. 29: Phae Muang Pi (ghost town bush) at Phrae, catastroloess column on terrace capped by 1-2 m gravel beds



Fig. 27: Phae Muang Pi (ghost town bush) at Phrae, catastroloess column on terrace capped by 1-2 m gravel beds (kindly supplied by Paveena Kitbutrawat)

fields along the trail to NNE, from south Kazakhstan to China, NE Thailand, Australia and disappeared in Mt, Darwin with crater. This was seriously wrongly understood, the tektite event originated by a cometary impact event, a terrestrial catastrophe from an extraterrestrial body from the deep space in the solar system (Wasson, 1985), apart from tektites, the most important was the catastroloess or was known also as ejecta. Chinese loesses are common above 35°N, above Huang Hue River, and around Beijing, not Kunming, which is newly recognised catastroloess in SW China.

We now know Chinese loess had little original connection with the impact although there may be some interbedding of catastroloess



Fig. 30: A-D. Sra Kaeo, eastern Thailand new discovered sao din (earth column) similar to those at Na Noi, 1000 km to the north. Lalu is its local name (kindly supplied by Paveena Kitbutrawat).



Fig. 31: Catastroloess in Kunming – similarly eroded to sao din in Thailand (from Discover the World, Royal Orchid Holidays-a tour guide 2008).

at L8 to L9. With microspherules reported (contamination), but later when related glaciation being developed after the impact, and regenerated as normal loess from glaciation in the whole columns (Li, C.; Ouyang, Z.; Liu, D. and An, Z., 1991). Hence, the Chinese columns contain mainly atmospheric or normal loesses and thin interval in L8 or L9? Contain catastroloess from 0.8 Ma impact with Microtektites and glass microspherules in loess (Li, C.; Ouyang, Z.; Liu, D. and An, Z., 1991).

The known Chinese loesses from “Yellow Mountain” washed into the Yellow River Huanghue River are the atmospheric loesses (‘normal’ loesses) from Pleistocene debris while the real ~0.8Ma Australasian catastroloess is found in Kunming to the southwest.

#### 3.4 Other areas of Thailand with possible catastroloess

Other areas in Thailand with loess include thick sections at Mae Tha in Lampang Province (Fig. 32) and along the Lomsak-Chumphae road (Fig. 33-34). At Mae Tha (Fig. 32), the widespread



Fig. 32: Km 26 to Mae Tha, Lampang, on opposite side young basalt flows on top gravels.




Fig. 33: Km 16, Lom Sak-Chumphae Road.




Fig. 34: Very fine sandy beds at higher levels of Fig. 33.

Lampang basalt is dated in part to about 0.8 Ma, has a reversed magnetism and lies above river gravels which contain an important collection of Acheulian (*Homo erectus*) artifacts (Barr and Macdonald, 1981).

### 3.5 The youngest Pliocene age for Tertiary deposits in Thailand, 1 km deep lake in Lampang tectonically immediately older than Pleistocene

Three diatomite samples from Lampang were brought to New Zealand (Bunopas, 1981) and the diatoms were kindly photographed by scanning electron microphotography and identified by Dr. Margaret Harper at Victoria University of Wellington. Dr. Harper (written comm.) stated that 99% or more of the frustules present are *Melosira granulata* (Ehrenberg) Rafts in Pritchard. Both spined terminal frustules and toothed terminal frustules of the coarse pored form are present. The predominance of this planktonic diatom suggests a deep lake with a silica concentration between 5 and 25 mg/l which is a high concentration (Kilham, 1971) and alkalinity between 10 and 1 meq/l which is relatively dilute. Although *M. granulata* is found in a wide range of climates the very thick walled forms with very coarse pores appear to be characteristic of warm climates (especially sub-tropical to tropical). Also they are better known in older deposits (early Holocene or older) but this may be an artefact of preservation and found only *M. granulata* and two closely related species in some Tertiary deposits in Canada.

A few volcanic glass shards are found with some sponge spicules, some smooth and some with rippled surfaces. The absence of species other than *M. granulata* is strange in terms of present day flora, but is likely to be due to solution of other more delicate frustules during deposition or diagenesis, coupled with a strong initial dominance of *M. granulata*.

A change from fine-grained facies to rudaceous facies marks the base of what is regarded as Quaternary, but the change is not well dated and could be above or below the real Tertiary-Quaternary boundary. It indicates the beginning of the rapid uplift of the present mountains.

The now Phaholyothin road at Km 2 from Lampang to Chiangrai (Fig. 35) exposed excellent outcrop of 3 diatomite beds 1 to 2 m thick each with no conglomerate overlying on



Fig. 35: Neogene diatomaceous beds formed before the catastroloess and uplifted before the impact, Lampang.

top but catastrophic loess indicating they were 1 km uplifted.

### 3.6 Miocene or younger fish fauna at Lomsak, Phetchabun Province, Thailand

Well preserved freshwater fish are well known and well described from shallow excavations near Lom Sak, Phetchabun Province (Figs. 36-37) and have been ascribed a Miocene age (Roberts and Jumnonthai, 1999). The abundant and diverse fish fauna is found in mudstones and siltstones and is overlain by sands here interpreted as loess (or catastroloess). We suggest the possibility that the fish fauna is actually Pleistocene in age and suffered a mass mortality due to the 0.78 Ma impact event.

## 4. Nakhon Ratchasima Province (Khorat) sandpits-evidence of the 08 Ma impact event

### 4.1 Ban Tha Chang (Khorat) sandpits

The sandpits or sand pumping mines at Tha Chang (Figs. 38-46) serve as the most perfect global classroom for the global impact which we have long searched for. These pits were later studied in detail by (Howard, 1999; Howard et al., 1999 and Howard et al., in 2000). The pit in late 1997, on the west side of River Mun south of Ban Tha Chang contains DCIMs reported by Bunopas et al. (1999) are as follows:

1) Numerous extensive burnt felled trees, all roots up, lower levels contain abundant charcoal and sulfides. Uprooted tree trunks up

to 1 metre in diameter are often shattered, branchless, snapped, and burnt to the core. Their presence in deposits throughout Thailand is evidence of widespread burning. These suggested extensive wild-forest-fire which were reported burnt trees or woods the whole of Southeast Asia.

2) Ancient Animals such as Mastodon, ancient species elephant, turtle, crocodiles and etc. suggesting a mass extinction event.

3) Tektites rare splash-form tektites from the pits.

4) Impact crater in Buntharik - though it is still dubious.

5) Catastrophic sands represent debris and ejecta that formed widespread sheet enclosed other DCIMs. This name is proposed "Catastroloess", for this impact sands washed and catastrophic flooded as sands at Ban Tha Chang, Howard (1999) added,

6) Microtektites, glasses, impact quartz and impact glasses etc. and,

7) Isotopic elements from INAA, ICP-MS measurements in tektites from Thailand, China and recent study from Ban Tha Chang sands .

Howard (1999) found the 7 DCIMs and a little later, Howard et al. (1999) emphasized many fluvial facies, microtektites and spherules.

The impact consistent with one from the Buntharik Impact Event at Ubon Ratchatani,



Fig. 36: Lom Sak, Phetchabun Province fish pond, the man stands on the previous bank of the pond where the pond was filled by catastroloess and buried fishes.



Fig. 37: A villager is hunting for the most beautiful fossil fishes. Same location as Fig.36.




Fig. 38: Map of NE Thailand (Khorat Plateau) showing sites and roads, Ford, (1988) and recorded tektite finds. Triangle showing sites of layered tektites around Buriรัมย์ area. (From Howard, 1999).

Thailand and catastrophic products enclosed in their findings and noted that Glass and Pizzutto (1994) and Glass and Wu (1993) had their record of the Australasian field microtektites in the seas.

Geochemical analyses of the sediments of the lower section of pit 1, Ban Tha Chang, reveal that the sediments possess a major oxide distribution that consists of two distinct components.  $\text{SiO}_2$  is the dominant component comprising on average 95.03 wt% of the analysed samples, this is reflected in the dominance of quartz revealed in petrographic analysis. The second component of the samples is a clay component that holds the remaining major oxides and the majority of trace elements. This second and minor component was revealed effectively by the use of Cs and Rb as analogues for clay content.

The sands of Ban Tha Chang possess a trace element geochemical signal that is similar to that typical of the upper crust of Earth (no enrichment in Ir, Ni or Co). However, these sediments are depleted in the elements Ba and Sr relative to most upper crustal sediments. Despite this depletion, barium (mean 108ppm) remains the most abundant element in the analysed samples followed by Zr (mean 60.2ppm), Ce (36.88ppm) and Rb (30.66ppm).

The REE present within the samples show minor light REE enrichment in the stratigraphically lowest portion of the section (sample 1). Generally, however, the samples possess REE patterns typical of the upper crust. The REE geochemistry of the Ban Tha Chang sands exhibit typical features of the REE in that they appear to behave in a uniform manner as indicated by a high degree of inter-correlation. This strongly suggests that these REE are distributed relatively pervasively throughout the mineral phases present in the Ban Tha Chang sands. The use of Cs as an analogue for clay content within the samples analysed reveals that Cs and clay content exhibits a general increase up-section. As may be expected, the abundances of other trace elements correlate well with increases in Cs content; implying that the fine clay component of the sample hosts most trace elements. However, Cs is not a perfect analogue for clay content and despite correlating with Cs, components of many trace elements abundances are not related to Cs. These




Fig. 39: Location of Ban Tha Chang and Chum Phuang sandpits




Fig. 40: Map showing location of sampled Ban Tha Chang sandpits 1 and 2, near Khorat City



Fig. 41: Looking into the southern pit at Ban Tha Chang Clive Burrett and the late Somboon Khositantont in 1997. (Flooded a year later, no stratigraphic measurement).



Fig. 42: Pit 1 east Nam Mun, south of Pit 2



Fig. 43: Pit 2, after being worked for a long time, east Nam Mun



Fig. 44: Pit 3 west Nam Mun (later flooded, not available for stratigraphic measurement).



Fig. 45 A and B: A Pit 2 left . B right. Panorama view of Pit 2L upper and 2R under.



Fig. 46: Sandpit or sand pumping mine at Tha Chang as the most perfect global impact long searching for. Later studied in detailed by Howard et al., 1999, Howard et al., in 2000; Haines et al., 2004).




Fig. 49: Sub-spherical non-tektite micro-impact glasses showing also numerous cracks. The glass shows some fracture plane on its surface, related to post-deformation transport. (Howard, 1999).




Fig. 47: Microtektite or microglass at Ban Tha Chang, Pit 2 (Howard, 1999).



Fig. 50: Close up of above pit. (Howard, 1999).




Fig. 48: Enlarged melted impact dust particle on to the molten glass particle (Howard, 1999).




Fig. 51: Non-tektite micro-impact glasses with pronounced pit structure at the top. (Howard, 1999).




Fig. 52 : Microglass at Ban Tha Chang. (Howard, 1999)




Fig. 53: Very small sub-spherical glass. (Howard, 1999).




Fig. 54: Irregular shape micro-glasses from Ban Tha Chang sands (Howard, 1999).




Fig. 55: Another irregular shaped glass, such extremely irregular morphologies are less common. These are the first terrestrial non-tektite impact glasses to be found in Thailand and are the only onshore non-tektite impact glasses (other than Darwin glass) to be directly associated, by stratigraphic position and geography, with the Australasian tektite strewn field. The analyses by XRD reinforce the overwhelming abundance of quartz in the sands, as well as the minor presence of the minerals rutile, baryte and several other, very minor, minerals. Baryte may itself represent a secondary product formed within the treatment process through reaction with  $H_2SO_4$  (Howard, 1999).

components strongly suggest that minerals such as K-feldspar, zircon and the mica family of minerals are present, in minor amounts, within the sands and are hosting these trace elements.

#### 4.2 *Microspherules, microtektites, microglasses, pdf in quartz crystals, shocked quartz at Ban Tha Chang*

Sandpits at Ban Tha Chang, south of Khorat, northeast Thailand, provide an ideal locality to study the environmental effects of the 0.788 Ma event. Based on the distribution of layered tektites that are believed to form close to the site of an impact, Ban Tha Chang is possibly 400km west or more of inferred impact centres. The sands exist above one of the regionally extensive tektite bearing layers of 788 ka and consist of six interbedded facies outcropping in two active sandpits. Facies 1-4 in the sands have chaotic grainsize distributions and were rapidly transported and deposited in flood events.

The sands contain abundant partially or wholly petrified and/or burnt trees, these were deposited during flood events and are confined to two distinct stratigraphic horizons. The trees




Fig. 56: PDF's in quartz close-up. (Howard, 1999).




Fig. 59: PDF's in quartz. (Howard, 1999).




Fig. 57: NW-SE orientated PDF's in quartz. Note: there is a second set of NW-NE planar fractures (PF's or PDF's ?) visible in the top left corner of this quartz grain (Howard, 1999).





Fig. 60: PDF's in quartz Ban Tha Chang, Scale of A) 0.5 mm and B) 1 mm (Howard, 1999)




Fig. 58: PDF's in quartz. (Howard, 1999)




Fig. 61: 'Cored grain' showing quartz grain with quenched margin. Note: deformation features are absent from the quartz grain. (Howard, 1999).

are branchless, snapped, uprooted, burnt to the core and are very similar to those at Tunguska, Siberia, site of a small impact event in 1908. The destruction of the trees at Ban Tha Chang is related to processes of impact induced shock and forest fires.

Microspherules (typically 100 $\mu$ m) glasses in sands at Ban Tha Chang and also Ratchaburi (west Thailand) are suspected to be of impact origin on the basis of their compositions and morphologies (Figs. 47-55). These glasses are typically sub-spherical to more irregular in form, to rarely teardrop shaped. The glasses may be divided into two distinct groups. Group A impact glasses are composed almost exclusively of SiO<sub>2</sub> (97.7%), while group B impact glasses possess low SiO<sub>2</sub> (37%), high Al<sub>2</sub>O<sub>3</sub> (28%) and anomalous fluorine (6.6%) contents. Group A impact glasses are believed to have formed from a quartz rich photolithography. If a primary chemical signature, the composition of group B impact glasses suggests that these formed from target material rich in fluorine, most likely implying a fluorite rich target e.g. granites. Quartz grains in sands at Ban Tha Chang contain planar fractures and in some grains these are spaced at less than 10 microns and resemble Planar Deformation Features (PDFs) that form at the lower end of impact induced shock pressures, from around 10 to greater than 30 GPa (Figs. 56-60). Also present in the Ban Tha Chang sands are partially melted grains of quartz that are surrounded by a glassy carapace up to 0.1 mm in width. The quartz grains contained within this carapace do not show deformation features but rather a relatively homogenous surface that grades into the outer, glassy carapace

surrounding crystalline quartz (Fig. 61). This texture is characteristic of 'cored grains' (French 1998).

Australasian tektites have been dated at 770  $\pm$  20 ka (Izett and Obradovich, 1992; Bollinger, 1993) and 786  $\pm$  12 ka (Kunz et al., 1995) and 788  $\pm$  3 ka (Jourdan et al., 2019) using <sup>40</sup>Ar-<sup>39</sup>Ar methods. Independent age evidence comes from a deep sea microtektite horizon widely believed to be associated with the tektites on land, which is situated just below the Brunhes-Matuyama magnetic reversal (Schneider et al., 1992). Ages of 793 ka (Lee and Wei, 2000) and 803  $\pm$  3 ka (Hou et al., 2000) have recently been suggested on this basis. Because of the minor variations in these published ages, we have referred to the event herein as the ca. 800 ka impact event. This is equivalent to latest Early Pleistocene extending to earliest Middle Pleistocene.

The main aims of the report are about the DCIMs from the impact which are known, though impact in a very substantial percentage of the World, all DCIMs were found only in Thailand. The DCIMs that formed by the Cometary Impact are mainly 95 % from Ban Tha Chang and Chum Phuang, described by Bunopas et al., 1999; Howard (1999), Howard et al., 2000, 2003; Haines et al. (2004).

#### 4.3 Sedimentology of ~0.8 Million-Year Old Log-Bearing Flood Deposits in Northeast Thailand and Mechanisms for pre-flood Deforestation

Pleistocene alluvial sands within the palaeochannels of major rivers in Northeast Thailand contain a very unusual abundance of large, burnt tree trunks (Figs. 62-67, 70, 82-83). Detailed studies in active sandpits near Khorat reveal 6 sedimentary facies (Figs. 71-81) with tree trunks confined to 2 stratigraphic horizons. Stratigraphy, unabraded tektites and vertebrates, including the pygmy elephant *Stegodon*, suggest a Pleistocene age for the upper sections of the pits with the burnt trees. Facies 1-4 were deposited in catastrophic mega-flood events. These are overlain by 2 facies characteristic of normal fluvial processes. The flood deposits may be associated with the formation of residual topographic features seen on Landsat images, which are similar to those of the channeled scabland of the USA.



Fig. 62: Devastation of a vast area of Siberia due to the 1908 Tunguska airblast leading to an estimated 80 million trees being demolished arranged in a radial pattern. Hundreds of reindeer were killed. Trees near ground-zero were still upright in 1927 but their limbs and bark had been stripped away (cf. Ban Tha Chang). Public domain photograph from the Kulik expedition of 1927. Wikipedia, Tunguska Event.



Fig. 65: Snapped and splintered tree end at Ban Tha Chang (Howard, 1999).



Fig. 63: Ban Tha Chang pit 3 studied by Haines et al. (2004). Note abundance of burnt logs. Photograph kindly supplied by Dr Peter Haines (Australia).



Fig. 66: Assemblage of felled and deposition of trees at Ban Tha Chang Pit 1. (Howard, 1999).



Fig. 64: Tree log, here pyrite appears as the grey colour to the right of the pencil. Note the crust of sulphosalt around the outside, calcite veins in centre.



Fig. 67: Burnt and shattered tree. (Howard,1999).




Fig. 68: Model for effects of initial shock front and thermal pulse on trees from impact. (Howard et al., 2003).

Uprooted tree trunks to > 1 metre in diameter are often shattered, branchless, snapped, and burnt to the core. Their presence in deposits throughout Thailand is evidence of widespread burning and are reminiscent of the huge forest devastation of the Tunguska, Siberia aerial blast in 1908 (Fig. 62).

We suggest a genetic relationship between the mega-flood and deposits, forest burning and the ca. 800 ka impact event (Figs. 68 and 69).

This forest destruction would have greatly accelerated the erosion regime in the area and increased sediment and wood supply to drainage channels including the palaeo-Mun River. Floods ultimately overwhelmed the drainage networks and led to breaching of existing banks and log-jam induced dams in a series of mega-flood events that represent major




Fig. 69: Depositional model for log bearing flood deposits.

A. Early stage log jam.

B. Log-jam density increase, reduced flow and deposition of mud lenses during this low energy period.

C. Pressure builds up behind log jam until the point of failure. On failure, a high energy flow resulted and deposits the sand, gravel, and logs in distinct stratigraphic horizons. (Howard et al., 2003)

topographic modification of the environment (Fig. 69).

Some logs have longitudinal fractures and snapped ends. Such logs show evidence of insect burrowing (Fig. 84) (Howard et al., 2003).

#### 4.4 Tektites in Khorat sandpits

Completely unabraded tektites were found throughout unit A (Fig. 85). These tektites show little evidence of having been transported. And are comparable, if not better, in their preservation than those from famous locality near Port Campbell, Victoria, Australia (Shoemaker and Uhlherr, 1999). Tektites were also found in unit B but these are abraded and have spent considerable time being reworked and transported. Geochemical analyses of tektites from Ban Tha Chang and Chum Phuang show that they belong to the Australasian strewn field. Haines et al. (2004) found one very fragile tektite at Chum Phuang that is extremely unlikely to have been reworked (Fig. 85).

#### 4.5 Palynology at Ban Tha Chang sandpits

Pollen sequences for two pits at Ban Tha



Fig. 70: Ban Tha Chang sands, Pit 2, with abundant burnt wood near bottom removed by sediment hosing.



Fig. 72 B: Pyrite on quartz grains. (Howard, 1999)



Fig. 71: Facies 1, cross bedded sands (Howard,1999).




Fig. 73: Pyrite in burnt trees. Pyrite occur as 'veinlets' in fracture within the trees fragment. Pyrite also forms a disseminated grains throughout charcoal in petrified trees at Ban Tha Chang sands. (Howard, 1999).




Fig. 72A: Facies 1 under binocular microscope, note the poor sorting in this facies (Howard, 1999).




Fig. 74: Charcoal in facies 1 (Howard, 1999).



Fig. 75: Facies 2: structure-less sands (Howard, 1999).



Fig. 76: Ban Tha Chang. Facies 3: silt. (Howard, 1999).




Fig. 77: Ban Tha Chang. Facies 4: Pebble/gravels. (Howard, 1999).




Fig. 78: Facies 5 and 6 on the upper part of Ban Tha Chang Pit 2. The distinct beds are of facies 5. Total thickness of facies 5 = 3m. (Howard, 1999).




Fig. 79: Type section at Pit 1, left 14 m high; Pit 2 right 15 m high. (Howard, 1999).




Fig. 80: Lithofacies on the Ban Tha Chang sands stratigraphy of Pit 1, left and Pit 2, right. (Howard, 1999).




Fig. 81 A: Ban Tha Chang. Pit 1 B: Ban Tha Chang. Pit 2 (Howard, 1999).




Fig. 82: Large limbless logs in flood deposits at Chum Phuang. Note man standing on log (1.87 m tall) for scale




Fig. 83: Variably burnt woody material in the flood deposits.

A. Log that is completely composed of charcoal.  
 B. Partially burnt log. The burnt outer margins are crumbling away from the non-burnt core.

C. Partially scorched log. The surface of this log has been variably singed by intense heat before the flames were extinguished.

D. Small burnt pieces of wood in sand at Ban Tha Chang, logs are more evenly distributed throughout the profiles.




Fig. 84: Insect burrows in non-burnt wood from Ban Tha Chang. (Howard et al., 2003) (Bar scale = 1 cm)




Fig. 85: Measured section from sand quarries at Ban Tha Chang and Chum Phuang showing the position of the main log horizon. The polarity of magnetic remanence in samples collected and analysed by Haines et al. (2004). Measured sections were terminated at the top when material disturbed by mining was encountered. Unpublished colour Fig. kindly provided by Dr Peter Haines (Australia).

Chang by Maloney and Howard (2000) are presented in Fig.86. These data are percentage-based and the Pinus pollen data should be ignored when characterising the past local vegetation because pine pollen are widely dispersed and may have been distantly sourced. Podocarpus, Castanopsis / Lithocarpus, and Quercus are likely to be from forests above 1,000 meters elevation. Salix tetrasperma and Elaeocarpus species are common along river courses.

Both sections show an upward decrease in pollen diversity and a simultaneous proportional increase in grasses. Gramineae pollen dominates all samples and this suggests that vegetation was open. Most Thai trees are pollinated by insects and animals and do not produce much pollen and maybe are significantly under-represented. Samples P2-7 and S3-13 contain phytoliths from Panicoid genera grasses. Some of these phytoliths from S3-13 contain carbon

inclusions that indicate they have been burnt (Maloney and Howard, 2000).

The occurrence of the fern *Pteris* sp. suggests re-growth after disturbance. The pollen record shows that there was forest stretching from the banks of the paleo-Mun to above 1,000 meters elevation before this was felled and the trees found their way into the river.

#### 4.6 Geochemistry of Ban Tha Chang sands

Results of major and trace element analyses of five samples taken from cross-bedded and sand at the base of section 1 are shown in Fig. 87, along with calculated chemical index of alteration weathering index values. The major oxide composition of the sediments indicates at two-component mixture between quartz and clay. The abundance of silica dioxide reflects a quartz-rich provenance and this is consistent with petrographic observations. In the study region, this is most likely the weathered basement sandstones of the Khorat Group. This provenance



Fig. 86: Pollen sum, total pollen , and pteridophyte spore Gramineae. Sample from Ban Tha Chang. Many are present-day Khao Yai species. (Maloney and Howard, 2000)

and the high degree of weathering indicated by chemical index of alteration values greater than 70 explain the low abundance of other oxides (Nesbitt and Young, 1982). Immobile elements typical of clay, such as cesium and rubidium, correlate strongly with oxides other than silicon dioxide. This also reflects the effects of weathering.

4.7 Ages of sediments, palaeomagnetism and fossil mammals at Ban Tha Chang and Chum Phuang.

4.7.1 Ages of sediments at Ban Tha Chang

Palynomorphs collected from Ban Tha Chang are consistent with a Quaternary age but do not allow for any finer resolution (Maloney and Howard, 2000). Charusiri et al. (2002) dated sand samples from Ban Tha Chang using thermoluminescence (TL) techniques. Their preliminary data produced a wide scatter in thermoluminescence model ages that range from 0.623 to 4.338 million years (Charusiri et al., 2002). However, the level with burnt tree

trunks and tektites gave a TL age of 0.748 Ma. along with a similar age for tektites.

“Radiometric dating of Australasian tektites repeatedly produces ages of about 0.8 million years (Kunz et al., 1995 ; Hou et al., 2000; Jourdan et al., 2019). The position of a deep-sea microtektite horizon confirms that the Australasian tektites were formed about 0.012-0.0165 million years prior to the Brunhes/Matuyama polarity reversal that is dated at 0.78 million years (Schneider et al., 1992; Lee and Wei, 2000). The presence of tektites in unit A constrains the maximum age to ca. 0.8 million years. Haines et al. (2004) have demonstrated that mud layers from unit A at both Ban Tha Chang and Chum Phuang were deposited under reversed polarity. This constrains the deposition of unit A to the period between the time of the tektite fall and the Brunhes/Matuyama polarity reversal. This is between about 0.8 million years and 0.78 million years. The completely unabraded nature of tektites in unit A suggests rapid deposition after the tektite-forming impact event.” (Haines et al., 2004).

Sample	1	2	3	4	5	Average	Sample	1	2	3	4	5	Average
SiO <sub>2</sub> (%)	95.0	86.7	91.0	90.3	85.2	89.6	Sb (ppm)	0.3	0.4	0.4	0.4	0.5	0.37
TiO <sub>2</sub> (%)	0.2	0.4	0.3	0.3	0.5	0.3	Cs (ppm)	0.9	2.8	1.7	2.2	3.6	2.22
Al <sub>2</sub> O <sub>3</sub> (%)	2.1	5.4	3.8	4.2	7.0	4.5	Ba (ppm)	90.6	114.2	104.0	102.9	128.8	108.1
Fe <sub>2</sub> O <sub>3</sub> (%)	0.6	0.1	0.9	1.0	0.1	0.5	La (ppm)	15.4	17.7	15.6	18.2	19.6	17.3
MnO (%)	<0.1	<0.1	<0.1	<0.1	<0.1	0.01	Ce (ppm)	34.8	35.5	32.8	41.6	39.7	36.9
MgO (%)	0.1	0.3	0.2	0.2	0.4	0.2	Pr (ppm)	3.9	3.9	3.6	4.4	4.3	4.0
CaO (%)	0.1	0.2	0.1	0.1	0.1	0.1	Nd (ppm)	15.7	15.1	13.6	17.6	16.6	15.7
Na <sub>2</sub> O (%)	0.4	0.1	0.6	0.5	0.7	0.4	Sm (ppm)	3.2	2.8	2.5	3.5	3.1	3.1
K <sub>2</sub> O (%)	0.3	0.7	0.5	0.5	0.8	0.5	Eú (ppm)	0.6	0.5	0.4	0.7	0.6	0.5
P <sub>2</sub> O <sub>5</sub> (%)	0.0	0.0	0.0	0.0	0.0	0.0	Gd (ppm)	2.1	1.8	1.6	2.4	2.0	1.9
Li (ppm)	9.1	14.8	12.1	12.5	18.0	13.3	Dy (ppm)	1.9	2.0	1.6	2.4	2.2	1.9
Be (ppm)	0.4	1.2	0.8	0.9	1.6	.9	Ho (ppm)	0.3	0.4	0.3	0.5	0.5	0.4
Sc (ppm)	1.7	4.9	3.5	4.0	6.8	4.1	Er (ppm)	0.8	1.1	0.8	1.1	1.3	1.1
V (ppm)	12.7	42.9	26.5	30.8	47.7	32.1	Yb (ppm)	0.7	1.3	0.9	1.1	1.4	1.1
Cu (ppm)	5.0	12.2	8.3	8.8	15.2	9.8	Lu (ppm)	0.1	0.2	0.1	0.2	0.2	0.1
Zn (ppm)	8.8	26.1	14.6	20.8	32.6	20.5	Hf (ppm)	0.9	2.2	1.6	1.6	2.5	1.7
Ga (ppm)	2.3	5.9	4.1	4.8	7.8	4.6	Ta (ppm)	0.2	0.5	0.4	0.4	0.6	0.4
Rb (ppm)	13.9	38.2	24.5	28.6	48.1	30.66	Pb (ppm)	6.0	8.0	6.9	8.2	10.0	7.8
Sr (ppm)	18.4	37.5	25.6	23.5	39.1	28.8	Th (ppm)	3.3	6.3	4.7	4.8	6.9	5.1
Y (ppm)	10.6	12.7	9.4	14.1	14.4	12.2	U (ppm)	1.0	1.5	1.0	1.4	1.7	1.3
Zr (ppm)	32.3	85.0	57.8	57.9	97.8	66.1	Co (ppm)	3.3	7.4	5.5	6.6	6.3	5.8
Nb (ppm)	2.7	7.2	5.2	4.9	8.5	5.6	Ni (ppm)	5.1	8.4	6.9	6.6	11.1	7.6
Mo (ppm)	0.2	0.2	0.1	0.1	0.2	0.19	Ir (ppb)	<1.1	<1.2	<2.0	<1.0	<2.0	<2.0
Sn (ppm)	0.5	1.2	0.8	0.9	1.4	0.97	CIA 72.1	85.4	76.9	80.5	81.13	79.2	79.2
							Total (%)	97.9	92.7	96.1	95.8	93.0	95.1

Fig. 87: Analytical results for five samples, TC1 to TC5, from the base of section 1 at Ban Tha Chang. Major elements by X-ray fluorescence and trace elements, excluding Ni, Co, and Ir, by solution inductively coupled plasma mass spectrometry at the University of Tasmania. Ni, Co, and Ir by ignition neutron activation analysis at NASA Johnson Space Center. Chemical index of alteration values calculated using equation of Nesbitt and Young (1982) suggest a high degree of weathering in these sand samples. (Howard et al., 2003)



Fig. 88: Field photograph of log and tektite bearing deposits in a gravel and sand quarry near Chum Phuang (15.381° N. 102.728°E). Unit B is overlain by material transported during excavation. (Haines et al., 2004). Photograph kindly supplied by Dr Peter Haines (Australia).



Fig. 89: Field photograph of a major log-bearing horizon at Chum Phuang. Some of the logs have been washed out of the enclosing sediments by water cannon operations. (Haines et al., 2004). Photograph kindly supplied by Dr Peter Haines (Australia).

#### 4.7.2 Palaeomagnetism of sediments at Ban Tha Chang and Chum Phuang

“Palaeomagnetic evidence including reversed polarity in mud lenses closely associated with unabraded tektites suggests that tektite-bearing flood deposits near Ban Ta Chang and Chum Phuang are penecontemporaneous with the impact event. The deposits include abundant organic debris, including whole tree trunks (Figs. 88-89) and mammal bones that were preserved due to reducing conditions, which are also responsible for the presence of abundant iron sulphides. Sedimentological observations

suggest a series of major flood events that are out of character with the modern meandering river system to which they are related. The deposits are consistent with the effects of regional deforestation, increased run off and erosion, and other environmental disruptions expected in the aftermath of a major impact event”. Haines et al. (2004).

Tektites by themselves only provide a oldest age limit for the deposits in which they occur, and claims of tektites embedded in sediments dating from the time of fall (e.g. Hou et al., 2000) have been criticized for lacking independent

age constraints (e.g. Koeberl and Glass, 2000). Previous authors (e.g. Howard et al., 2000) have suggested that some ancient flood deposits in Thailand may be related to the ~ 0.8 Ma impact event. Haines et al. (2004) confirmed the presence of in situ Australasian tektites within these deposits and used palaeomagnetic and other evidence to provide independent minimum-age constraints.

“The sand-pit deposits are very similar at both locations, comprising distinct lower and upper units, separated by a marked discontinuity (Fig. 85). The upper unit (unit B), is typically about 5 m in thickness and continuing to the surface, comprises sand, fine gravel and silt considered to represent normal channel and floodplain deposition produced by the migrating meander belts of the Mun River. This unit is oxidized, has a brown-yellow color, and contains minimum organic matter, limited to minor isolated fragments and logs of wood, the latter possibly reworked from the underlying unit. The lower unit (unit A) is markedly different, comprising crudely stratified and cross-bedded sands and gravels, with minor lenticular mud horizons. Logs of wood, including entire limbless tree trunks, some in excess of 10 m in length and complete with tap roots are common throughout the unit, but are particularly abundant in discrete horizons. Some of the wood is partly or wholly replaced by silica and pyrite, but most is preserved as original woody tissue. The local presence of charcoal on the same logs is confirmed by Environmental Scanning Electron Microscope (ESEM) studies (Howard et al., 2003). Fragmentary mammal bones are also present. The entire lower unit is chemically reduced, with a grey color and ubiquitous pyrite/marcasite present as disseminated material and as encrustations around wood and bone.” Haines et al. (2004).

Several previous workers have reported tektites in the sand quarries at Ban Tha Chang e.g. (Charusiri et al., 2002; Sato, 2002). “We searched all measured stratigraphic sections and adjacent exposures for tektites and were successful in locating four tektites in situ; one from unit A and two from unit B at Ban Tha Chang, and one from unit A at Chum Phuang. The specimens from unit A have sharp broken edges and no signs of fluvial abrasion while both from unit B have clearly been abraded by prolonged fluvial action. In the former, the sharp

delicate broken edges show evidence of light chemical etching indicating that breakage preceded burial and is not of modern origin.” Haines et al. (2004).

#### 4.7.3 Palaeomagnetic studies of sand section from the pit

Palaeomagnetic studies were carried out on oriented mud samples collected from unit A along three measured quarry sections; two at Ban Tha Chang and one at Chum Phuang. “Available sedimentological evidence strongly suggests that at least the upper well exposed section of unit A was deposited rapidly, but probably in several discrete pulses as indicated by the lenticular mud interbeds. Two independent lines of age evidence are broadly supportive of our age interpretations. Bones that we collected within unit A at Ban Tha Chang include common *Stegodon* sp. (J. de Vos, personal communication, 2002) indicating a mid-Pleistocene or older age, although it is possible that some bones could be reworked from a lower horizon. Thermoluminescence dating of quartz grains from tektite-bearing sands at Ban Tha Chang gives a reported age of  $0.75 \pm 0.13$  Ma (Charusiri et al., 2002; Howard et al., 2003). Unit A is interpreted as a series of high-energy flood deposits. They are unusual in their preservation of abundant organic debris, most notably large complete tree trunks. Fast flowing river channels and sheet floods are normally oxidized environments. The reduced nature of these high-energy deposits is thus somewhat paradoxical, unless the large amounts of entrained organic matter were buried and sealed rapidly, before significant aerobic decay was possible. Under such sealed conditions, oxygen is quickly exhausted during incomplete aerobic decay leading to reducing conditions and development of iron sulfides by the sulfate reduction activity of anaerobic bacteria (Saunders et al., 1997). The lack of oxidized, rooted or bioturbated surfaces, suggests that the flood pulses were closely spaced in time, rapidly infilling the pre-existing incised river channel or valley, probably with the aid of frequent debris-induced stream damming.”

“Large impact events will be accompanied by regional environmental destruction, including regional deforestation and other changes to the landscape (Toon et al., 1997). Inevitably this will lead to greatly increased erosion and runoff




Fig. 90: Location of fossil locality (sand pit), Ban Non Man Thet, in Nakhon Ratchasima. (Reproduced from 1/50000 scale map Amphoe Non Sung, 5439/U (Left) and columnar section of fossil locality, new pit (right). (Sato, 2002).

resulting in floods of unusual magnitude, carrying higher than normal sediment loads and plant debris resultant from the deforestation. Such floods would be most likely in the years immediately following the event, before significant recovery of soil binding vegetation. Combining palaeomagnetic evidence with the presence of tektites leads to the conclusion that some or all of unit A was most likely deposited very soon after the ~ 0.8 Ma tektite producing impact event in Southeast Asia. As such, flooding may have been a direct result of increased runoff following catastrophic deforestation, as can be predicted for any large impact event on or near land.” Haines et al. (2004).

[Editor’s note: These palaeomagnetic, age and palaeoenvironmental conclusions are markedly different from those of Yang and Grote (2018 a, b) who reported thermoluminescence dating on 7 samples and two C14 dates ranging from 172,739 ± 22,400 a BP to 27,332 ± 3000 a BP in a 3.25 m section from about 13.5 m to 10.0 m depths which covers the same depth interval in the Ban Tha Chang sandpits as the

study by Haines et al. (2004). From palynological studies, Yang and Grote (2018 b) suggest major vegetational and palaeoclimatic changes from cool to warm to cool, and correlation of the sediments to parts of two glacial periods (Riss and Wurm) during the deposition of 3.25 m of sediments which Howard et al. (2003) and Haines et al. (2004) argue were deposited within, at most, a few years.]

#### 4.7.4 Fossil mammals at Ban Tha Chang by Sato

“At Ban Non Man Thet, west of Ban Tha Chang, Khorat an extraordinary rich fauna of fossil mammals have been found in the eastern part of Nakhon Ratchasima. A tektite is also found in an upper horizon in the same locality (Fig. 90). The occurrences of fossils are allochthonous. Molars and bones are scattered in the sediments, but they are not derived fossils. Mammal fossils consist of bones and teeth. Lower than the tektite horizon, every horizon yields fossils. Almost of bones are scattered in the beds and thin bones like scapula, ilium and

ribs, are deposited parallel to bedding plane. Though the fragmentation of bones is frequent, serious abrasions of the bone surfaces are not observed. Vertebrate bones also are not articulated, but all the processes are nearly preserved. Skulls have not been discovered in this new pit. In the course of study, the following taxa are identified: *Rhinoceros* sp., *Stegodon* sp., *S. (Eostegodon)* sp., *Bubalus* sp., *Cervus* sp., *Equus* sp., *Mastodon* sp. and *Stegolophodon* sp.. At this moment, specimens are not enough to determine species. In species, for the identification of species belonging to Stegodontidae and Mastodontidae, it is necessary to obtain more molar specimens through growth stages.

“In this locality excepting fossils, geological age assignment is only the tektite. In Fig. 90 (below), tektite yielding horizon is the third horizon in the columnar section. On the other hand, obvious unconformities are observed between the boundary second horizon and third horizon, third horizon and fourth horizon and fourth horizon and fifth horizon. According to Ducrocq et al. 1994) the Mastodon horizon in north Thailand is the late early Miocene. The stratigraphical distribution of *Stegodon* and *Eostegodon* is also restricted to Pliocene (Kawamura, 1998). So the geological age of the third horizon is considered to be Early Pleistocene, the fourth horizon is late Pliocene and the fifth horizon is early Miocene” Sato (2002).

“Most part of this formation is Miocene. In the present course of study, only a few molars of *Mastodon* and *Stegolophodon* were collected from only the fifth, lowest horizon.”

“The old pit was much deeper than a new pit. Total number of thickness may be larger than 50 m. So, thickness of fifth horizon is expected to be larger than 30 m. The mammal fossils of *Elephantoidea* deposited in DMR and universities are supposed to be collected from the fifth horizon.”

“It is very dangerous to determine the geological age by using only mammal fossils. In this locality, credible data concerning the geological age is only tektite, 0.7 Ma. It needs another method, for example, fission track method by using zircon to fix the geological age.” (Sato 2002)

#### 4.8 Neogene mammalian fauna and reptiles

*in other areas in Thailand and neighbors – mass mortality in ~0.8 ma impact*

Neogene mammalian fauna in Thailand are started researches, and found interesting results that could be relevant to the later ~0.8 Ma impact.

The Japan-Thailand Fossil Expedition Team, reported their paleontological survey in the Late Cenozoic basins of Thailand during the 1996 to 2002 field seasons, and re-examined the geology and mammalian faunas of the following fossil sites: Mae Soi, Chiang Muan, and Sop Mae Tham Northern Thailand); and Tha Chang (Northeastern Thailand). From the sand pits in Tha Chang they found three new Late Cenozoic mammalian faunas at least: the middle Miocene fauna consists of amebelodontid gomphothere and *Prodeinotlurium*: the latest Miocene to early Pliocene fauna yields *Hipparion*, *Stegolophodon*, and *Merycopotarnus*; and the early Pleistocene fauna yields advanced *Stegodon* (Figs. 96-98). The discoveries of Pliocene and Pleistocene (in Tha Chang) mammalian faunas are the first reports of mammalian faunas of these ages in Thailand. (Nayake et al., 2003)

#### 4.9 Khok Sung locality, Nakhon Ratchasima Province

“Khok Sung sand pit is situated in Nakorn Ratchasima Province, Northeastern Thailand. Concerning the Pleistocene, a rich and diversified Middle Pleistocene mammalian fauna has been excavated from ancient river deposits at Khok Sung sandpit in Nakhon Ratchasima Province which has delivered outstanding fossils like a complete skull of a spotted hyaena which is extinct in Asia (Chaimanee, 2006), several complete skulls of a long snouted gavia, a fish eating crocodile which is also extinct in Thailand. Associated is an extinct elephant, *Stegodon* and deer, buffalos, gaur and banteng remains, it represents the best preserved Middle Pleistocene fossil assemblage from Thailand. (Fig. 99). Usually these animal remains are only documented from caves where their remains have been fragmented by predators and by porcupines. Therefore mostly isolated teeth are left in cave sediments. From such caves we have also discovered spotted hyaena in the Peninsular Thailand near Nakhon Si Thammarat Province and in several other places in Northern Thailand. Where they are associated with remains of giant





Fig. 94: Large turtle from Siam sandpit Khorat. Note hammer for scale.



Fig. 95: Turtle as Fig. 94. Siam sandpit Khorat.

deposits are over represented due to the numerous Tertiary basins. Climate dramatically changed during the ice ages in the Pleistocene (Figs. 91-95), where important latitudinal shifts occurred, some taxa of northern areas such as the giant panda and *Hadromys humei* (an Assam rat) moved southwards to the Thai-Malay border (Chaimanee & Jaeger, 2000).” (Chaimanee, 2007).

(Ed: More recent work at Khok Sung by Duangkrayom et al. (2014), Duval et al. (2019) and Suraprasit et al. (2015, 2016) have conducted



Fig. 96: Tusk of an elephant in a sandpit in Khorat.



Fig. 97: Stegodon reconstruction fibre-glass models of Sivalik Hills, Nepal specimens. Tusks are 5m long. From: I, Vjdchauan, CC BY-SA 3.0, <https://commons.wikimedia.org/w/index.php?curid=2449818>).



Fig. 98: Stegodon teeth, Khorat sand-pit.



Fig. 99: Reconstruction of the environs of savannah forest of Ban Khok Sung along the palaeo-Mun River, in Khorat (Nakhon Ratchasima), NE Thailand at c. 217 ka or 130 ka (reconstruction from Chaimanee, 2004). A broadly similar reconstruction is possible for the probably older Pleistocene deposits c.788 ka at Ban Tha Chang.



Fig. 100: Early operation of open pit Bo Phloi gem mine western Thailand with yellow sands (loess) on top and numerous burnt felled trees on gravels with sapphire grains.




Fig. 101: Bo Phloi gem mine, western Thailand upper (A) sands (loess), lower (A) weathered gravels, (B) gravels with weathered basalt with sapphire. Burnt trees are underlying sands on top of gravels.



Fig. 102: Basaltic flow sheet melted underlying catastroloess as many rolling tennis balls stick to sheet in the lower sheets of basalt hills south of Buriram town (sample by courtesy of Dr. Chumpol Vichiansilp of Buriram Rajabhat University).

*magnetostratigraphy, updated the taxonomic composition of the fauna, provided a good sedimentological analysis and have suggested two possible dates at about  $217 \pm 36$  ka or  $130 \pm 29$  ka based on ESR dating and magnetostratigraphy).*

**4.10 Bo Phloi sapphire quarries, Kanchanaburi Province, western Thailand.**

The sapphire bearing sand-pits at Bo Phloi in Kanchanaburi Province, western Thailand have long been known to contain abundant apparently burnt trees but these deposits have not yet been dated. The upper horizon above the abundant burnt trees (Figs. 100 -101) is probably loess and may, also be

**4.11 Buriram basalt enclosed by the catastroloess to form rolling balls.**

An interesting sample of basaltic sample with firmly fused by a half golf size sandy loess (catastroloess to be more accurately) was given to us, during Buriram basalt, among many varied sizes of rolling balls of semi-melted loess. All of these samples indicated flowing bands of high heated basaltic flows flown over catastroloess horizon and were meted by basaltic flows flown across. Some loess stick under the flows and some appeared as many rolling balls.

Fall from the impact ended, and were over flowed by later basaltic flows. The sample indicates that parts of Buriram basalt must be younger than 0.8 Ma, which was the age of the catastroloess (Fig.102). Older  $^{40}\text{K}/^{39}\text{Ar}$  dates by Barr and Macdonald (in Suthirat, 1999) was  $0.92 \pm 0.3$  Ma and 0.43 Ma. More recent Ar/Ar work by Chualaowanich et al. (2008) also give dates of 0.49-0.43 Ma. If these dates and



Fig. 103 A: Mudflow gravels that buried petrified burnt felled trees in contemporaneous mudflows and avalanches with gravels at south Khorat at a terrace at Ban Krok Duen Ha. The gravels represent previous vegetation surface on common terraces when impact arrived. B: A splashed tektite specimen was found embedded in the mudflow covering the tree. The terrace gravels seen under, nearby catastrophes to east and west.



Fig. 104: Petrified tree within river terrace gravels at Ban Krok Duen Ha, Khorat. Note man for scale.



Fig. 105: Tree (73 m long) found in river terrace gravel at Ban Tak District, Tak, NW Thailand. Pleistocene - exact age unknown. Ban Tak District west of Tak, *Azelia xylocarpa* (Kurz) (Photo kindly supplied by Cherdchan Pothichaiya)



Fig. 106: Same as Fig. 105. Tree found in terrace gravel at Tak, NW Thailand. Pleistocene (Photo kindly supplied by Cherdchan Pothichaiya)

conclusions are correct, an age constraint on the loess is provided of  $>0.43$  Ma.

#### 4.12 Ban Krok Duen Ha, Khorat. Tree and tektite in terrace deposit. Huge tree in terrace deposit in Tak, northwest Thailand

Buried petrified burnt tree are found in the contemporaneous mudflows and avalanches with gravels at south Khorat at a terrace at Ban Krok Duen Ha (Fig. 103). A splash form tektite specimen was found embedded in the mudflow covering the tree (Fig. 104). The terrace gravels marked the impact surface, seen under nearby catastrophes, uprooted hard wood tree trunk, burned, was buried under terrifying earthflows, later formed gravel beds at near Wat Krok Duan



Fig. 107: Photo of a new lignite excavation prospecting taken in January 2000 at immediately south of the road from Hod to Mae Sariang at the beginning of the road from Suan Son to Omkoi in inferred Precambrian terrain.

Ha, at Khorat Petrified trees Museum site in Khorat on the Khorat Plateau. When impact all trees grown on older gravel terraces were suddenly uprooted fallen flat and almost all of the catastrophic burned or burnt trunks pointing their roots ends to the blasting impact on Ubon Ratchatani etc., and some are entrained by surface slump, landslide and earthflows. The gravels were mixed with atmospheric falls of catastrophic loess, splashed tektites until none but catastrophic loess or the catastroloess for more than 10m. We nominate the as this Q1/Q2 boundary the new and more advanced datings ~800 ka.

*4.13 Suan Son Lignite Quarry (Bo Luang Prospect), Hod, 90 km south of Chiangmai*

Suan Son Lignite Quarry (Bo Luang Prospect, now Lam Thong Mining in 2002) is recently known (later than 1990, after the Geological Mapping Program of Baum et al., 1970; and Hahn et al., 1986) in the previous Tertiary basins in the Precambrian terrain of the Western Mountains described and the study above by Uttamo et al. (1999). Tertiary beds were truncated by grey muddy sands (Fig.107) while uplifting to more than 1000 m above sea level and were finally overlain by brownish red

catastroloess sands dating to 790 ka. This was indicated that the general uplifting of this area was active before and during 790 ka, as similar to the uplifting of diatomaceous beds that was uplifted, in Fig. 107, before the catastroloess was falling in. This also implied that the uplifting of the Western Mountains were uplifting during Lower Pleistocene as was real-time correlated dating from the catastroloess. Almost everywhere catastroloess were over the high to medium terraces excepting floodplain. See Uttamo et al. (1999) for more new information on Tertiary sequence on the Western Mountains.

At Bo Luang, Suan Son (Pine trees forest) in the “former western mountains and the peninsular parts”, tilted and slightly folded Tertiary with the unconformably overlying 2 meters dark grey muddy-sandy soil is conformable with the grey beds by three meters of reddish yellow catastrophic sands (catastroloess) and topsoil found in early 2000. There is not any conglomerate between folded lignite beds or the grey topsoil and the reddish yellow catastrophic sands (catastroloess). It is most likely means when the lignite beds folded the measures soon covered by topsoil and was relatively uplifting while the nearby areas deposited gravel beds (Bunopas,



Fig. 108: Sand beds at sand quarry west of Na Thawi, southern Thailand also showing a little more than 6 m (top parts above distinctive fault). Lower parts are faulted sandy gravel beds of pre-~0.8Ma impact.

1981).

Probable loess or catastroloess is also found in southern Thailand as at Na Thawi (Fig.108).

#### THE INTHANON EPEIROGENY IN THAILAND

Another effect of the immediately pre-castroloess was shifting of courses of rivers such as the Mun and Chi. Mekong River that was already moved eastwards from the Grand Lake in Cambodia towards southern Laos and northeast Cambodia. The Grand Lake (Tonle Sap) itself became overfilled and deserted. Hartung (1990) and Stauffer (1978) suggested a tektite source crater in the Tonle Sap, but this is considered very unlikely.

The Khorat Plateau Early Quaternary Vertical Uplifting was after Long Cenozoic Cratonization of the Mesozoic Khorat Group. Its Northeast Tilting Diverted the Mae Khong River before the 0.8 Ma Cometary Impact Catastroloess and the overflowing Mid-Pleistocene Buriram Basalt. New discoveries in northern Thailand suggest the late Pliocene to lower Quaternary Inthanon Epeirogeny and uplift from 200 m to 1,000 m, or even to 2,600 m in the western mountains of Thailand (Bunopas et al., 2004). The Khorat Plateau early Quaternary vertical uplift, after the

long Cenozoic cratonization of the Mesozoic Khorat Group, its northeast tilting eastwards diverted the Mae Khong River before the 0.8 ma cometary impact catastroloess and the overflowing Mid-Pleistocene Buriram and Indochina basalts. Palaeomagnetic reversal polarity results from the catastroloess ensemble show the real-time catastroloess from the Buntharik Event of ~0.8Ma cometary impact are absolutely well chronologically constrained.

The base of the impact in the Buntharik Event is now absolutely certain at the level of ancient felled burnt trees, covering by earth and mudflows on the blasting and heat wave followed shortly by thunder storms and heaviest rains on earth resulting in the buryial of trees. In the avalanche debris at Ban Krok Duen Ha contain pioneer splash tektites, and this level on ancient high and medium terraces in Khorat is only few m. below lateritic sands in Ubon Ratchatani known by Fiske et al. (1996) near the Chulaphorn Dam filled with splash tektite in a row of catastroloess. Above the horizon are massive clayey sands here called catastroloess for some meters.

Settling of enormous debris and fine sands in the later stage given rise to the accumulation



Fig. 109: *Homo erectus* skull (Peking Man also Java Man in Indonesia). Photograph by Yan Li of specimen in Paleozoological Museum of China ([https://en.wikipedia.org/wiki/Homo\\_erectus](https://en.wikipedia.org/wiki/Homo_erectus)).

of catastroloess field, of considerable thickness, accumulated evenly in a sheet to all earth surface in differ local environments from tops of mountains to bottoms of the seas as constructive material burying most of the “Old World” surface. The catastroloess did varied from places to places concerning chiefly upon local environs where it rested on. Catastrophically, when it falls it was globally fallen down to differ qualitatively and quantitatively devastated and disaster surfaces. The process on the formation of the catastroloess needed to have precisely studied dynamically. Since it was previously globally known and was seriously interpreted with the strong conception for normal local geological windblown sands or flooded river sands, or ordinary loess, for more than a century in Thailand or abroad.

In fact, catastroloess was the accumulation of ejecta, impactites and previous erosion debris, but there were unknown of such considerable thickness from previous geological record. Ordinary loess does not quantitatively contains such products of ejecta, impactites, impact sands, impact glasses, microtektites and dust, but local or exotic erosion debris with/or without minor local glacial outwashes blown, traditionally in Europe, by lateral low atmospheric windblown or gusty storm.

#### 4.14 Tektites in Southeast Asia and *Homo erectus*

[Editor’s note: The predecessor species to our species *Homo sapiens* - *Homo erectus* in Java is now well studied and the tektites found in the *H.erectus* beds and associated volcanic horizons are well dated. There can be little doubt

that *Homo erectus* (a.k.a. *Pithecanthropus*, *Java Man*, *Peking Man* (Fig. 109)) witnessed the Australasian tektite event both in Yunnan and Java and would have been strongly affected by the event. Hou et al. (2000) have studied the convincingly unreworkeed tektites and also the stone tools from the Bose Basin in Yunnan which were almost certainly made by *Homo erectus* and conclude “..the presence of abundant charcoal and silicified wood fragments, detected during excavation and laboratory study, {are} in precisely the same sediments containing the tektites and stone artifacts. On the basis of the co-occurrence of these remains, we suggest that the Paleolithic artifacts of Bose signal a behavioral adaptation to an episode of woody plant burning and widespread forest destruction initiated by the tektite event, which exposed cobble outcrops throughout the basin.” and subsequent use of the cobbles by incoming *H.erectus* populations.

The following is a discussion of the stratigraphic position of *Homo erectus* in Java from Sibava (1993). “The Jetis fauna containing early Javanese *Homo erectus* is Early Middle Pleistocene according to Hooijer, Kurten and Kahlke (in Nilsson, 1983). Beds following the Trinil fauna possess a rich faunal assemblage with 440,000 to 600,000 years for basalt layers interbedded with marine deposits containing the Trinil fauna. Tektite dated about 700,000 years have been found some The late Cenozoic strata in the Sangiran area have been divided into an Upper and Lower Kalibeng Formation, followed by the Early Pleistocene Pucangan Formation, and the Early to Middle Pleistocene Kabuh and Notopuro Formations (Watanabe and Kadar, editors, 1985). Tektites represent another marker horizon, suitable for wider regional correlation. The tektites found in situ in the middle part of the Kabuh Formation in the Sangiran area of Java have been dated by the fission track method at  $0.71 \pm 0.09$  million years and 0.71 million years respectively (Itihara et al., 1985; Suzuki et al., 1985). The tektite horizon in the Sangiran area of Java is close to the base of the Middle Pleistocene, as the underlying Middle Tuff has also been dated at 0.78 million years. The position of stratigraphic horizons of hominid finds in the Sangiran area based on the fluorine method has been discussed by Matsu’ura (1985). This method confirms the stratigraphic assignment of *Homo erectus* finds to a horizon

between the Middle and Lower Tuff of the Kabuh Formation (Homo erectus VII and VIII). Homo erectus corresponds to the base of the Kabuh Formation. Following this concept, Homo erectus modjokertensis together with Homo erectus are accompanied by the Early Pleistocene Jetis fauna." (Sibava, 1993).

### **5. Impact craters, glasses and tektites possibly the same age as the Australasian strewn field impact(s).**

The Australasian Tektite Strewn Field (Ford, 1988; Glass, 1990; Glass and Pizzutto, 1994; Wasson, 1991; Wasson et al., 1995) may speculatively be divided into 3 impact fields (see below): Kazakhstan, Indochina and Australia-Tasmania fields. After thorough field work on tektites, examined tektite expedition from Vietnam (Izokh and An, 1988) and tektites from southeast China and Hainan, it was found that that Muong Nong type tektite or layered tektites are characteristic of the Indochina field and restricted to a relatively small area. On the other hand glass in Tasmania exhibited greater temperature towards Darwin Glass an impactite glass scattered asymmetrically about Darwin crater, Tasmania (Fudali and Ford, 1979; Howard, 2004) possibly from the coma part and ended the long journey landing of the unnamed comet to the Earth.

#### *5.1 Zhamanshin Crater and irghizites, Kazakhstan*

Further evidence that tektites can be produced by an impact lies in the discovery of tektite-like glass (irghizites) around the 14 km diameter Zhamanshin impact crater in Kazakhstan (Florensky and Dabizha, 1980). They have compositions and ages similar to the Australasian tektites, but many have a composite form consisting of numerous droplets welded together. Microirghizites show a greater range in composition than the irghizites. O'Keefe (1987) suggested that the Zhamanshin crater was not the source of the irghizites, but that the crater was produced by the impact of a large mass of glass of irghizite composition and that the irghizites are merely fragments of the impacting body. However, most authors believe that the impacting body was a meteorite and that the irghizites are tektite-like glasses produced by impact melting of the surface deposits during the impact that produced the

crater (e.g., Taylor and McLennan, 1979; Bouška et al., 1981; Glass et al., 1983). The glasses are not well dated but the Zhamanshin Crater was formed at  $900 \text{ ka} \pm 100 \text{ ka}$ .

#### *5.2 Darwin Crater and Darwin Glass, Tasmania, Australia*

Darwin Crater has been known for a long time since Ford (1988) but later little had been carried on and there was extremely limited information. Darwin Glass is an impact glass found in a strewn field near Mt Darwin, western Tasmania, and Australia. It has been dated at  $816 \pm 7 \text{ ka}$  by Ar-Ar methods. A 1.2 km circular depression, named Darwin Crater ( $42^{\circ}18.39'S$ ,  $145^{\circ}39.41'E$ ), has previously been suggested as the source crater for the glass. The structure sits in a remote valley in Siluro-Devonian (Eldon Group) quartzite and slate. Earlier geophysical investigations demonstrated that the structure is an almost circular sediment-filled basin.

Drill core intersected fine grained lacustrine sediments (-60m thick) overlying poorly sorted crater-fill deposits. The pre-lacustrine crater-fill stratigraphy comprises an uppermost polymict breccia (~40m thick) of angular quartz and country rock, which contains very rare (<<1%) glass fragments (Crater-fill Facies A). Beneath the polymict breccia facies, the drill core intersected monomict sandy breccias of angular quartz (Crater-fill Facies B), and a complicated package of deformed slates (Crater-fill Facies C). One core penetrated to a maximum depth of ~230m, at which point coherent slate was encountered. Quartz grains in the crater-fill samples contain abundant irregular fractures. In some of the most deformed quartz grains, sub-planar fractures define zones of alternating extinction. Kinked micas are also present. The deformation observed in the crater-fill facies is far greater than in rocks cropping out around the crater. However, diagnostic shock indicators (e.g. planar deformation features in quartz) are absent, preventing confirmation of an impact origin by petrographic analysis of crater-fill samples alone.

Geochemical analyses of the glass reveal two compositional groups. Group I is close to bulk average Darwin Glass and is highly variable in composition. its major element compositional range is:  $\text{SiO}_2$  (80.6-93.9%),  $\text{Al}_2\text{O}_3$  (3.1-10.6%),  $\text{TiO}_2$  (0.2-0.7%),  $\text{FeO}$  (0.8-4.2%),  $\text{MgO}$  (0.25-2.3%) and  $\text{K}_2\text{O}$  (0.7- 2.7%). Group I glass is

predominantly light green to dark green or white. Group 2 glass is almost always black. Group 2 has a lower average  $SiO_2$  (81.1%) content, and a decreased range in  $SiO_2$  (76.4-84.4) concentrations. Average  $Al_2O_3$  (8.2%) is also greater than in Group 1. Group 2 glass is also significantly enriched in FeO (+1.5%), MgO (+1.3 %) and Ni, Co and Cr relative to Group 1. Average Ni (416 ppm), Co (31 ppm), and Cr (162 ppm) concentrations in Group 2 glass are beyond the range expected in average sedimentary rocks. The remaining trace element data show affinity with typical upper crustal sediments including pronounced negative Eu anomalies ( $Eu/Eu = 0.48 - 0.66$ ) and LREE enrichment ( $La/Lu^* = 5.8 - 8.87$ ).

Sr and Nd isotope data indicate that a mixture of the Eldon Group lithologies (suspected target rocks) can form Darwin Glass. Mixing calculations using average Eldon Group compositions also successfully model the glass composition. Such models result in significant errors only for Ni, and to lesser extents CO, MgO, Cr and FeO in some Group 2 glass samples. Enrichments in these elements require an ultramafic contribution. However, mixing models using a component of Tasmanian dunite, pyroxenite, or lamprophyre fail to produce the required glass compositions, and can thus be ruled out as significant components of the target rock stratigraphy. The observed composition of Group 2 glass samples can only be explained by mixing with up to 9% of a chondrite or chondrite-like projectile. The distribution of projectile material in the glass is extremely heterogeneous, and the amount of this contribution is varied. Only the transition metals are enriched, with no simultaneous enrichment in the highly siderophile elements (HSE) that are present in crustal abundances.

Geochemical and isotopic data are considered to be consistent with Darwin Crater being the sole source of the glass. The crater-fill facies are also interpreted as consistent with impact processes. The argument for a genetic connection is strengthened by the observed trends in the distribution of glass relative to the crater. In the proposed model of the impact event, the polymict matrix supported breccia of Crater-fill Facies A is interpreted to have formed from non-melted angular quartz and country rock fragments that were blasted outwards and upward along the cavity floor, before collapsing




Fig. 110: Location of Darwin Crater in SW Tasmania, Australia. Historically glass was found in the state railway track to Kelly Basin. (Howard, 2004)



Fig. 111: Dense trees in the study area. (Howard, 2004)



Fig. 112: Darwin glass collected near Mt. McCall in 2002. Scale bar= 5 cm. (Howard, 2004).




Fig. 114: Irregular Darwin Glass, A-F) scale bar = 2cm. Note the parallel layering that is predominantly planar and common presence of vesicles best illustrated in Fig. 13.9B. These specimens have a superficial resemblance to the Australasian Strewn field. (Howard, 2004)



Fig. 113: Darwin Glass, A) Irregular and ropy glass fragments (scale bar 5 cm) size; B) Irregular white glass fragments (scale bar = 1 cm.); C) Splash-form Darwin Glass (scale bar = 2.5 cm.), D) Splash-form mini glasses (scale bar = 5 cm), (Howard, 2004)

inwards and mixing. Crater-fill Facies B and C are interpreted as representing shattered quartzite and plastically deformed slate (<5GPa), sourced from slumping of the cavity walls.

Physical trends in glass distribution relative to distance from the crater can also be used to test the genetic relationship between the strewn field, and show that the ejected melt cooled and rained down as glass fragments over more than 410km<sup>2</sup> of western Tasmania. In rare cases glass fragments exceeded 1 kg, but typically were only a few grams in size. In a 50 km<sup>2</sup> area surrounding the crater (~1/8th of the strewn field), and it is estimated that the total glass volume is at least 11,250 m<sup>3</sup> Therefore, relative to the size of the crater, Darwin Glass is the most abundant impact glass on Earth. The wide distribution and anomalously high abundance of glass in the strewn field is explained as relating to groundwater infiltration of the target rocks along fractures and faults prior to impact. Surface swamps are interpreted to have been present in the study area throughout the Pleistocene, and thus were a likely feature of the pre-impact environment. The abundance of water would have produced a highly volatile- charged target stratigraphy. This volatile enhancement is interpreted to have increased the explosiveness

of the impact, and the efficiency of melt dispersal and ejection. Australian tektites are very different to Darwin Glass as the following Figs 110-123 show.

Analyses of more than 4000 fragments of glass recovered in situ around the crater show:

- 1) the largest recovered fragments are found closest to the crater;
- 2) a decrease in the proportion of fine glass fragments away from the crater;
- 3) size distribution data for the recovered glass specimens are strongly skewed towards outlying large fragments;
- 4) an increase in the proportion of black glass away from the crater;
- 5) an increase in the proportion of splash-form, relative to irregular or ropey shapes away from the crater: and
- 6) splash-form shapes are preferentially black.

In the impact model, impact glass size distribution data are considered to be consistent with ballistic ejection of melt. The poor size sorting is interpreted to indicate that the ballistic ejection of melt from the crater was as a highly turbulent plume and that large and small fragments were deposited together, on the breakdown of turbulent cells in the plume. The increase in the proportion of black glass fragments with increasing distance from the crater is related to the depth of excavation. Black glass is interpreted to form from melting of pelitic layers in the Keel Quartzite, which is the uppermost target formation and it is expected that the uppermost target rocks that will be ejected farthest during impact.

Splash-form shapes form during aerial transport, increasing the distance of melt ejection will promote development of such shapes, and this is in turn consistent with the preference for splash-form shapes to be black. The expected lower viscosity of the black melt (based on  $\text{SiO}_2$  content) is also interpreted to have promoted the development of splash-form shapes. Deriving of black glass from the uppermost target rocks, close to the target-projectile interface, also aids in explaining the evidence for preferential projectile contamination of some black glass specimens.




Fig. 115: Irregular Darwin Glass, G-I) The largest known fragment of Darwin glass that weighs 946g after a slice has been removed for sectioning. The fragment was collected by R. J. Ford. This specimen also has a superficial resemblance to layered tektites of the Australasian strewn field. (Howard, 2004)

## 6. CONCLUSIONS: Pleistocene cometary impact in Thailand or nearby caused the Australasian tektite field; as suggested by catastrophic loess, mass mortality and forest fires

New discoveries in 1997 (Bunopas, S.; J. T. Wasson, Vella, P., H. Fontaine; S. Hada; C. Burrett Th. Suphajunya and S. Khositantont, 1999) added to previous research on tektites and impact craters in Thailand, which suggested that common continental fires; mass mortality of




Fig. 116: Internal features of Darwin Glass, A) Schlieren defining a wave or droplet shape (scale bar = 100µm; B) Close up of A (scale bar = 100µm). These features reflect turbulent circulation in the melt jet. C) Tail of another droplet shape feature (scale bar = 100µm); D) Close up of C (scale bar = 100µm). (Howard, 2004)




Fig. 117: Ropy Darwin Glass. A-F) scale bar = 2cm. Twisted ropy feature reflects the extreme topographic expression of the parallel layering characteristic of the irregular glasses. This texture reflects quenching from a rapidly stretching viscous melt. (Howard, 2004)

trees, mammals, and reptiles, as well as the presence of thick catastrophic loess, are linked in an extraordinary global catastrophic event. Evidence from these new discoveries have been superimposed on an area larger than the region known as the Australasian tektite strewn-field. Radiometric dating by various methods of tektites from this field gave ages between 0.709 - 0.770 ± 0.020 Ma. [788 ± 3 ka Jourdan et al., 2019] This date is probably consistent with related evidence of an early Quaternary event and possibly correlated to a glacial stage at O-isotope stage 20.2 (~NN20). 300 km west of the probable impact center in Ubon Ratchatani,



Fig. 118: Elongate Darwin Glass, A) scale bar = 1cm; B) scale bar = 2cm; C) scale bar = 2cm; D) scale bar = 1 cm; E) scale bar = 1 cm; F) scale bar = 1 cm; G) scale bar = 2cm. These are usually planar and rod-like but a= samples with bulbous end and also found along very rare and usually poorly developed dumbbells. Such elongate samples almost always have broken ends. (Howard, 2004)

sand pumping pits were investigated in Tha Chang and Chum Phuang; respectively 16 km east and 100 km northeast of Khorat in northeast Thailand. Pits >12 m were sufficiently deep to reveal irregular piles of partially to completely burnt and petrified logs, trunks, and trees, apparently pushed and pulled down abruptly to the ground by tremendous force.

Nearby at Tha Chang, two newer sandpits produced complete ancient elephant teeth and *Stegodon* sp. fossils of Pleistocene age were also found, distributed among other mammal bones, crocodile teeth and bones, as well as burnt trees in the black reduced beds and in the bottom of the pits.

This evidence suggests sudden ground melting resulting from an extra-terrestrial impact, contemporaneous with the formation of tektites in the Australasian field, though geochemical analyses have not indicated the addition of any




Fig. 119: Spheroid and droplet shaped Darwin Glass, A) scale bar = 1cm; B) scale bar = 1cm; C) scale bar = 0.5cm; D) scale bar = 0.5cm; E) scale bar = 1cm; F) scale bar = 2cm; G) scale = 2cm; H) scale bar = 2cm. These spherical glasses appear to have the most highly pitted surfaces of all of the glass shapes (e.g. B). These droplets are almost always asymmetric and typically have broken tails. Examples like F, G and H bear a very strong resemblance to splash-form tektites of the Australasian strewn field, particularly Thailandites. (Howard, 2004)

new elements or minerals. A rapid succession of enormous global explosions followed the proposed impact, causing global distribution of various tektites. Consequent continental forest fires and build-up of atmospheric sands (or dusty debris) caused great loss of life over a very large area.

Small, medium, and (rarely) large layered tektites are found near an impact centre, and less commonly outside. This evidence may substantiate the enormous terrestrial impact of a big comet, indicated by the considerable size on and large numbers of multiple craters in an 800 x 1140-km impact centre between Hainan and Cambodia.

The presence of microtektites and glass



Fig. 120: Colour variation of Darwin Glass, A) Typical white glass (scale bar = 1cm); B) Typical light green glass (scale bar = 2cm); C) Typical dark green glass (scale bar = 2cm); D) Typical black glass (scale bar = 2cm) (Howard, 2004).

microspherules in north Chinese loess confirms that late generation of loess enabled the collection of minute material suspended in the atmosphere. The catastroloess is related to both an early Pleistocene mass mortality event, as well as to the addition of thick sediments to normal local geological processes.

Incorporating the 1997 evidence from Thailand allows a new Comet theory to be proposed: The Cometary Impact Theory encompasses the effects of a large comet disintegrating and impacting on Earth, causing great global shocks and quakes, disastrous continental fires, and a mass mortality that affected much of the extant flora and fauna. Numerous violent impact explosions caused rapid exhaustion of atmospheric oxygen, which, in turn resulted in vast quantities of debris and dust being sucked up into the atmosphere. Subsequent cooler conditions on Earth were accompanied by the late steady fallout of dry earth-sands settling and forming thick and extensive catastroloess. A modern analogy for this catastrophe would be a country impacted by many hundreds of nuclear bombs, allowing resident communities little chance of survival.

Tektites may be found at any level in the sands that cover the basement. We contend the variation in this level reflects repositioning either during the catastrophe or as a result of geological




Fig. 121 : Geology of the Darwin Crater Strewn field. Based on Corbett and Brown (1975) and Corbett et al. (1993).

processes at some stage between the time of catastrophic events and the present day. The catastrophe should be known as the Buntharik Event, name derived from a town south of Ubon Ratchathani, Thailand where all evidence outlined in this study can be examined in a few hours. The Buntharik Event which is responsible for the studied tektites, also caused the outbreak of global continental fires, mass extinction flora and fauna, and the development of a catastroloess that dramatically altered the lower Quaternary landscape. The impact may also have triggered vast basaltic volcanism nearby (Ford, 1988) and facilitates demarcation of the lower-middle Pleistocene boundary in Southeast Asia




Fig. 122: Darwin Crater geology. Based on field mapping in this study and by R.J. Ford; 1: 25 000 aerial photographs; Corbett and Brown (1975) and Corbett et al. (1993).



Fig. 123: Physiography of Mount Darwin western Tasmania, Australia. (Howard, 2004).

(Bunopas et al., 1999a; Sibrava, 1993 etc.) and China (Han et al., 1997; Wang et al., 1985). Dating of tektites reveals the age of the event between 0.709 - 0.770 ±0.020 Ma (Blum et al., 1992; Gentner et al., 1967; 1969; Izett an Obradovich, 1992 and Zahringer, 1963). [788 ± 3 ka Jourdan et al., 2019]. The global disaster contributes to the theory of extra-terrestrial tectonics and allows an understanding of the origin of earth's common exotic loess sands. We contend that such sands are not the normal weathering

products in an arid region, but rather represent reworked catastroloess.

In final conclusion, such a catastrophe may comprise occurrences in the following categories:

1) **Numerous extensive burnt felled trees**, all root up and most are petrified, lower levels contain abundant soot and sulfides. Uprooted tree trunks up to 2 metres in diameter are often shattered, branchless, snapped, and burnt to the core. Their presence in deposits throughout Thailand is evidence of widespread burning. These suggested extensive wild-forest-fire which were reported bunt trees or woods the whole of Southeast Asia.

2) **Ancient animals** such as Stegodon turtles, crocodiles etc. (Chaimanee, 2004), all suggesting a mass mortality event.

3) **Tektites** are rare splashed tektites from the pits but many layered tektites in Ubon Ratchatani, 233 km east.

4) **Impact crater** in Buntharik though it is still dubious.

5) **Catastrophic dusty sands** represent debris and ejecta that formed widespread sheet enclosed other DCIMs. This name is proposed "Catastroloess", for this impact sands washed and catastrophic flooding as sands at Ban Tha Chang, Howard (1999) added.

6) **Microtektites, glasses, impact quartz and impact glasses etc.** (Howard, 1999)

7) **Isotopic elements from INAA, ICP-MS measurements**

in tektites from Thailand, China and recent study from Ban Tha Chang sands (Howard, 1999).

8) **Reversal paleomagnetism with fragile (unreworked) tektite** (Haines et al., 2004).

9) **Mass Mortality Event.** The great impact mass killed numerous ancient vertebrate and knocked down and burnt previous flourishing forest trees. Bunopas et al. (1999, 2007).

Evidence constituted by these criteria supports the theory of a cometary impact during the early Quaternary. We believe that evidence of the Buntharik Event, including phenomena of tektites, impact craters, burnt woody fossil material, and catastroloess all further our

understanding of mass mortality events and possibly extinctions in general.

The impact occurred at almost the end of a Lower Pleistocene time (Q1). The impact surface is marked by ground forested layer gravels of latest Lower Pleistocene, continued by sandy impact debris belonging mainly to the early parts of Middle Pleistocene, consistent with the stratigraphic age and the age of the tektites (latest Q1 and mainly Q2). An analysis of the Quaternary stratigraphy indicates that all badlands that make a good touring place, and mega-flooded BTC sands at Khorat, and above the fish beds at Lomsak, Pha Wing Choo, south Chiangmai and faulted sand beds at Na Thawi in Songkhla, the Southern Peninsula etc., are marked as Middle Pleistocene age.

Continuation of the regional uplifting had resulted in considerable thickness of common covering or capping gravels and late raising of impact layer to a higher level. All gravels under the "muddy gravels" on the top of lower terrace gravels mark the 770 ka "avalanches and mudflows mixed with the earliest onset of catastroloess sands" surface. The gravels must be former impact forestry surface. These "muddy gravels" buried felled trees at Krok Duan Ha, Khorat and also at school yard, Ban Non Thung, Buntharik, containing rare splash and abundant un-oriented, small to medium layer tektites, respectively, and are correlated




Fig. 124: Projection of Earth Showing Host Continents, Asia and Australia When they were Impacted by the Australasian Comet at Kazakhstan with irghizites at Zhamanshin structure, south China and Indochina (South-East Asia), and Australia with tektites, then collided, terrestrially to Mount Darwin making the Darwin Crater in Tasmania with Darwin glass. DCIMs, most are misinterpreted, and are newly known in Thailand, Vietnam, China, Australia and Antarctica etc.



Fig. 125: Proposed track of the 0.788 Ma comet. 1. Zhamanshin (Kazakhstan), 2. Indochina, 3. Australia including Tasmania, Darwin Crater.



Fig. 126: The icy missiles with solid cores made fantastic Jupiter impact in 1994 (in this Fig. it is represented by Halley's Comet). The coma of the comet is the halo or nebulous envelope of gas, ice and rock bursting from the rocky-ice-gas nucleus at the head of the comet.

to Bo Phloi, Kanchanaburi, on 23m thick older gravels with sapphires.

All terrace gravels in most regions of Thailand are dated as Lower Pleistocene, the tectonic uplifting stage. Lower Pleistocene includes also high terrace and lower terrace gravels under Ban Tha Chang Sands (Howard et al., 1999) and river sands in the Mun and Chi rivers. In the Khorat plateau "old gravels" were thought to be below the tektites. The top of the gravel is represented by old tree growth and form the disastrous surfaces during the impact with original first generation tektites fine sandy-silty and lateritic layer with tektite in Ubon Ratchatani (Fiske et al., 1996) is only a few metres above the avalanche position in a locality few km to the south. Finally, the horizon produces high atmospheric explosion size frequency pattern

and consistent REE pattern similar to those for tektites (Wasson, 1991) derived from siliceous sediments in previous impact surface.

The Buntharik Event is proposed for the event that produced tektites in the Australasian field covering 1/10, and the catastroloess covering much of the Earth's surface (Bunopas et al., 1998). The catastroloess admixed to those local vital normal conditions in mountainous valleys; rejuvenated the current physiography of all valley and flood plain areas; and made the extinction of many older plants and in lowlands or floodplain areas or paddy fields everywhere in Thailand contain some thickness of catastroloess or/and the secondary enriched equivalent sands several metres thick. This is applied also to the floors of the Gulf of Thailand that were filled with these sands.

The 0.788 Ma impact event was likely caused by the disintegration of a comet and the impact of its fragments, first in Kazakhstan to form the Zhamanshin Crater, then in Indochina perhaps forming numerous small craters and then in western Tasmania to form the Darwin Crater (Figs. 124-126). If this model is correct, then Thailand contains unique evidence for the terrestrial impact and the catastrophic effects of a large extraterrestrial object.

### SUGGESTED FURTHER READING

- Barnes, V.E. and Pitakpaivan, K. 1962: Origin of Indochinese tektites. *Proceedings of the National Academy of Science USA*. 48, 947-955.
- Chaimanee, Yaovaluk, 2004. Fossils from Khok Soong and Ban Tha Chang, Khorat. In: *Numerous fossils in Paleontology of Thailand, in 114 years of the Department of Mineral Resources.* (in Thai)
- Duangkrayom, J., Ratanasthien, B., Jintasakul P., Carling, P.A. 2014. Sedimentary facies and paleoenvironment of a Pleistocene fossil site in Nakhon Ratchasima province, northeastern Thailand. *Quaternary International* 325: 220-238. doi: 10.1016/j.quaint.2013.07.048
- Glass, B.P., Folco, P., and Campanale, F., 2020 Coesite in a Muong Nong-type tektite from Muong Phin, Laos: Description, formation, and survival. *Meteoritics & Planetary Science* 1-21, doi: 10.1111/maps.13433
- Haines, P. W., K. T. Howard, J. R. Ali, C. F. Burrett, Sangad Bunopas, 2004. Flood deposits penecontemporaneous with 0.8 Ma tektite fall in NE Thailand: impact-induced environmental effects? *Earth and Planetary Science Letters* 225, 19-28.
- Howard, K. T. 2004: Origin of Darwin Glass. Unpublished Ph. D. Thesis, University of Tasmania, Australia. 362 pp
- Howard, K.T., 2011. Tektites in: Ridd, M.F., Barber, A.J., Crow, M.J. (Eds.), *The Geology of Thailand*, The Geological Society of London, pp. 574-591.

DOI:<https://doi.org/10.1144/GOTH.21>

- Howard, K. T., P.W. Haines. C. F. Burrett, J. R. Ali, S. Bunopas, 2003. Sedimentology of 0.8 Ma log-bearing flood deposits in northeast Thailand and mechanisms for pre-flood deforestation, Proceedings, 8th International Congress on Pacific Neogene Stratigraphy, Chang Mai, Thailand, 2003, pp. 49-67.
- Koeberl C., 1994. Tektite origin by hypervelocity asteroid or cometary impact: Target rocks, source craters and mechanism. In Large Meteorite Impact and Planetary Evolution (eds. B. O. Dressler, R. A. F. Grieve and V. L. Sharpton), pp. 133-151. Geol. Soc. Am. Spec. Pap. 293, Boulder, Colorado.
- McCall, G.J.H. 2001. Tektites in the Geological Record: Showers of Glass from the Sky.: London (The Geological Society), 2001, viii + 256 pp. ISBN 1 86239 085 1.
- Nichol, J.E., Nichol, D. W., 2015. Character and provenance of aeolian sediments in northeast Thailand. *Aeolian Research*, 19, 5-14. <https://doi.org/10.1016/j.aeolia.2015.09.001>
- Sieh, S., Herrin, J., Jicha, B., Schonwalder, D., Moore, J., Banerjee, P., Wiwegwin, W., Sihavong, V., Singer, B., Chualaowanich T., and Charusiri, P., 2020. Australasian impact crater buried under the Bolaven volcanic field, Southern Laos. Proceedings of the National Academy of Science, 117 (3) 1346-1353; <https://doi.org/10.1073/pnas.1904368116>
- Suraprasit K, Jaeger J-J, Chaimanee Y, Chavasseau O, Yamee C, Tian P, Panha S. 2016. The Middle Pleistocene vertebrate fauna from Khok Sung (Nakhon Ratchasima, Thailand): biochronological and paleobiogeographical implications. *ZooKeys* 613: 1-157. doi: 10.3897/zookeys.613.83
- Wasson, J. T., 1985. Meteorites: Their Record of Early Solar System History, Freeman, New York.
- Wasson, J. T., Pitakpaivan, K., Putthapiban, P., Salyapongse, S., Thathimthong, B., and McHorne, J. F. 1995. Field recovery of layered tektites in northeast Thailand: Evidence of a large scale melted sheet. *J. Geophys. Res.*, 100. 14385-14390.
- Yang, F.-C., & Grote P.J., 2018. Riverine vegetation and environments of a Late Pleistocene river terrace, Khorat Plateau, Southeast Asia. *Palynology*, 42:2, 158-167, doi: 10.1080/01916122.2017.1296044
- and the Himalaya and their implications on Gondwana and Cathaysia/Laurasia continent configurations. *Geology and mineral resources of Southeast Asia* (ed. Prinya Nutalaya), pp. 121-7. Asian Instit. Technology, Bangkok.
- Adam, D. P., 1988. Developing for a climate record for the past three million years. United States Geological Survey Year Book Fiscal Year 1988. United States Printing Office, 1989. 134 pp., 50 - 52.
- Adams, E. W. and Huffaker, R. M., 1964. Aerodynamic analysis of the tektite problem. *Geochim. Cosmochim. Acta* 28, 881-892.
- Adinegoro, U. and Hartoyo, P. (1974). Paleogeography of northeast Sumatra. Proceedings of the Indonesian Petroleum Association Third Annual Conuention, Jakarta, June 1974, 45-52.
- Aghassi, B. 1962. *Astro. Contrib. Boston Univ.*, ser II, No. 6, 8 pp.
- Ahrendt, H., C. Chonglakmani, B. T. Hansen, and D. Helmcke 1993. Geochronological cross section through northern Thailand, *J. SE Asia Earth Sc.*, 8, 207-2 17, 1993.
- Alderson, A; Holmes, M. A. and Murphy, C., 1994. A summary of the biostratigraphy and biofacies for the Tertiary sequence, Gulf of Thailand. Proc.of the Intern. Symp.on Stratigraphic Correlation of SE Asia. 15-20 Nov. 1994, Bangkok, Thailand.
- Aldiss, D. T. and Ghazali, S. A. (1934). The regional geology and evolution of the Toba volcano-dipression, Indonesia. *Journal of the Geological Society London* 141,487-500.
- Allen, C. R. 1952. Circum-pacific faulting in the Philippines -Taiwan region. *Journal of Geophysic Research* 67,4795-812.
- Allen, C. R., Gillespie, A. R., Han Yuan, Sieh, K. E., Zhang Buchun, and Zhu Chengna. 1984. Red River and associated faults Yunnan Province, China: Quaternary geology, slip rates and seismic hazard- Geological Society of America Bulletin 95. 686-700.
- Alexander, J. B. (1958). The geology and mineral resources of the neighbourhood of Bentong, Pahang, and adjoining porrions of Selangor and Negri Sembilan. *Memoirs of the Geological Survey Department West Malaysia* 8, Ipoh.
- Alexseev, M. N. and Takaya, Y., 1967. An outline of the Upper Cenozoic deposits in the Chao Phraya basin, Central Thailand. *Southeast Asia Studies*. V. 5, no. 2, pp. 106-124
- Alvarez, W., 1986. Towards a theory of impact crises. *EOS Trans. AGU*, 67 (no. 35) 649, 653-655. 658.
- Alvarez, W., Alvarez, L. W., Asaro, F. and Michel, H. V., 1979, 1979a. Experimental evidence in support of an extra-terrestrial trigger for the Cretaceous-Tertiary extinctions (abstract), *EOS Trans. AGU*, 60, 734.
- Alvarez, W., Alvarez, L. W., Asaro, F. and Michel, H. V., 1979b. Anomalous iridium levels at the Cretaceous-Tertiary boundary at Gubbio, Italy: Negative tests for a super nova origin. In T. Birkelund (ed.) *Cretaceous-Tertiary Boundary Events Symposium*, 2, 69, University of Copenhagen, Copenhagen.
- Alvarez, L. W., Alvarez, W., Asaro, F. and Michel, H. V., 1980. Extraterrestrial cause for Cretaceous-Tertiary extinction. *Science*, 206, 1095-1107.
- Anderson, R.N., 1980. Update of heat flow in the East and Southeast Asian seas and islands. *Am, Geophys. Union Geophys. Monogr. Ser.*, 23, 319-326.

## REFERENCES AND BIBLIOGRAPHY

- Abbott, M. J., and Chamalaun, F. H. 1981. Geochronology of some Banda Arc volcanics. *Indonesia Geological Research and Development Centre Special Publication* 2, 253-68.
- Achache, J. and Courtillot, V. (1987). A preliminary Upper Triassic paleomagnetic pole for the Khorat Plateau (Thailand): consequences for the accretion of Indochina against Eurasia. *Earth and planetary Science Letters* 73, 147-57.
- Achache, J., Courtillot, V., and Besse, J. 1983. Paleomagnetic constraints on the Late Cretaceous and Cenozoic tectonics of Southeast Asia. *Earth and Planetary Science Letters* 63, 123-136.
- Achache, J., Courtillot, Y., Zhou, Y. X., Lu, L. Z., and Yuan, X. G. (1982). The paleolatitude and geographic extent of southern Tibet in Middle Cretaceous-new palaeomagnetic data. *Palaeomagnetic research in Southeast and East Asia*, C. C. O. P/TP 13, Bangkok, 47-57.
- Acharyya, S. K. (1978). Mobile belts of the Burma-Malaya

- Armijo, R., P. Tapponnier, J. L. Mereier, and H. Tonglin 1986. Quaternary extension in southern Tibet : field observations and tectonic implications, *J. Geophys. Res.*, 91, 13,803-13, 872, 1986.
- Armijo, R., P. Tapponnier, and H. Tonglin, 1989. Late Cenozoic right-lateral strike-slip faulting in southern Tibet, *J. Geophys. Res.*, 94, 2787-2838, 1989.
- An, Z. S., Wang, J. D. & Li, H.M., 1977. Paleomagnetic research of the Luchuan loess section. *Geochimica* 4, 239-249. (in Chinese with English abstract)
- Anderson, J. B.. 1983: Ancient glacial-marine deposits: Their spacial and temporal distribution. In Molnia, B.E (ed.): *Glacio-marine sedimentation*. Plenum Press, New York, p. 3-92.
- Angier, N. Reported by R. Burderi, B. and C.Gorman, 1985. Did Comets Kill the Dinosaurs? *Science Section, TIME*, May, 1985, 50-61.
- Archibald, J. D. and Clemens, W. A., 1982. Late Cretaceous extinction. *Amer. Sci.* 70, 377-385.
- Arpandi, D. and Patmosukismo, S. (1975). The Cibulakan Formation as one of the mosr prospective stratigraphic units in the northwest Java bisinal area. *Proceedings of the Indonesian petroleum Association I*, 1,81-211.
- Aitchison, J. C. and A. M. Davis, 2001. When did the India - Asia Collision Really Happen? *Gondwana Research*, 4 (4) 2001: 460-461.
- Asaro, F., Alvarez, L., Alvarez, W. and Michfi, Fl. 1982. Geochemical anomalies near the Focene/Oligocene and Permian - Triassic boundaries. *GSA Spec. Paper* 190, 517-528.
- ASCOPE (1981). Tertiary sedimentary basins of the Gulf of Thailand and South China Sea: stratigraphy, structure and hydrocarbon occurrences. ASEAN Council on Petroleum, Jakarta.
- Audley-Charles, M. G., 1987. Dispersal of Gondwanaland: relevance to evolution of the angiosperms. In: *Biogeographical Evolution of the Malay Archipelago*. (Ed. T. C. Whitmore.) 5-25. (Clarendon Press: Oxford.)
- Audley-Chartes, M. G., 1988, Evolution of the southern margin of Tethys (North Australia region) from the Early Permian to Late Cretaceous, in Audley-Charles, M., and Hallam, A., eds., *Gondwana and Tethys: Geological Society of London Special Publication* 37, p. 79-100.
- Audley-Charles, M. G., Carter, D. J., and Milsom, J. S. (1972). Tectonic development of eastern Indonesia in relation to Gondwanaland dispersal. *Nature, London (Physical Science)* 239, 35-9.
- Aylmer, D., M. Guyton. G.F. Herzog, B.M. Clement, i. Klein and R. Middleton. Beryllium-IO contents of sediments associated with the Matuyama- Brunhes geomagnetic reversal (abstract). *Trans. Am. Geophys. Union* 69, 34t. 1988.
- Baker, G. 1958. The role of aerodynamical phenomena in shaping and sculpturing Australian tektites. *American Journal of Science* 256, 369-383.
- Baker, O. 1967. Structure of well preserved Australite buttons from Port Campbell, Victoria. Australia. *Meteoritics* 3, 179-217.
- Baldwin, R. B. 1949. *The Face of the Moon*. University of Chicago Press, Chicago.
- Baldwin, R. B. 1963. *The Measure of the Moon*. University of Chicago Press, Chicago.
- Banner, F. T., and W. H. Blow, 1967. The oigin, evolution and taxonomy of the foraminiferal genuus *Pullenialatina* Cushman, 1927. *Micropaleontology*, v/ 13, pp, 133.
- Barber, A.J. 1995. Late Permian ans Mesozoic Statigraphy of Timor, Eastern Indonesia. *Journal of Geology*, special issue, Series B, no. 5-6 (1995), p. 153.
- Barber,.A. J. 1981. Strucural interpretarions of the island of Timor, eastern Indonesia. *The geology and tectonics of eastern Indonesia*, Geological Research and\_ Development Centre, Special publication 2 (ed. A. J. Barber and S. Wiryosujono) , pp” 1g3\_97. Bandung.
- Barber, A. J. and Wiryosujono, S. (eds.), 1951. *Tbe geology and tectonics of eastern Indonesia*. Geological Research and Developmenr Cenrre, Special publication 2. Bandung.
- Barbour, G.B., 1930. The Paote and Chingshui physiographic stage and their significance. *Bull. Geol. Soc. China* 9, 45-48.
- Barkatt, A. A., Saad, E. E., Adiga, R., Sousanpour, W., Barkatt, Al. and Alterescu, S. 1986. Leaching of microtektite glass compositions in seawater. *Advace in Ceram.* 20: Nucl. Waste Manage. 11, 6.1-687.
- Barnes, V. E., 1961. *Scientific Amercan*, 205 (5), p, 56-65.
- Barnes, V. E., 1963. Detrital mineral grains in tektites. *Science* 142, 1651-1652.
- Barnes, V. E., 1964. Variation of petrographical and chemical characteristics of indochinite -tektites within the strewn field. *Geochim. Cosmochim. Acta* 28, 893-913.
- Barnes, V. E. 1967. Tektites. *International Dictionary of geophysics* 2. 1507-1518.
- Barnes, V. E. 1969. Progress of tektite studies in China. *Trans. Am. Geophysic. Union.* 50. 704-709.
- Barnes, V. E., 1989. Origin of Tektites. *Tex. J. Sci.* 41, 5- 33.
- Barnes, V. E. 1971. Description and origin of 12.8-kg layered tektite from Thailand (abstract). *Meteoritics* 6 (4), 249
- Barnes, V.E. and Pitakpaivan, K. 1962: Origin of Indochinite tektites. *Proc. Nat. Acad. Sci. USA* 48: 947-955.
- Barnes, V. E. Description and Origin of Large Tektite from Thailand with comment]. (in Tektites. p. 392-398, illus., DOWDEN. HUTCHINSON & ROSS. Stroudsburg. 1973) Barnes. Virgil E. Description and origin of 12.8-kg layered tektite from Thailand [abstr.]. (METEORITICS. Vol. 6, No. 4, p.249. 19711
- Barnes, Virgii E. Description and Origin of Large Tektite from Thailand fwith comment]. (in Tektites. p. 392-398, illus., DOWDEN, HUTCHINSON & ROSS. Stroudsburg- 1973)
- Barnes, Virgil E. 1971. Description and Origin of Large Tektite from Thailand. (CHEM. ERDE. Vol.30, No. 1-4, p. 13-19, illus.)
- Barnes. Virgil E. Description and origin of 12.8-kg layered tektite from Thaiiand [abstr.]. (METEORITICS. Vol. 6. No. 4. p. 2-19. 1971.
- Barnes, Virgil E. Tektites; a world-wide geological investigation. (GEO TIMES (AM. GEOL. INST.). v.6, no.2. p.8-i2’ 38, iilus.; addendum, no’ 4, p’37’ 1961)
- Barr, SM., and MacDonald, A.S., 1978. Geochemistry and petrogenesis of Late Cenozoic alkaline basalts of Thai/and. *Bull. Geol. Soc. Malaysia*, 10, 25-53.
- Barr, SM., and MacDonald, A.S., 1981. Geochemistry and geochronology of late Cenozoic basalts of southeast Asia. *Geol. Soc. Am. Bull.* 92, 1069-1142.
- Batygin, K., and Brown, M., 2016. Evidence for a distant giant planet in the Solar System. *The Astronomical Journal*. 2016, 151(22), 1-12. Doi: 10. 3847/0004-6256/ 151/2/22.
- Benammi, M., Chaimanee, Y., Jaeger, J.-J., Suteethom, V. & Ducrocq, S., 2001. Eocene Krabi basin (southern

- Thailand): Paleontology and magnetostratigraphy. *GSA Bull.* 113 (2), 265-273.
- Benammi, M., Urrutia-Fucugauchi, J., Alva-Valdivia, L. M., Chaimanee, Y., Triamwichanon, S., and Jaeger, i-i., 2002. Magnetostratigraphy of the Middle Miocene continental sedimentary sequences of the Mae Moh basin in northern Thailand: evidence for counterclockwise block rotation. *Earth and Planetary Science Letters* 204, 373-383.
- Benainini, M., Chaimanee, Y., Urrutia-Fucugauchi, J., and Jaeger, J.-J., 2004. Magnetostratigraphy study of the continental sedimentary sequence of the Chiang Muan basin, Northern Thailand; Implication for the age of the first Miocene Hominoids from Thailand. *International Geology Review* 46. 646-654.
- Berger, W. H. and Heath., 1968. 'Vertical mixing in pelagic sediments', *Journal of Marine Research*, 26, 134-143.
- Berry, R. F. and Grady, A. E. 1987. Mesoscopic structures produced by Plio-Pleistocene wrench faulting in South Sulawesi, Indonesia. *Journal of Structural Geology* 9, 563-71.
- Berry, R. F., and McDougall, I. 1986. Interpretation of <sup>40</sup>Ar / <sup>39</sup>Ar and KAr dating evidence from the Aileu Formation, Last Timor, Indonesia. *Chemical Geology (Isotope Geoscience section)* 59, 43-58.
- Bhongsuwan, T., 1993. Paleomagnetic Investigation in Thailand. Lulia University of Technology. Lulia, Sweden. Ph. D. Thesis.
- Bignell, J. D. and Snelling, N. J. 0977a. Geochronology of Malayan garnets.- Overseas' geology and mineral resources 47, Institute of Geological Sciences, London.
- Bignell, J. O. 1d Snelling, N. J. 1977b). K:Ar ages on some basic igneous rocks from peninsular Malaysia and Thailand. *Geological Society of Malaysia Bulletin* 8,89-93.
- Bird, E. C F. 1965. A geomorphological study of the Gippsland Lakes. Publ. Dep. Geogr. Aust.natn Uni. Canberra G/I
- Birkeland, P. E. 1984. Soil and Geomorphology. Oxford, New York, 372 pp.
- Bohor, B. F. and Glass B. P. 1995 Origin and diagenesis of K/T impact spherules from Haiti to Wyoming and beyond. *Meteoritics* 30, 182-198.
- Boillinger, W. and de Ruiter, P. A. C. (1976). Geology of the south central Java offshore area. Proceeding's, of the 4th annual meeting of the Indonesian petroleum Association, Jakarta 5, 67 -8 I.
- Borax, E. and Stewart, R. D. (19dd). Note on the Paleozoic stratigraphy of northeastern Thailand. Report for ECAFE meeting in Bangkok, 1965 mimeograph.
- Bostrom, R. C. 1978. Motion of the pacific plate and formation of marginal basins: asymmetric flow induction. *Journal of Physics of the Earth* 26 Supplement, S103-22.
- Bowin, C., et al. 1980. Arc-continent collision in the Banda Sea region. *American Association of Petroleum Geologists Bulletin* G4, 1 68-9 I S
- Blaine Cecil, C. B., 1988. Modern analogs of coal formation. *United States Geological Survey Year Book Fiscal Year 1988*. United States Printing Office, 1989. 134 pp.,81-84.
- Blum, J. D., Papamastassiou, D. A., Koerberl, C. and Wasserberg, G., 1992. Neodymium and Strontium Isotope Study of the Australasian tektites: New constraints on the provenance and age of target materials. *Geochim. Cosmochim. Acta* 56, 483-492.
- Bohor, B. F., Betterton, W. J., and Ford, E. E., 1988, Coesite, glass, and shocked quartz and feldspar at DSDP Site 612: Evidence for nearby impact in the late Eocene (abs.): *Lunar and Planetary Science XIX*, p. 114 115.
- Bollinger, K., 1993. <sup>40</sup>Ar-<sup>39</sup>Ar Datierung von Tektiten. Unpublished Diploma Thesis, Max Planck Institute, Heidelberg, 84pp.
- Boonsener, M., 1983. The stratigraphy of Quaternary deposit in Khon Kaen Province, Northeastern Thailand. In *Conference of Geology and Mineral Resources of Thailand*, Bangkok, 1-5.
- Boonsener, M., 1985 Review on Quaternary Geology in Northeastern, Thailand. *Proceedings of Conference on Geology and Mineral Resources Development of the Northeast, Thailand*. Department of Mineral Resources, Bangkok..
- Boonsener, M., 1991. The Quaternary stratigraphy of Northeast Thailand. *Journal of Thai Geosciences*, 1: 23-32
- Boreham, A. U. E., 1963. Some problems concerning the application of the Lower Nukumaruan (Hautawan) Substage (Pleistocene, New Zealand). *N.Z. J. Geol. Geophys.* 6, 3-27.
- Boudka. V. V., Povondra. P., Florensky, P.V. and Randa. Z.. 1981. Irghezites and Zhamanshinites: Zhamanshin Crater. *USSR. Meteoritics*. 16. 171-184
- Bouska, V., Povondra, P., Floresnky, P. V. & Randa, Z. 1981. Irghezites and zhamanshinites:Zhamanshin Crater, U.S.S.R. *Meteoritics* 16, 171-184.
- Bender, F., 1983, *Geology of Buma*: Berlin, Borntraeger,293 p.
- Bretz, J.H., 1923. The channeled scablands of the Colorado Plateau. *Geology*, 31, 617-49
- Briaes, A., 1989. Climatique d'ouverture de la Mer de Chine du Sud (Nan Hai). Implications pour la tectonique tertiaire de l'Asie, Doctorat thesis Thesis, 239 pp., UniversitParis 6, Paris, Nov. 1989.
- Briaes, A., P. Patriat, and P. Tapponnier, 1993. Updated interpretation of magnetic anomalies and seafloor spreading stages in the South China Sea, implications for the Tertiary tectonics of SE Asia, *J. Geophys. Res.*, 98, 6299-6..
- Brothers, R. N. 1954. The relative Pleistocene chronology of the South Kaipapa District, New Zealand. *Trans Roy. Soc. N. Z.* 82, 677-94.
- Brown, D. A., Campbell, K. S. W., and K. A. W. Crook. 1969. *The Geological of Australia and New Zealand*. Pergamon Press, Sydney, London.
- Brown, A. G., 1985, Traditional and multivariate techniques in the interpretation of, floodplain sediment grain size variations: *Earth Surface Processes and Landforms*, v.70, p. 281-291.
- Brown, G. F., et al. 1951. Geologic reconnaissance of the mineral deposits of Thailand. *U.S. Geological Survey Bulletin* 984.
- Browne, W. R., 1957. Pleistocene glaciation in the Commonwealth of Australia. *J. Glacio.* 3 (22), 111-15.
- Brunnschweiler, R. O. (1966). On the geology of the Indoburman ranges. *Journal of the Geological Society of Australia* 13,137-94.
- Brunnschweiler, R. O. 1970. Contributions to the post-Silurian geology of Burma (northern Shan State and Karen State). *Journal of the Geological Society of Australia* 17 , 59-79.
- Brunnschweiler, R. O. 1974. Indoburman ranges.

- Mesozoic-Cenozoic orogenic belts, Geological-Society of London Special Publication 4 (ed. A. M. Spencer), pp.279-99
- Buffetaut, E., 1984. Selective extinction and terminal Cretaceous event. *Nature*, London, 310, 276.
- Buffetaut, E., R. Helmcke-Ingavat, J.-J. Jaeger, Y. Jongkanchanasoontorn, and V Suteethorn. 1988. Mastocene remains from the Mae Teep Basin and their biostratigraphic significance. *C. r. Acad Sci Paris Serie II*, 245-254.
- Buffetaut, E., Suteethorn, V. and Khansubha, S. 2007. The Ceratosian Dinosaur Psittacosaurus in the Early Cretaceous of Southeast Asia: A Review of Old and Recent Finds. (GEOETHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. Geothai 2007, pp. 338-343. 21-22 November. Imeral Hotel, Bangkok, Thailand
- Bunopas, S., 1966. Upper Miocene – Pleistocene Stratigraphy and the Paleontology of Gladstone, Wararapa New Zealand. Unpublished M. Sc.Thesis at the Geology Department, Victoria University of Wellington, New Zealand.
- Bunopas, S., 1981. Paleogeographic history of western Thailand and adjacent parts of Southeast Asia - a plate tectonics interpretation, Ph.D Thesis, 810 pp., University of Wellington, New Zealand, 1981. -*Geol. Surv. Pap.*, 5 (spec. Iss.): 810 pp., 164 figs., 21 tabs.; Bangkok. [Typescript]
- Bunopas, S., and P. Vella, 1983. Opening of the Gulf of Thailand - rifting of continental Southeast Asia, and late Cenozoic Tectonics, *J. Geol. Soc. Thailand*, 6, 1-12, 1983b.
- Bunopas, S. and Vella, P., 1984. Phuket-Kaeng Krachan Group, A rifted continental margin deposit with effect of ice-rafting from Gondwana. Abstr. paper presented at GEOSEA V., 9-13 April, Kuala Lumpur.
- Bunopas, S. and Vella, P., 1989. "Pebbly mudstones", A composited assemblages of ice-rafted sediments on the western rifted marginal basin of Shan-Thai while drifting away from Gondwana during the Upper Paleozoic. Proceedings of the 4th Intern. Symp. On Pre-Jurassic East Asia, IGCP Project 224. Hat Yai and Langkawi Island, Thailand and Malaysia, 24 April-2 May 1989
- Bunopas, S., 1989. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Pacific Science Congress, Abstr. Seul, Korea.
- Bunopas, S., 1990. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Proc. of the 16th Conf. on Sciences and Technology of Thailand, 25-27 October, 1990, Bangkok. p. 512-513.
- Bunopas, S., 1992. Tektites - Their origin and the Early Quaternary catastrophic destruction in Northeastern Thailand and south Indochina. Proc. 29th Intern. Geol. Congress. 24 Aug.- 3 Sept. 1992. Kyoto, Japan, Abstr. Vol. 3(3), II-12-4 0-8 2114, p. 650.
- Bunopas, S.; Kositanont, S. and Wasson, J. T., 1997. Evidence of the Early Global Disaster and Destruction from Extraterrestrial Impacts Comet in Northeast Thailand and South Indochina within the Australian Tektites Field: The last Mass Extinction, (extended abstract). Proc. of the national conf. on stratigraphy and tectonics evolution of Southeast Asia and the South Pacific. 19-24 Augt. 1997, Bangkok, Thailand, 434-435.
- Bunopas, S and Kositanont, S. 1999: Geology and National Development in accordance with the King's new theory for self sufficient agriculture as viewed from the Buntharik event (in Thai). In: The Special Memorial Book for the King's 72 Birthday in 2542 (1999), contribution made to the King, "The Geographer of The Nation" by The Rajabhat Institute Buriram: 3 pp.
- Bunopas, S.; Wasson, J. T.; Vella, P.; Fontaine, H.; Hada, S.; Burrett, C.; Suphajanya, Th. and Kositanont, S. 1999a: catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. *Journ. Geol. Soc. of Thailand* 1999 (1): 1-17 2000, October 28-29 1999, Bangkok, Thailand: 142-151.
- Bunopas, S.; Wasson, J. T.; Vella, P.; Fontaine, H.; Hada, S.; Burrett, C.; Suphajanya, Th. and Kositanont, S. 1999c: Ancient analogs of burial alive extinction of the mastodons in catastroloess in Thailand, and of the last dinosaurs (in eggs) in Gobi Desert: further on tektites. Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000, October 28-29 1999, Bangkok, Thailand: 168-177.
- Bunopas, S.; Wasson, J. T.; Vella, P.; Fontaine, H.; Hada, S.; Burrett, C. and Kositanont, S., 1998. The early extraterrestrial collision with a whole comet in the Australasian tektite field, apparent evidences and discoveries from Thailand and East Asia (abstract). Geosea '98. Ninth Regional Congress on Geology, Mineral and Energy Resources of Southeast Asia. 17-19 August 1998, Shangri-La Hotel, Kuala Lumpur, Malaysia. Programme and Abstracts. 191-192.
- Bunopas, S.; Wasson, J. T.; Vella, P.; Fontaine, H.; Hada, S.; Burrett, C.; Suphajanya, Th. and Kositanont, S. 1999a: catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. *Journ. Geol. Soc. of Thailand* 1999 (1): 1-17.
- Bunopas, S.; Yaemniyom, Ng. and Kositanont, S. 1999b: Catastroloess and its derivatives, the life time sustainable construction sands was originally high atmospheric settling in the Quaternary cometary impact in Thailand and Asia. . Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year
- Bunopas, S.; Vella, P.; Fontaine, H.; Hada, S.; Burrett, C.; Haines, P.; Kositanont, S. and K. T. Howard 2001. Australasian Tektites and Catastrophic Products in Impact Ejecta Horizon from the Buntharik Events in Thailand. *Gondwana Res.*, v. 4 (4): 586.
- Bunopas, S.; P Vella; S. Hada; H. Fontaine; C. Burrett; P. Haines; S. Khositanont and K. T. Howard, 2002. Australasian Tektites and catastrophic products enclosed in impact ejecta horizon from the Buntharik Impact Event in Thailand on old surface gravel beds marked real-time Asian Quaternary correlation. *Ibid.* p. 181-187.
- Bunopas, S., Vella, P., H. Fontaine; S. Hada; C. Burrett; P. Hines S. Potisat; Ch.. Hinthiong; T. Wongwanich; P. Chaodamrong; K. T. Howard; A. Surinkum and S. Khositanont. 2002. New epeirogenesis, rapid vertical uplifting acme of Doi Inthanon in NW Thailand and late Himalayan extrusional continuum in the Lower to early Middle Pleistocene: An application of real-time layer. Program and Abstract of the Fourth Symposium

- of the IGCP Project No. 411, 17-25 November 2002. Phitsanulok, Thailand. P.110-119
- Bunopas, S., P. Vella; H. Fontaine, S. Hada, C. Burrett ; P. Haines ; S. Potisat, Th. Wongwanich' P. Chaodatrong, K. T. Howard, A. Surinkum and S. Khositantont, 2002. Shan-Thai and Indochina, Lower Paleozoic Gondwana derived Paired-Micro-continents growing Pangea by first continent-continent collision in late Norian with South China suffered second continent- continent collision of India to Asia. *Ibid* p..120-133
- Bunopas, S., P. Vella; H. Fontaine; S. Hada; C. Burrett; P. Haines; S. Potisat, Th. Wongwanich; P. Chaodamrong, K. T. Howard and S. Khositantont, 2003. The Khorat Plateau Early Quaternary Vertical Uplifting After Long Cenozoic Cratonization of the Mesozoic Khorat Group, Its northeast Tilting Diverted Mae Khong River Before the 0.8 Ma Cometary Impact Catastrophe and The Overflowing Mid Pleistocene Burirum Basalt. 1st International Conference on Paleontology of Southeast Asia. Mahasarakham University Journal, vol. 22. Special Issue 2003.
- Bunopas, S., P. Vella, H. Fontaine, S. Hada, C. Burrett, P. Haines, K. T. Howard and S. Khositantont., 2004.. Did Shan-Thai Twice Merge with Indochina and India? Discovery Quaternary Pushed-Pulled Himalayan Extensional Continuum Heritage to Major vertical Crustal Uplifting Inthanon Epeirogenesis, That Also Tilted Khorat Plateau Northeastwards Diverting Mae Khong River From Tonle Sap Prior to Buntharik Event, 0.8 MA Cometary Impact, Birirum Basalt And Asian Glacial Age. Internatinal Conf. on the Geologic Evolution of East and Southeast Asia. Department of Geology, Faculty of Science, Chulalongkorn University, 8-14 Febr. 2004, Asia Hotel, Bangkok. (Full paper in the print in 2005)
- Bunopas, S., P. Vella, H. Fontaine, S. Hada, C. Burrett, P. Haines, K. T. Howard and S. Khositantont., 2005. Latest Pliocene to early M. Pleistocene Inthanon Epeirogenesis, 2.0-0.65 Ma, in north Thailand connecting to late Himalayan extrusion continuum: An application of extraterrestrial / terrestrial real-time layer of catastrophe from Buntharik Cometary Impact Event (new). Proc. of the Int. Conference on Geology, Geotechnology and Mineral Resources of Indochina. GEOINDO 2005, pp. 468-499. Kosa Hotel, Khon Kaen, 28-30 Nov. 2005.
- Bunopas, S., P. Vella; H. Fontaine; S. Hada; C. Burrett; P. Haines; S. Khositantont; P. Chintasakul, P. Chaodamrong, P. Charusiri and K. T. Howard, 2007. Australasian Cometary Impact, 0.8ma3 Catastrophe Buried Alive Miocene1, Pliocene2 Faunas In Thailand And Central Australia, And Tektite-Bearing Flood Deposits In Ne. Thailand, New World's Discovery Extinction Age In Thailand (Indochina), and Australia Impact Fields4. (GEOETHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. Geothai 2007, pp. 30-43. 21-22 November. Imeral Hotel, Bangkok, Thailand.
- Burchfiel, B. C., P. Zhang, Y. Wang, W. Zhang, D. Jiao, F. Song, Q. Deng, P. Molnar, and L. H. Royden, Geology of the Haiyuan fault zone, Ningxia-Hui autonomous region, China, and its relation to the evolution of the northeastern margin of the Tibetan Plateau, *Tectonics*, 10, 1091-1110, 1991.
- Burek, P. J., 1984. 'Impacts, magnetic field reversals and tectonic episodes', Abstracts and Papers of the 15th Lunar and Planetary Science Conference, 102-103.
- Burns, C. A., 1989. Timing between a large impact and a geomagnetic reversal and the depth of NRM acquisition in deep-sea sediments. In: *Geomag. Palaeomag*, edited by F. J. Lowes, pp. 253-261.
- Burns, C. A. and B. P. Glass (1989), Source region for the Australasian tektite strewn field. *Meteoritics*. 24, 257.
- Burns C. A. 1990. The Australasian microtektite layer: Implications concerning its source and relationship to the Brunhes/Matuyama geomagnetic reversal. Ph.D. Thesis, Univ. of Delaware, Newark. 210 pp
- Burns. C A. and Glass, B. P. 1987. Microtektites and geomagnetic reversals revisited. *Meteoritics*. 22: 346-347 (Abstr.).
- Burrett, C. F., 1974, Plate tectonics and the fusion of Asia: *Earth and Planetary Science Letters*, v. 21, p.181-189.
- Burrett, C F., and Stait, B., 1985, South-east Asia as part of an Ordovician Gondwanaland- A paleogeographic test of a tectonic hypothesis: *Earth and Planetary Science Letters*, v. 75, p. 184-190.
- Burrett, CF., and Stait, B., 1986, China and southeast Asia as part of the Tethyan margin of Cambro-Ordovician Gondwanaland, in McKenzie, K., ed., *Shallow Tethys 2: Rotterdam, Balkema*, p. 65-77.
- Burrett, C., Long, J., and Stait, B. 1990. Early-Middle Palaeozoic biogeography of Asian terrains derived from Gondwana. In *Palaeozoic Palaeogeography and Biogeography*. Geological Society Memoir N12. (Eds W. S. McKerrow and C. R. Scotese.) 163 - 74.
- Burrett, C.; N. Duhig, R. Berty and R. Varne, 1991. Asian and South-western Pacific Continental Terranes Derived from Gondwana, and their Biogeographic Significance. *Australia. Syst. Bot.*, 4, 13-24.
- Burton, C. K. 1970. The palaeotectonic status of the Malay Peninsula. *Palaeogeography Palaeoclimatology, Palaeoecology* 7, 51-60
- Butler, B. E. 1956. Parna, an aeolian clay. *Aust. j. Sd.*, 18, 145-51.
- Butler, B. E. 1959. Periodic phenomena in landscapes as a basis for soil studies. *Soil Publ. C.S.I.R.O. Aust.* 14.
- Butler, B. E. 1964. The place of soils in studies of Quaternary chronology in southern Australia. *Revue Geomorph. dyn.* 14, 160-5.
- Butler, B. E. 1966. Soil periodicity in relation to land-form development in south eastern Australia, in Jennings, J. N. and Mabbutt, J. A. (Eds.), *Landform Studies of Australia and New Guinea*, Aust. natn. Univ., Canberra.
- Cambell, M. J., 1995. Permian – Triassic Links Between North-East Asia and New Zealand. *Journal of Geology, special issue, Series B, no. 5-6 (1995)*, pp. 304-305.
- Cameron, N. R., Clark., M. C. G., Aldiss, D. T., Aspden, J. A. and Djunuddin, A. 1980. Geologic evolution of northern Sumatra. proceedings of the Indonesian petroleum Association 9th Annual convention, Jakarta, 149\_97.
- Cande, S., D. V. Kent, 1995. Revised calibration of the geomagnetic polarity time scale for the Late Cretaceous and Cenozoic, *J. Geophys. Res.* 100(1995), 6093. 6095,
- Cardwell, R. K. and Isacks, B. L. (1978). Geometry of subducted lithosphere beneath the Banda Sea in eastern Indonesia from seismicity and fault plane solutions. *Journal of geophysical research* 83, 2825-38.
- Cardwell, R. K. and Isacks, B. L. (1981). A review of the

- configuration of the lithosphere subducted beneath the eastern Indonesian and Philippine islands. The geology and tectonics of eastern Indonesia. Geological Research and Development Centre Special Publication 2 (ed. A. J. Barber and S. Wiryosujono), pp. 3147. Bandung.
- Cardwell, R. K., Isacks, B. L., and Karig, D. E. (1980). The spatial distribution of earthquakes, focal mechanism solutions and subducted lithosphere in the Philippine and northeastern Indonesian islands' The tectonic and geologic evolution of Southeast Asian seas and islands. Geophysical monograph 23 (ed. D. E. Hayes), pp. 1-35. American Geophysical Union, Washington.
- Cassidy, W. A., Villar, L. M., Bunch, T. E., Kohman, T. P. and Milton, D. J., 1965. Meteorite and craters of the Campo del Cielo, Argentina. *Science* 149, 1055-1064.
- Cassidy W. A., Glass B. & Heezen, B. C. 1969. Physical and chemical properties of Australasia microtektites. *Journal of Geophysical Research* 74, 1008-1014.
- Cavosie, A.J., Nicholas E. Timms, N., Erickson, T., and Christian Koeberl, C., 2018. New clues from Earth's most elusive impact crater: Evidence of reidite in Australasian tektites from Thailand. *Geology* 46 (3): 203-206. <https://doi.org/10.1130/G39711.1>
- Chalmers, R. O., Handerson, E P. and Mason, B., 1976. Occurrence, distribution and age of Australasian tektites. *Smithson. Contrib. Earth Sci.*, 17: 46 pp.
- Chao, E. C. T., 1963. The petrographic and chemical characteristics of tektites. In: J.A. O'Keefe (Editor), *Tektites*. Univ. Chicago Press. Chicago. Ill.. pp. 51-94.
- Chao, E. C. T., Dwornik E. J. and Littler, J.. 1964. New data on the nickel-iron spherules from Southeast Asian tektites, and their implications. *Geochim. Cosmochim. Acta.* 28: 971-980.
- Chaodumrong, P, 1992, Stratigraphy, sedimentology and tectonic implication of the Lampang Group, central-north Thailand, Ph. D. thesis:- University of Tasmania, Australia, 203 p.
- Chapman, D. R. 1960. Recent re-entry research and the cosmic origin of tektites. *Nature* 188, 353.
- Chapman D. R. 1964. On the unity and origin of the Australasian tektites. *Geochim. Cosmochim. Acta* 28, 84 1-880.
- Chapman, D. R., 1971. Australasian tektite geographic pattern, crater and ray origin, and theory of tektite events. *Jour. Geophys. Resear.* 76, 6309-6338.
- Chapman, D. R. and Larson, H. K. 1963. On the Lunar origin of tektites. *J. Geophys. Res.* 68: 4305 - 4358.
- Chapman, D. R. and Scheiber, L. C., 1969. Chemical investigation of Australasian tektites. *J. Geophys. Res.* 74, 6737-6776.
- Chapman, D. R. and Scheiber L. C. 1989. Tables of tektite chemical analyses conducted in 1966/69. Unpublished data filed at the Smithsonian Museum, Washington. D. C. with the NASA/Ames collection of tektites.
- Chapman, D.R., Zimmermann, B., Centolanzi F. S. and Scheiber, L.C. 1967. Effects of etching and of earth landing on aerodynamically ablated tektites (abstract). *Ann. Meet Meteorit. Soc.* 30th, NASA Ames Research Center, Moffett Field, California.
- Chaimanee, Y., 1998. Plio-Pleistocene rodents of Thailand. *Thai Studies in Biodiversity*, 3, 1-3 13.
- Chaimanee, Y., 2004. *Siamopithecus eocaenus*, anthropoid primate from the Late Eocene of Krabi, Thailand. In *Anthropoid origins New Visions*, C. F. Ross and R. F. Kay, eds. (New York; Kluwer Academic/ Plenum Publishers), pp. 341-368.
- Chaimanee, Y., 2006. Late Pleistocene vertebrate records from Southeast Asia. In *Encyclopedia of Quaternary*, pp. 3189-3197. Elsevier.
- Chaimanee, Y. & Jaeger, i-i., 2000. Occurrence of *Hadromys humei* (Rodentia: Muridae) during the Pleistocene in Thailand. *J. Mammalogy* 81(3), 659-665.
- Chaimanee, Y., Suteethorn, V., Jaeger, J.-J. & Ducrocq, S., 1997. A new Late Eocene anthropoid primate from Thailand. *Nature* 385, 429-431.
- Chaimanee, Y., Khansubha, S. & Jaeger, J.-J. 2000. A new lower jaw of *Siamopithecus eocaenus* from the Late Eocene of Thailand. *C. R. Acad. Sci.* 323, 235-241.
- Chaimanee, Y., Jolly, D., Benammi, M., Tafforeau, P., Duzer, D., Moussa, I. & Jaeger, J.-J., 2003. A new middle Miocene hominoid from Thailand and orangutan origins. *Nature* 422, 61-65.
- Chaimanee, Y., Suteethorn, V., Jintasakul, P., Vidthayanon, C., Marandat, B. & Jaeger, J.-J., 2004. A new orang-utan relative from the Late Miocene of Thailand. *Nature* 427, 439-44 1.
- Chaimanee, Yaovaluk, 2004. Fossils from Khok Soong and Ban Tha Chang, Khorat. In: *Numerous fossils in Paleontology of Thailand, in 114 years of the Department of Mineral Resources.* (in Thai)
- Chaimanee, Yaovaluk, 2007. Diversity of Cenozoic Mammals in Thailand: Paleoenvironment and Age Updated. (GEOTHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. *Geothai 2007*, pp. 73-79. 21-22 November. Imeral Hotel, Bangkok, Thailand.
- Chaimanee, Yaovaluk, Chotima Yami, Panipa Saethien, Kitti Kaovises 2002. Diversity of Ancient Elephants in Nakorn Ratchasima. Director Office of Geology.
- Chakraborty, K. R. 1980. On the evolution of the nepheline to hypersthene normative alkali basaltic rocks of Kuantan, Pahang, Peninsular Malaysia. *Geological Society of Malaysia Bulletin*, 79-85.
- Chakraborty, K. R. and Kamineni, D. C. 1978. Chemical characteristics and formation of Segamat volcanics. *Geological Society of Malaysia Newsletter* 4, No. 3, SS-9i.
- Chalmers, R. O., Henderson, E, P., Mason, B. (1976): occurrence, distribution, and age of Australian tektites. - *Smithsonian Contrib. to the Earth - Sciences*, 17; 46.
- Chamalaun, F. H., Grady, A. E., von der Borch, C' C', and Hartono, H. M. S. (1981). The tectonic significance of Sumba. *Geological Research and Development Centre Bulletin* 5 , 7-20.
- Charusiri, P., Daorerk, V., Krowchan, V., Klongsara, N., Kosuwan, S., Srirattanachatchawan, V. and Santatiwongchai, U., 2002. Quaternary tektites and their sediment hosts at Ban Tachang sand pit, Chaloe Prakit, Nakhon Ratchasima, NE Thailand: stratigraphy and TL ages. *The Symposium on Geology of Thailand, 26-31 August 2002, Bangkok, Thailand.*
- Chaussidon, M., and C. Koeberl (1995), Composition of tektites and impact glasses: Constraints on source regions, *Geochim. Cosmochim. Acta*, 59, 613-624.
- Chen, J. C. (1975). Geochemistry of andesites from the Coastal Range., eastern Taiwan. *Geological Society of China Proceedings* 18, 73-88'
- Chen D.N., Lu, Y.C. & An, Z.S., 1982. Snail assemblages in loess strata and their environmental implication.

- in: Proc. 3rd Nat. Quat. Conf. China, 7-15, Science Press, Beijing.
- "The K/Ar age determination of Quaternary geological samples, the Ar38 dilution method." 1981. (Annual Reports - Institute Of Geochemistry, Academia Sinica: Ching-Kuo K'o Hsueeh Yuan. Ti Ch'iu Hua Hsueeh Yen Chiu So: Vol. 1978-1 979. p. 40-41)
- Chhibber, H. L., 1934. The Geology of Burma. MacMillan & Co., London.
- Chualaowanich, T., Saisuthichai, D., Sarapanchotewittaya, P., Charusiri, P., Sutthirat, C., Lo, C.-H., Lee, T.-Y., and Yeh, M.-W., 2008, New 40Ar/39Ar Ages of some Cenozoic basalts from the east and northeast of Thailand : In Proceedings, International Symposia on Geoscience Resources and Environments of Asian Terranes (GREAT 2008), Bangkok, Thailand, p. 225-229
- Chunlai, L., Ziyuang, O.. and Dongsheng. L.. 1993. Microtektites and glassy microspherules in loess: Their discoveries and implications: Science in China: Ser. B, v. 36, no. 9, p. 1141-1152.
- Claude, J., Naksri, W., Boonchai, N., Buffetaut, E., Duangkrayom, J., Laojumpon, C., Jintasakul, P., Lauprasert, K., Martin, J., Suteethorn, V., and Tong, H. 2011. Neogene reptiles of northeastern Thailand and their paleogeographical significance. *Annales de Paléontologie*, 97:113–131. <https://doi.org/10.1016/j.annpal.2011.08.002>
- Clement, B. M. and D.V. Kent, 1991. A southern hemisphere record of the Matuyama-Brunhes polarity reversal. *Geophys. Res. Lett.* 18, 18-84, 1991.
- Clement, B. M., Kent, D. V. and Opdyke, N. D., 1982. 'Brunhes-Matuyama polarity transition in three deep-sea sediment cores', *Phil. Trans. Roy. Soc.*, A306, 113-119.
- Cleverly W. H. 1986. Australites from Hampton Hill Station, W.A. *Journal of the Royal Society of Western Australia*, 68, 81-93,
- Coaldrake, J. E., 1961. The ecosystem of coastal lowland ("Wallum") of Sothern Quensland. *Bull. Commonw. Scient. Ind. Res. Org.* 263.
- Cobbold, P., and P. Davy, Indentation tectonics in nature and experiment, 2. Central Asia, *Bull. Inst. Uppsala*, 14, 143-162, 1988.
- Cohen, A. J., 1963. Asteroid- or comet-impact hypothesis of tektite origin: the moldavite-strewn fields: In: *Tektites*, J. A. O'Keefe, ed., Univ. Chicago Press, Chicago, 189-211.
- Condie, K.C. (2001) Supercontinents, superplumes and continental growth: the record in the Neoproterozoic supercontinent Rodinia. *Geol. Soc. Lond., Spl. Publ.* (in press).
- Condie, K.C., Des Marais, D.J. and Abbott, D (2001) Precambrian superplumes and supercontinents: a record in black shales, carbon isotope, and paleoclimates. *Precamb. Res.* (in press).
- Cope, M. J., and Chaloner, W.G. 1985. Wildfire, an interaction of biological and physical processes, in Tiffriey, B. H., ed., *Geological Factors and the Evolution of Plants*: New Haven, Yale University Press, p. 257 -277
- Corbett K.D., and Brown A.V. 1975. *Geological Atlas 1:250 000 Series*, Queenstown. Geological Survey of Tasmania, Department of Mines
- Corbett K.D., Pemberton J., and Vicary M.J. 1993. Map 13. *Geology of the Mt Jukes-Mt Darwin Area, 1:25 000*. Geological Survey of Tasmania, Department of Mines.
- Cotton, C. A., 1963. The question of high Pleistocene shorelines. *Trans. Roy. Soc. N. Z. (Geol.)* 2, 51-62.
- Cox, A., Dalrymple, G. B. and Doell, R. R., 1967. Reversals of the Earth's magnetic field. *Sci. Am.*, 44-54.
- Clarke, W.B., Beg, MA., and Craig, H., 1969. Excess 3He in the sea: evidence for terrestrial primordial helium. *Earth Planet. Sci. Lett.* 6, 213-220.
- Clement, B. M., DV. Kent and N. D. Opdyke. 1982. Brunhes-Matuyama polarity transition in three deep-sea sediment cores, *Philos. Trans. R. Soc. A306*, 113-119, 1982.
- Craig, H., 1991. 3He in MORB, BABB, and oceanic and continental hotspots: an overview and the case for primitive mantle. *Cal.Tech. Plume Symposium*, May, 1991.
- Craig, H., and Rison, W., 1982. Helium 3: Indian Ocean Hot Spots and the East African Rift (abstract). *EQS Trans.* 69. (Spring Meeting Suppl.), 1144.
- Cranbrook, F. O. 1981. The vertebrate faunas. In 'Wallace's Line and Plate Tectonics Tectonics. (Ed. T. C. Whitmore.) 57-69. (Clarendon Press: Oxford.)
- Crawford, A. R. 1974a. The Indus suture line, the Himalaya and Gondwanaland. *Geological Magazine* 111, 369-83.
- Crawford, A. R. 1974b. A greater Gondwanaland. *Science* 184, 1179 -81.
- Crawford' A. R."Tektites plobablv wholly terrestrial and related to continental movement.,' July 1979. (*GEOL. MAG.* ; Vot. I 16, No. 4, p. 261-276)
- Credner, W. 1935. *Siam, Das Land der Tai*, J. Engelhorn's Nachr., Stuttgart, Germany.
- Croft, S.K. 1982. *Proc. Lunar P1anet Sci. Conf.* 15th, C828-C842
- Crostella, A. 1977. Geosynclines and plate tectonics in Banda arcs, eastern Indonesia. *American Association of Petroleum Geologists Bulletin* 57, 2063-81.
- Crostella, A. 1931. Malacca Strait wrench fault controlled Lalang and Mengkapan oil fields' Offshore S.E. Asia oilfield, proceedings 2, S.E.A.P.E.X., Singapore, t-12.
- Geophysics of the Banda Arc and Adjacent Areas, Part I, Snellius II Symposium Proceedings: Netherlands Sea Research Journal, v. 24,p. 13 1-140.
- Curray, J. R., 1989, The Sunda arc: a model for oblique plate convergence, in *Geology and*
- Curray, I. R., Emmel, F. J., Moore, D. G., and Raitt, R. W. 1982. Structure, tectonics and geological history of the northeastern Indian Ocean. The ocean basins and margins: the Indian ocean 6 (ed. A. E. M. Nairn and F. G. Stehli), pp. 399-450. Plenum Press, New York.
- Curray, J. R., Moore, D. G., Lawver, L. A., Emmel, F. J., Raitt, R. S7., Henry, M., and Kiekhefer, R. 1979. Tectonics of the Andaman Sea and Burma. *Geologi-cal and geophysical investigations of continental slopes and rises*, American Association of petroleum Geologists Memoirs 29 (ed. J. S. Watkins, L. Montadert, and P. W. Dickerson); pp. l89-98.
- Curray, J. R. and Moore, D. G. 1974. Sedimentary and tectonic processes in the Bengal deep-sea fan and geosyncline. *The geology of continental margins* (ed. C. A. Burke and C. L. Drake), pp. 6 17-237. Springer-Verlag, Berlin.
- Curry, J. R., Emmel, F. J., Mooie, D. G., and Raitt, R. W. 1982. Structure, tectonics and geological history of the northeastern Indian Ocean. The Indian Ocean basins and margins: the Indian ocean 6 (ed. A. E. M.

- Nairn and F. G. Stehli), pp.399-450. Plenum Press, New York.
- Cuttitta, F., Clarke. R. S., Jr., Carron M. K. and Annell, C.S., 1967. Marthas Vineyard and selected Georgia tektites: new chemical data. *J. Geophys. Res.* 72: 1343-1349.
- Cuttitta, F., Carron, M. K. and Annell. C. S., 1972. New data on selected Ivory Coast tektites. *Geochim. Cosmochim. Acta.*
- Daines, S. R. 1985. Structural history of the W Natuna Basin and the tectonic evolution of the Sunda region. Proceedings of the Indonesian Petroleum Association 14th annual convention, Jakarta, October 1985 1, 39-61.
- Darbyshire, D. P. S. 1975. K/Ar determination of basaltic samples from Vietnam. Unpublished report of. the isotope geology unit, Institute Geological Sciences, London, 75 /2.
- Darwin, C. 1844. Geological observations on the volcanic islands and parts of South America visited during the voyage of HMS Beagle. Smith, Elder & Co.
- Darwin, C. 1859. On the origin of species by mean of natural selection or the preservation of favoured races in the struggle for life. John Murray, London, 513 p.
- David T. W. E., Summers H. S. & Ampt G. A. 1927. The Tasmanian tektite - Darwin glass. Proceedings of the Royal Society of Victoria 39, 167- 190.
- Davies, H. L. and Jaques, A. L. 1994. Emplacement of ophiolite on Papua New Guinea. Ophiolites and oceanic lithosphere (ed. I. G. Gass, S. J. Lippard, and A. W. Shelton), pp. 341-9. Geological Society, London.
- Davies, P. R. 1984. Tertiary structural evolution and related hydrocarbon occurrences, North Sumatra Basin. Proceedings Indonesian Petroleum Association 13th Annual Convention, Jakarta, May 1984 1, 19-49.
- De Beaumont, G. 1973. Guide des vertebres fossils, Detachaux et Neistle, 476pp.
- De Boer, J., Odom, L. A., Ragland, P. C., Snider, F. G., and Tilford, N. R. 1980) The Baiaan origin., eastwards subduction, tectonic rotations and volcanism in the western Pacific (Philippines). *Tectonophysics* 67, 25 I-BZ.
- De Coster, G. L. 1975. The geology of the central and south Sumatran basins. Proceedings of the Indonesian Petroleum Association Third Annual Convention, Jakarta, June 1974, 77-110.
- De Gasparis, A. A., Fuller, M. and Cassidy, W. A., 1975. Natural remanent magnetism of tektites of Muong-Nong type and its bearing on models of their origin. *Geology* 3, 605-607.
- De Master, D. J. and J.K. Cochran. 1982. Particle mixing rates in deep-sea sediments determined from excess <sup>210</sup>Pb and <sup>32</sup>Si profiles, *Earth Planet. Sci. Lett.* 61, 257-271..
- De Menocal, P. B., Ruddiman, W. F. and D. V. Kent, 1990. Depth of post-depositional remanence acquisition in deep sea sediments: a case study of the Brunhes -Matuyama reversal and oxygen isotopic Stage 19.1. *Earth Planet. Sci. Lett.*, 99, 1-13.
- Denny, C. S., compiler, 1970. Glacial Geology of Coterminous United States, in The National Atlas of the United States of America: U.S. Department of Interior, Geological Survey, p. 76.
- Dehn J., Farrell, J., and Schmincke, H. U., 1991. Neogene tephro chronology from site 758 on Northern Ninetyeast Ridge: Indonesian arc volcanism of the past 5 Ma. *Proc. ODP, Sci, Results.* 121, 273-295.
- Des Marais, D.J., Strauss, H., Summons, R'E and Hayes' J M' 1992. Carbon isotope evidence for the stepwise oxidation of the Proterozoic environment. *Nature*, v. 359, pp' 605-609'
- Desprairies, A., Riviere, M., and Pubellier, M. 1991. Diagenetic evolution of Neogene volcanic ashes (Celebes and Sulu Seas) - *Proc. ODP, Sci. Results* 121, 489-503.
- Ding, Z. L. & Liu, T.S., 1989. Progresses of loess research in China, 1. Loess stratigraphy. *Quat. Sci.* 1, 24-35. (in Chinese with English abstract)
- Ding, Z.L. & Liu, T.S., Liu, X.M., Chen, M.Y. & An, Z.S., 1989. 37 climatic cycles during the past 2.5 Ma. *Clim. Sd. Bull.* 34, 1494-1496. (in Chinese)
- Ding, Z.L., Rutter, N. & Liu, T.S., 1992. Pedostratigraphy of Chinese loess deposits and climatic cycles in the last 2.5 Myr. *Catena* 20, 73-91.
- Divis, A. F. 1980. The petrology and tectonics of Recent volcanism in the central Philippine islands. The tectonic and geologic evolution of Southeast Asian seas and islands, Geophysical monograph, 23(ed. D. E. Hayes), pp.12744. American Geophysical Union, Washington.
- Dou Chengxun. et al. "Preliminary report on tektite in Dianbai, Guangdong." Oct. 1989. (*Dizhi Lun'ing* = Geological Review; Vol. 35, No. 5, p. p. 414-422).
- Duangkrayom J, Ratanasthien B, Jintasakul P, Carling P.A. 2014. Sedimentary facies and paleoenvironment of a Pleistocene fossil site in Nakhon Ratchasima province, northeastern Thailand. *Quaternary International* 325: 220–238. doi: 10.1016/j.quaint.2013.07.048
- Duangkrayom, J., Wang, S.Q., Deng, T., Jintasakul, P., 2017. The first Neogene record of Zygodontodon (Mammalia, Proboscidea) in Thailand: implications for the mammutid evolution and dispersal in Southeast Asia. *Journal of Paleontology* 91 (1), 179–193.
- Duarah, R., Saikia, M. M., and Bhattacharjee, C. C. 1983. Occurrence of the ophiolitic complexes along the Indoburman orogenic belt. 120, 175-82.
- Ducrocq, S. 1992. Etude biochronologique de bassins continentaux tertiaires du Sud-est Asiatique: Contribution des faunes de mammifères, Université Montpellier II.
- Ducrocq, S. Chaimanee, Y., Suteethorn, V., and Jaeger, J.-J.. 1994. Ages and paleoenvironment of Miocene mammalian faunas from Thailand. *Paleogeography Palaeoclimatology Paleocology.* 108: 149-163.
- Durrani, J. A. and H. A. Khan, 1971. Ivory Coast microtektites Fission-track ages and geomagnetic reversals, *Nature* 232. 320-323, 1971.
- Duval, Mathieu, Fang, Suraprasit, K., Jaeger, J, Benammi, M., Chaimanee, Y., Iglesias C., Javier, and Grün, Rainer. 2019. Direct ESR dating of the Pleistocene vertebrate assemblage from Khok Sung locality, Nakhon Ratchasima Province, Northeast Thailand. *Palaeontologia Electronica* 22.3.69 1-25. [https://doi.org/10.26879/941\\_palaeo-electronica.org/content/2019/2794-esr-dating-of-khok-sung-site](https://doi.org/10.26879/941_palaeo-electronica.org/content/2019/2794-esr-dating-of-khok-sung-site)
- Emilia, D. A., D. F. Heinrich. 1969. Ocean floor spreading: Oldavai and Gilsa Events in the Matuyama Epoch. *Science*, v 154, p. 149.
- Emiliani, C., Kraus, E. B., and Shoemaker, E. M., 1981. Sudden death at the end of the Mesozoic. *Earth Planet. Sci. Lett.*, 55, 317.

- England, P.C., and Houseman, G., 1989. Extension during continental convergence, with application to the Tibetan Plateau. *J. Geophys. Res.* 94, 17561-17579.
- England, P., and P. Molnar, 1990. Right-lateral shear and rotation as the explanation for strike-slip faulting in eastern Tibet, *Nature*, 344, 140-142, 1990.
- Einsale, G. and Seilacher, A. (Eds). 1982. *Cyclic and Event Stratigraphy*. Springer Verlag, Berlin.
- Falvey, D. A., and Greene, H. G. 1988. Origin and evolution of the sedimentary basins of the Ness Hebrides arc. in *Geology and Offshore Resources of Pacific Island Arcs Vanuatu Region*, Circum-Pacific Council for Energy and Mineral Resources Earth Science Series. Volume 8'. (Eds H. G. Greene and F. L. Wong.) 413-42. (Circum-Pacific Council for Energy and Mineral Resources: Houston, Texas.)
- Falvey, D. A., and Mutter, J. C. 1981. Regional plate tectonics and the evolution of Australia's passive continental margins. *BMR. Journal of Australian Geology and Geophysics* 6, 1-29.
- Faul, H., 1966. Tektites and terrestrial. *Science* 152, 1341-1345.
- Fiske, P. S., Lowe, T., 1995. Large Compositional Variations in Layered Tektites of NE Thailand: Implications for the Dynamics of Large Impact-Induced Explosions on Earth. *Meteoritics* 30, 504.
- Fiske, P. S.; Putthapiban, P. and Wasson, J. T. 1996. Excavation and analysis of layered tektites from Northeast Thailand: Results of 1994 field expedition. *Meteoritics and Planet. Sci.*, 31, 36-41.
- Fiske, P.S., Schnetzler, C.C., McHone, J., Chanthavaichith, K.K., Homsombath, I., Phouthakayalat, T., Khenthavong, B., Xuan, P.T., 1999. Layered tektites of southeast Asia: Field studies in central Laos and Vietnam. *Meteoritics & Planetary Science* 34, 757-761.
- Fleischer, R. L. et al. Fission track evidence for the simultaneous origin of tektites and other natural glasses. (*Geochim. Et Cosmochim. Acta*. v. 28, no. 6, p.755-760. 1964)
- Fleischer, R. L., P. B. Price and R. M. Walker, 1965. On the simultaneous origin of tektites and other natural glasses. *Geochim. Cosmochim. Acta* 29, 161-166. 1965.
- Fleming, C. A., 1953. The Geology of Wanganui Subdivision. *Bull. Geol. Survey, N. Z.* 62.
- Fleming, C. A. 1962. New Zealand biogeography: a paleontologist's approach. *Tuatara* 10, 53-108.
- Flint, R. F. 1971. *Glacial and Quaternary Geology*, Willy, New York.
- Florensky, P.V and Dabizha, A.I., 1980. Zhamanshin meteorite Crater (Meteoritny Krater Zhamanshin). *Nauka, Moscow*, 127 pp.
- Flower, M., 1991. Cenozoic magmatism in Indochina and the South China Sea. Second Conference on Geology of Indochina, 11-13 November 1991, Hanoi.
- Flower, M.F.J., Hoang, N., Yem, N.T., Bao, N.X., McCabe, R.J., and Harder, S.H., 1993. Cenozoic magmatism in Indochina: lithosphere extension and mantle potential temperature. *Geol. Soc. Malaysia Bull.* 33, 211-222.
- Foden, J. D. 1985. The petrology of Tamboro volcano, Indonesia: a model for the 1815 eruption. *Journal of Volcanology and Geothermal Research* 27, 141.
- Foden, J. D. and Varne, R. (1981). Petrogenetic and tectonic implications of near coeval calc-alkaline to highly alkaline volcanism on Lombok and Sumbawa islands in the eastern Sunda arc. The geology and tectonics of eastern Indonesia (ed. A. J. Barber and S. Wiryosujono), pp. 135-52. Geological Research and Development Centre, Bandung.
- Folco, L.M. D'Orazio, M. Tiepolo, S. Tonarini, L. Ottolini, N. Perchiazzi, P. Rochette, B.P. Glass 2009 Transantarctic Mountain microtektites. Geochemical affinity with Australasian microtektites *Geochim. Cosmochim. Acta*, 73, 3694-3720.
- Folco, L., B.P. Glass, M. D'Orazio, P. Rochette 2010 A common volatilization trend in Transantarctic Mountain and Australasian microtektites: implications for their formation model and parent crater location *Earth Planet. Sci. Lett.*, 293 135-139.
- Folco, L., Perchiazzi, N., D'Orazio, M., Frezzotti, M.L., Glass, B.P., Rochette, P., 2010. Shocked quartz and other mineral inclusions in Australasian microtektites. *Geology* 38, 211-214
- Folco, L., Rochette, P., Perchiazzi, N., D'Orazio, M., Laurenzi, M. A., Tiepolo, M. 2008. Microtektites from Victoria Land Transantarctic Mountains. *Geology*, 36 (4), 291-294, doi: 10.1130/G24528A.1.
- Folco, L., Rochette, P., Suavel, C., and Gattacceca, J. 2006. Micrometeorites from Frontier Mountain (Antarctica): Abstracts, *Meteoritics and Planetary Science*, v. 41, P. A56
- Folk, R.L. and Ward, W.C. (1957) A Study in the Significance of Grain-Size Parameters. *Journal of Sedimentary Petrology*, 27, 3-26. <https://doi.org/10.1306/74D70646-2B21-11D7-8648000102C1865D>
- Fontaine, H. 1973. *Bibliographie des sciences de la terre pour le Vietnam, le Laos et le Cambodge*. Archives géologiques du Vietnam 16, Saigon.
- Fontaine, H. 1979. Note on the geology of the Calamian islands, north Palawan, Philippines. *C.C.O.P. Newsletter* 6, No.2, 4047.
- Fontaine, H. 1985. Discovery of Lower Permian corals in Sumatra. *Geological Society of Malaysia Bulletin* 19, 183-91.
- Fontaine, H. and Workman, D. R. 1978. Review of the geology and mineral resources of Cambodia, Laos and Vietnam. *Geology and mineral resources of Southeast Asia* (ed. Prinya Nutalaya), pp. 538-603. Asian Instit. Technology, Bangkok.
- Fontaine, H. and Hoang Thi Than, 1971. Alluvions anciennes du Nam Phan septentrional. *Arch. Geol. Du Vietnam*. N 14, p. 145-168.
- Fontaine, H., 1966. Decouverte de tectites sans formes Fig.es dans la province de Bien-Hoa. *Archives Des Geologie du Viet-Nam* 9, 3-4.
- Fontaine, H. & Workman, D. R., 1978. Review of the geology and mineral resources of the Cambodia, Laos, and Vietnam. In P Nutalaya P. ed. *Proceeding of the Third Regional Conference and Mineral Resources on Southeast Asia*, pp. 541-572. Asian Institute of Technology, Bangkok
- Ford, R., J. 1988: An empirical model for the Australasian tektite field. *Australian J. Earth Sci.* 35: 483-490.
- Fraser, G. S., and Bleuer, N. K., 1988, Sedimentologic consequences of two floods of extreme magnitude in the late wisconsinan, in Clifton, H. E., ed., *Sedimentologic Consequences of Conclusive Geologic Event: Geological Society of America Bulletin*, p. 111-132.
- French, B.M., 1998. *Traces of Catastrophe: A Handbook of Shock-Metamorphic Effects in Terrestrial Meteorite*

- Impact Structures. LPI Contribution No. 954, Lunar and Planetary Institute, Houston.
- Frey, F. A., C.M. Spooner and P.A. Baedeker, 1970. Micro-tektites and tektites: a chemical comparison. *Science* 170, 845-847. 1970.
- Frye, J. C., H. B. William, M. Robin and R. F. Black. 1968. Definition of Wisconsin Stage: U. S. Geological Survey Bull. 1274-E, Washington. U.S. Government-printing.
- Fudali, R. F. and Ford, R. J., 1979. Darwin glass and Darwin crater; a progress report. *Meteoritics*, 14, 283-296
- Fudali, R.F., Dyar, M.D., Griscom, D.L. and Schreiber, H.D., 1987. The oxidation state of iron in tektite glass. *Geochimica et Cosmochimica Acta* 51, 2749-2756.
- Fudali, R.F., Fredriksson, K., 1992. Tektite-like bodies at Lonar Crater, India? Very unlikely. *Meteoritics* 27, 99-100.
- Fudali R. F., 1993. The stratigraphic age of australites revisited. *Meteoritics* 28, 114-119.
- Geological map of Vietnam, Cambodia, Laos. Scale 1:2,000,000. National Geographic Directorate of Vietnam, Dalat.
- Fuller, M., and Lin Jin-Lu 1984. Paleomagnetism of an accretionary margin: S. China and S.E. Asia. *Geological Society of Malaysia: GEOSEA V, Abstracts of Papers*, 10-11.
- Fuller, M., et al., 1983. Paleomagnetism of Luzon. Hayes, D. E. (ed.) *The tectonic and geologic evolution of Southeast Asian seas and islands: part 2, Geophysical monographs* 27 (ed. D. E. Hayes), pp. 79-94. American Geophysical Union, Washington.
- Futrell, D. S. and Fredriksson, K., 1983. Brecciated Muong-Nong type tektites. *Meteoritics* 18, 15-17.
- Futrell, D. S. 1986. Implication of welded breccia in Muong Nong-type tektite. *Nature*, London. 319, 663-665.
- Futrell, D. S. "Accumulations of inclusions inside a muong nong-type tektite." *Oct. 27, 1992. (Eos, Transactions, American Geophysical Union: Vol.73, No.43, Suppl., p. 328)*
- Futrell, D. S. and Wasson, J. T., 1993. A 10.8-kg layer (Muong-Nong-type) tektite from Wenchang, Hainan, China. *Meteoritics* 28, no. 1, 136-137.
- Gansser, A. 1964. *Geology of the Himalayas*. John Wiley-Interscience, New York.
- Gansser, A. 1966. The Indian Ocean and the Himalayas, a geological interpretation. *Eclogae Geologicae Helvetiae* 59, 831-48.
- Gansser, A. 1980. The significance of the Himalayan suture zone. *Tectonophysics* 62, 37-52. Garson, M. S., Young, B., Mitchell, A. H. G., and Tait, B. A. R. 1975. The geology of the tinbelt in Peninsular Thailand around Phuket, Phangnga and Takua Pa. *Overseas memoir* 1.
- Garson, M. S., Amos, B. J., and Mitchell, A. H. G. (1975). The geology of the area around Nyaungga and Ye-ngan, southern Shan States, Burma. *Overseas memoir* 2.
- Gage, M. 1958. Late Pleistocene glaciations of the Waimakariri Valley, Canterbury, New Zealand. *N.Z. J. Geol. Geophys.* 1, 123-55.
- Gage, M. 1961. New Zealand glaciations and the duration of the Pleistocene. *J. Glaciol.* 3, 940-3.
- Gage, M. 1965. Some characteristics of Pleistocene cold climates in New Zealand. *Trans.R. Soc. N.Z. (Geol.)* 3 (2), 11-21.
- Gage, M., and Suggate, R. P. 1958. Glacial chronology of the New Zealand Pleistocene. *Bull. geol. Soc. Am.* 69, 589-98.
- Galloway, R. W., 1965. Quaternary climate of the Australian region. *J. Geol.* 3. 603-18
- Garlong, V. and Kuttikul, P., 1999. The application of ground penetrating radar (GPR) for petrified wood mapping. *Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000, October 28-29 1999, Bangkok, Thailand: 135-141.*
- Ganapathy, R., 1982. Evidence for a major meteorite impact on Earth 34 million years ago: Implication for Eocene extinction. *Science* 216, 885-886.
- Ganapathy, R. and Larimer, J. W., 1983. Nickel-iron spherules in Tektites and other glass. *Earth Planet. Sci. Lett.*, 65., 225-228.
- Gangadharam, E. W., et al., 1978. Tektites from Burma? A suspected new tektite locality in Southeast Asia. *Wata Geol. Newsletters of Geol. Soc. of Malaya*, vol 4 (1), 13-17.
- Garlik, G. D., Naeser, C. W. and O'neil, J. R., 1971. A Cuban tektites. *Geochim. Cosmochim. Acta* 35. 731-734.
- Gatinsky, Y. G. and Hutchison, C. S. 1984. Cathaysia, Gondwanaland and the paleotethys in the evolution of continental Southeast Asia. *Geological Society of Malaysia: GEOSEA V, Abstracts of Papers*, Kuala Lumpur, 11-12.
- Gatinsky, Y. G., Mischina, A. V., Vinogradov, I. V., and Kovalev, A. A. 1978. The main metallogenic belts of Southeast Asia as a result of different geodynamics conditions interference. *Geology and mineral Resources of Southeast Asia (ed. Prinya Nutalaya)*, pp. 313-18. Asian Institute of Technology, Bangkok.
- Gatinsky, Y. G., Hutchison, C. S., Nguyen, Minh, N.N., and Tri, T. V. 1984. Tectonic evolution of Southeast Asia. *27th International Geology Congress Reports 5, Colloquium 5, Tectonics of Asia*, 225-40.
- Gaudemer, Y., P. Tapponnier, B. Meyer, G. Peltzer, Guo Shunmin, Chen Zhitai, Dai Huagang, and I. Cifuentes, 1995. Partitioning of crustal slip between linked, active faults in the eastern Qilian Shan, and evidence for a major seismic gap, the "Tianzhu gap", on the western Haiyuan Fault, Gansu (China), *Geophys. J. Res.*, 120, 599-645, 1995.
- Gehrels, G. E., and Saleehy, J. B. 1987. Geologic framework, tectonic evolution, and displacement history of the Alexander terrane. *Tectonics* 6, 1-74
- Gentner, W., 1966. Auf der suche nach Kraterglassern, Tektiten und Meteoriten in Africa, *Naturwissenschaften* 12, 285-289, 1966.
- Gentner, W., Kleinman, B. and Wagner, G. A., 1967. New K-Ar and fission track of impact glasses and tektites. *Earth Planet. Sci. Lett.* 2, 82-86.
- Gentner, W.; H. J. Lippolt and O. Muller, 1964. K-Ar-Alter des Bosumtwi-Kraters in Ghana mid die chemische Beschaffenheit seiner Gläser, *Z. Naturforsch.* 19a. 150-153, 1964.
- Gentner, W., Storzer, D. and Wagner, G. A., 1969. New fission-track ages of tektites and related glasses. *Geochimica et Cosmochimica Acta* 33, 1075-1081.
- Gentner, W., Glass, B. P., Storzer, D. and Wagner, G. A., 1970. Fission ages and ages deposition of deep sea microtektites. *Science* 168, 359-361.
- Geological Bureau of Xizang, 1982. *Tectonics of Yarlung-Zangbo suture zone Xizang (Tibet)-guide to geological excursion*. Geological Bureau of Xizang Geological Society of Xizang, China.

- Gerling, E. K. and Ni. L. Yaschenko, The age of tektites, Dokl. Akad.Nauk SSSR, 83, 901-902, 1952
- Geronimo, I. Di; R. Sanfilippo; N. Chaimanee. 2002. The Quaternary Fauna from Ban Pak Bo (Western Thailand). The Symp. on Geology of Thailand. 26-31 August 2002, Bangkok, Thailand. Additional.
- Gherbrant, E., Sudre, J., and Capetta, H. 1996. Paleocene proboscidean from Morocco. *Nature*, 383: 68-70.
- Giggenbach, W.F. and Poreda, A.J., 1993. Helium isotopic and chemical composition of gasses from volcanic - hydrothermal systems in the Philippines. *Geothermics* 22, 369-380.
- Gilchrist, J.; Thorpe, A. N. and Senftle, F.E., 1969. Infrared analysis of water in tektites and other glasses. *J. Geophys. Res.*, 74, 1475-1483.
- Ginsburg, L. and Tassy, P., 1985. The fossil mammals and the age of the Lignite beds in the intramontane basins of northern Thailand. *Jour Geo. Soc Thailand*. 8(1-2): 13-27.
- Gladenkov, Y. B. 2003. New Stratigraphic Schemes for the Cenozoic of the Russian far east and the time scales of Geologic Events. *Proceedings, 8th International Congress on Pacific Neogene Stratigraphy, Chang Mai, Thailand, 2003*, p.10-19
- Gradstein, F. and Ludden, J. 1988. Leg 123. Site 765. Ocean Drilling Program
- Glass, B.H. . 1968. Glassy objects (microtektites?) from deep-sea sediments near the Ivory Coast. *Science* 161, 891-893.
- Glass, B. P. 1968. Microtektites and the origin of the Australasian tektite strewn field, *Publ. 6, 22 pp.*, Cent. for Meteorite Stud., Ariz. State Univ., Tempe.
- Glass, B. H. 1969. Chemical composition of Ivory Coast microtektites. *Geochem. Cosmochim. Acta* 33, 1135-1147, 1969.
- Glass, B. P., 1970. Zircon and chromite crystals in Muong nong-type tektite. *Science* 169, 766-769.
- Glass, B. P., 1970. Comparison of chemical compositions in flanged australite with chemical variation among normal Austral-Asian microtektites. *Earth and Planet Sci. Lett.* 1970. V.9, N. 3, p. 240-246.
- Glass, B. P., 1972, Australasian microtektites in deep-sea sediments, in Hayes, D.E., ed., *Antarctic oceanology, II. The Australasian-New Zealand sector: American Geophysical Union Antarctic Research Series*, v. 19, p. 335-348.
- Glass, B. P., 1972. Crystalline inclusions in a Muong Nong-type tektite. *Earth Planet. Sci. Lett.* 16, 23-26.
- Glass, B. P., 1976. No evidence for 0.8-0.9 m.y. old australite layer in deep sea cores. *Earth Planet. Sci. Lett.* 77, 428-433.
- Glass, Billy P. Zircon and Chromite Crystals in a Muong Nong-Type Tektite with comment. (in *Tektites*, p. 81-84, illus., Dowden, Hutchinson & Ross, Stroudsburg. 1973)
- Glass, B. P., 1979. The Ivory Coast microtektite- strewnfield. *Earth Planet. Sci. Lett.* 43. 336-342.
- Glass, B. P., 1979. Zhamanshin crater: possible source of Austral-Asian tektite (?). *Geology*, 1979. N. 7, p. 351-353.
- Glass, B. P., 1982a. *Introduction to Planetary Geology*. 469 pp., Cambridge University Press. New York. N. Y.
- Glass, B.P. 1982b. Possible correlations between tektite events and climatic changes? *Geol. Soc. Am. Spec. Pap.* 190: 251-256.
- Glass B. P. 1984. Solution of naturally - occurring glasses in the geological environment. *J. Non-crystal. Solids*. 67,285-286.
- Glass. B. R. 1986. No evidence for a 0.8-0.9 m.y. old microaustralite layer in deep-sea cores. *Earth Planet. Sci. Lett.* 77.428-433. 1986.
- Glass B. P. 1987. Coesite association with North American tektite debris in DSDP Site 612 in the continental slope off New Jersey. *Lunar Planet. Sci.* 18. 238-329 (Abstr.).
- Glass. B. P. 1988. Montagnais impact crater; possible source of the North American tektite Strewn field. (Abstracts Of Papers Submitted To The Lunar And Planetary Science Conference; 19(1), 391-392.
- Glass B. P. 1989. North American tektite debris and impact ejecta from DSDP Site 612. *Meteoritics* 24. 209-2 18.
- Glass, B. P. 1990: Tektites and microtektites: key fact and interferences. In: L. O. Nicolysen and W. U. Reimold (eds.), *Cryptoexplosions and catastrophes in the Geological Record with a Special Focus on the Vredefort Structure*. *Tectonophysics*, 171: 393-404.
- Glass B. P., 1993. Geographic variations in abundance of Australasian microtektites: Implications concerning the location and size of the source crater. *Meteoritics* 28, 354 (abst.).
- Glass, B. P. 2003. Australasian microtektites in the South China Sea: implications regarding the location and size of the source crater, *Lunar Planet. Sci. Conf.* 34 (Abstract no. 1092).
- Glass, B. P. and Barlow, R. A., 1979. Mineral inclusions in Muong Nong-type indochinites: Implications concerning parent material and process of the formation. *Meteoritics* 14, 55-67.
- Glass, B. P. and Heezen, B. C., 1967. Tektites and geomagnetic reversals. *Sci. Am.*, 33-38.
- Glass, B. P., and Koeberl, C. (2006), Australasian microtektites and associated impact ejecta in the South China Sea and the Middle Pleistocene supereruption of Toba. *Meteoritic and Planet. Sci.*, 41. 305-326.
- Glass, B.P., Folco, P., and Campanale, F., 2020 Coesite in a Muong Nong-type tektite from Muong Phin, Laos: Description, formation, and survival *Meteoritics & Planetary Science* 1–21, doi: 10.1111/maps.13433
- Glass, B.P., Fredriksson, K. and Florensky, P.V., 1983. Microirghizites recovered from a sediment sample from the Zhamanshin Impact Structure. 88(S01): B319-B330.
- Glass, B. P., Huber, H., and Koeberl, C., 2004. Geochemistry of Cenozoic microtektites and clinopyroxene-bearing spherules: *Geochimica et Cosmochimica Acta*.v. 68, p. 3971-4006, doi: 10. 10 16/j. gca.2004.02.026.
- Glass, B. P., Muenow, D. W. and Aggrey, K. E., 1986. Further evidence for the impact origin of tektites. *Meteoritics* 21, 369-370.
- Glass, B. P., D. R. Chapman and M. S. Prasad. 1996. Ablated tektite from the central Indian Ocean. *Meteoritics & Planetary Science* 31, 365-369.
- Glass, B. P. and Pizzutto, J. E., 1994. Geographic variation in Australasian microtektites concentration: Implication concerning the location and size of the source. *J. Geophys. Res.* 99 (E9), 9,075- 19,081.
- Glass, B. P., Swincki, M. B. and Zwart, P.A., 1979. Australasian, Ivory Coast and North American tektite strewn fields: Size, mass and correlation with geomagnetic reversals and other earth events. *Proc. Lunar Planet. Sci. Conf.* 10th, 2535-2545.
- Glass, B. P. and Koeberl, C., 1989. Trace element study of high- and low- refractive index Muong Nong-type tektites from Indochina. *Meteoritics* 24, 143-146.

- Glass, B. P., and C. Koeberl, 2006, Australasian microtektites and associated impact ejecta in the South China Sea and the Middle Pleistocene supereruption of Toba, *Meteorit. Planet. Sci.*, 41, 305-326.
- Glass, B. P. and Zwart, M. J., 1979. North American Microtektites in Deep Sea Drilling Project cores from Caribbean Sea and Gulf of Mexico. *Bull. Soc. Am.*, 90: 595-602.
- Glass, B. P. and Wu, J. 1993. Coesite and shocked quartz discovered in Australasian and North America microtectes layers. *Geology* 21: 435-438.
- Glass, B.P., and Koeberl. C., 2006, Australasian microtektites and associated impact ejecta in the South China Sea and the Middle Pleistocene supereruption of Toba: *Meteoritics and Planetary Science*, v. 41, p. 305-326.
- Glass, B.P. and Koeberl, C., 2006, Australasian microtektites and associated impact ejecta in the South China Sea and the Middle Pleistocene supereruption of Toba: *Meteoritics and Planetary Science*, v. 41, p.305-326.
- Glasstone, S., and Doll, P. J., 1977, *The effects of nuclear weaponr (3rd edition)*: U. S. Department of Defense.
- Gobbett, D. J. and Tjia, H. D., 1973. *Tectonic history Geology of the Malay Peninsula* (ed. D. J. Gobbett and C. S. Hutchison), pp. 305-30. John Wiley Interscience, New York.
- Grant-Taylor, T. L. and Rafter, T. A., 1962. New Zealand radiocarbon measurements. *N.Z. J. Geol. Geophys.* 5, 331-59.
- Grieve, R. A. F., 1982. The record of impacts on Earth: Implications for a major Cretaceous/Tertiary impact event. *Geol. Soc. Amer. Spec. Pap.* 190, 25-38.
- Grieve R.A.F. & Robertson, P.B. 1979. *Icarus* 38, 212-229.
- Grieve R.A.F., Dence M.R. & Robertson P.B. 1976. In: *Impact and Explosion Cratering* (Roddy, D.J., Pepin, R.O., and Merrill, R.R., eds) Pergamon, New York, 791-834.
- Grossman J. N. and Baedeker P. N., 1986. Computer graphics for quality control in the INAA of geological samples. In *Proc. 7Th Intl. Conf. on Modern Trends in Activation Analysis*, pp. 571-578.
- Grote P. 2007. Studies of fruits and seeds from the Pleistocene of northeastern Thailand. *Courier Forschungsinstitut Senckenberg* 258: 171-181.
- Guinasso, N.L. and Schink, D.R., 1975. 'Quantitative estimates of biological mixing rates in abyssal sediments', *J. Geophys. Res.*, 80, 3032-3043.
- Haaq, B.U. 1998. Gas hydrates: greenhouse nightmare? energy panacea or pipe dream? *GSA Today*, v. 8, pp. 1-6.
- Haile, N., 1980. Palaeomagnetic evidence from the Ordovician and Silurian of northwest Peninsular Malaysia. *Earth Planet. Sci. Letts.*, 48: 233-236.
- Haile, N. S. and Briden, J. C. 1982. Past and future palaeomagnetic research and tectonic history of East and Southeast Asia. Palaeomagnetic research in Southeast and East Asia, CCOP Technical publication 13, pp. 25-46. United Nations Building, Bangkok, pp. 25-46.
- Haile, N. S., Stauffer, P. H., Krishnan, T P. and Ong, G. B. 1977. Palaeozoic redbeds and radiolarian chert: reinterpretation of their relationships in the Bentong and Raub areas, West Pahang, Peninsular Malaysia. *Bull. geol. Soc. Malaysia*, 8: 45-60.
- Haile N. S., M.W. McElhinny and I. McDougall., 1977. Paleomagnetic data and radiometric ages from the Cretaceous of West Kalimantan (Borneo), and their significance in interpreting regional structure, *J. Geol. Soc. London* 133 (1977) 133-144,
- Haile, N. S., Wong, N. P. Y., and Nuttall, C. P. 1985. The geology and mineral resources of Dent Peninsula, Sabah. Geological survey of the Borneo region, Malaysia, memoirs 16.
- Haile, N. S., Beckinsale, R. D., Chakraborty, K. R., Abdul Hanif Hussein, and Hardjono, T. (1983). Palaeomagnetism, geochronology and petrology of the dolerite dykes and basaltic lavas from Kuantan, West Malaysia. *Geological Society of Malaysia Bulletin* 16, 71-85.
- Haines, P. W., K. T. Howard, J. R. Ali, C. F. Burrett and Sangad Bunopas, 2004. Flood deposits penecontemporaneous with ~0.8 Ma tektite fall in NE Thailand: impact-induced environmental effects? *Earth and Planetary Science Letters* 225(1), 19-28.
- Hamano, Y. 1980. An experiment on the post-depositional remanent magnetization in artificial and natural sediments. *Earth Planet. Sci. Lett.* 51, 221-232. 1980.
- Hamburger, M.W., Cardwell, R. K., and Isacks, B. L., 1983. *Seismotectonics of the northern Philippine island arc. The tectonic and geologic evolution of Southeast Asian seas and islands, part 2, Geophysical monograph 27* (ed. D. Hayes), pp. 1-22. American Geophysical Union, Washington.
- Hamilton, W. B. 1979. *Tectonics of the Indonesian region. United States Geological Survey Professional Paper* 107\$, 1-345.
- Hamilton, W. B. 1988. Plate tectonics and island arcs. *Geological Society of America Bulletin* 100, 1503-27.
- Han, J.M., 1991. Palaeoclimatic impact on the magnetic and stable isotope characteristics of the Chinese loess. Ph.D. Thesis of Vrije Universiteit Brussel, Belgium.
- Han, J.M., Jiang, W.Y., Liu, T.S., Li, H.Y., Guo, Z.T. & Wu, N.Q., 1996. Carbonate isotopic records of paleoclimate changes in Chinese loess. *Science in China* 39, 460-467.
- Haneda, Y., Okada, M., and Kitamura, T., 2020. A full sequence of the Matuyama-Brunhes reversal in the Chiba composite section, Central Japan. *Progress in Earth and Planetary Sciences*. 7 no.44, doi: [10.1186/s40645-020-00354-y](https://doi.org/10.1186/s40645-020-00354-y)
- Han Jiamao, Jiang Wenying and Chu Jun. 1997: *Stratigraphy of the Chinese Loess. Proc. Intern. Conf. on Stratigraphy and Tectonic Evolution of Southeast Asia and Associated Meetings of IGCP 359 and IGP 383. Siam City Hotel, Bangkok, August 1977. Vol. 1 & 2 : 223-231.*
- Hanta, Rattanaphorn; B. Ratanasathien; Y. Konimatsu; H. Saegusa; H. Nakaya; P. Chintasakul. 2005. Description of the Tha Chang Merycopotamus and Its Preserved Condition. *International Conference on Geology, Geotechnology and Mineral Resources of Indochina (GEOINDO 2005)*, pp. 600-605. 28-30 November 2005, Khon Kaen, Thailand
- Hanta, Rattanaphorn; Benjavun Ratanasathi, Tanaka Satoshi, Yutaka Kunimatsu, Hideo Nakaya. 2007. A Depositional Environment of Late Miocene Mammalian Fossil, Sop Mae Tham, Lampang Province, Northern Thailand., *GEOTHAI'07 International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy*
- Harder, S. H., Mauri, S.J., and Marshall-Arrazola, S., 1993. Gravity modelling of extensional basins in southeast Asia. *Bull. Geol. Soc. Malaysia* 33, 153-162.
- Harbury, N. A., Jones. M. E., Audley-Charles, M. O.,

- Metcalf, I., and Mohamed, K. I. 1990. Structural evolution of Mesozoic Peninsular Malaysia. *Journal of the Geological Society, London* 147, 11-26.
- Harris, R. A. 1990. Time variant distribution of strain in the active Banda orogen: a reconciliation of rival hypotheses. in 'Orogenesis in Action. Symposium at the Geological Society, London, 15-20th April. 1990'. (Eds R. Hall, G. Nichols and C. Rangin.)
- Harrison, T. M., P. H. Leloup, F. J. Ryerson, P. Tapponnier, R. Lacassin, and Chen Wenji, 1996. Diachronous initiation of Transtension along the Ailao Shan-Red River Shear zone, Yunnan and Vietnam, in: *The Tectonics of Asia*, edited by Harrison, T. M., and An Yin, pp. Cambridge University Press, New York, 1996.
- Hartung, J. B., 1990. Australasian tektite source crater? Tonle Sap, Cambodia. *Meteoritics* 25(4), 369-370.
- Hartung, J. B. and Rivolo, a. R. 1979. A possible source in Cambodia for Australasian tektites. *Meteoritics* 14, 153-160.
- Hartung, J. B. "Australasian tektite source crater? Tonle Sap, Cambodia." 1990. (METEORITICS :Voi.25, No.4, p. 369-370)
- Hartung, J. B., et al., 1994. In search of the Australian tektite source crater; the Tonle Sap hypothesis. *Meteoritics* 29 (3), 411-416.
- Hashimoto, V. and Sato, T. 1968. Contribution to the geology of Mindoro and neighbouring islands, the Philippines. *Geology and paleontology of Southeast Asia 5* (ed. T. Kobayashi et al.), pp. 192-220. University of Tokyo Press.
- Hashimoto, W. and Sato, T. 1973. Geological structure of north Palawan and its bearing on the geologic history of the Philippines. *Geology and paleontology of Southeast Asia 13* (ed. T. Kobayashi et al.), pp. 145-61. University of Tokyo Press.
- Hatherton, T. and Dickinson, W. R. 1969. The relationship between andesitic volcanism and seismicity in Indonesia, the Lesser Antilles, and other island arcs. *Journal of Geophysical Research* 74, S301-9.
- Hatton, F. 1885. *North Borneo: explorations and adventures on the Equator*. Samson, Low, Marston, Searle and Rivington, London.
- Hawkesworth, C.J., and Gallagher, K., 1993. Mantle hot spots, plumes and regional tectonics as causes of intraplate magmatism. *Terra Nova*, 5, nr. 6, 552-559.
- Hawkins, J. W. and Evans, C. A. 1983. *Geology of the Zambales Range, Luzon, Philippine islands: ophiolite derived from an island arc - back arc basin pair. Tite tectonic and geologic evolution of Southeast Asian seas and islands, part 2, Geophysical monographs 27* (ed. D. E. Hayes), pp. 95-123. American Geophysical Union, Washington.
- Hayes, D. E. and Taylor, B. 1978. *Tectonics. A geophysical atlas of the East and Southeast Asian seas (5 sheets) MC-25* (director, D. E. Hayes). Geological Society of America.
- Hehuwat, F. 1976. Isotopic age determinations in Indonesia: the state of the art. proceeding of isotope geology, Technical publication 3, pp. 135-57. U.N. ESCAP/CCOP, Bangkok.
- Heim, A. and Hirschi, H. 1939. A section of the mountain ranges of north-western Siam, *Ecolgae Geologicae Helvetiae* 32, No. 1.
- Heins, W. A.; An, Z.; Zhang, X.; Li, C.; Ouyang, Z. and Wasson, J. T., 1991. Search for microtektites and associated noble metals in North China Loess (abstract), INQUA Conference Beijing, p. 132.
- Heller, F. & Liu, T.S., 1982. Magnetostratigraphical dating of loess deposits in China. *Nature* 300, 43 1-433
- Heirtzler, J. R., et al. 1978. The Argo abyssal plain. *Earth and Planetary Science Letters* 41,2t-31.
- Hennecke, E. W., Manuel, O. K. and Sabu, D. D., 1975. Noble gases in Thailand tektites. *J. Geophys. Res.* 80, 2931-2934.
- Hennecke, E. W., and O. K. Manuel, 1980. Mass fractionation and the isotopic anomalies of xenon and krypton in ordinary chondrites. *Z.Naturforsch.* .1, 26, 1980-1986, 1971.
- Herberer, G., 1968. *Der Ursprung des Menschen*. Gustav Fischer, Stuttgart, 43p.
- Herman, B. M., Anderson, R. N., and Truchan, M. (1979). Extensional tectonics in the Okinawa Trough. *Geological and geophysical investigations of continental margins, American Association of Petroleum Geologists memoirs 29* (ed. J. S. Watkins, L. Montadert, and P. f. Dickerson), 199-208.
- Hermes, J. J. 1974. West Irian. Mesozoic-Cenozoic orogenic belts, *Geological Society of London Special Publication 4* (ed. A. M. Spencer), 475-90. Hiller, K. and Elahi, M. 1984. Structural development and hydrocarbon entrapment in the Burma Basin/Bangladesh (Northwest Indo-Burman fold belt). 5th offshore Southeast Asia conference proceedings s.E.A.P.E.X., 5.50-6.53
- Hildebrand, A. R., et al., 1994. Phnum Voene; source crater for Australasian strewnfield?. Program with Abstracts-Geological Association of Canada; Mineral Association of Canada; Canadian Geophysical Union, Joint meeting; v. 19, p. 50.
- Hills, J. G., 1981 Comet showers and the steady-state infall of comets from the Oort cloud. *Astron. J.*, 86, 1730-1740.
- Hills, J. G., and Goda, M. P, 1993, The fragmentation of small asteroids in the atmosphere: *Astronomy Journal*, v. 1 05, p. 1114-1144.
- Hilton, D. R., and Craig, H., 1989. A helium isotope transect along the Indonesian archipelago. *Nature* 343, 906-908.
- Hilton, D. R., Grönvold, K., O'Nions, R.K., and Oxburgh, E. R., 1990. Regional distribution of <sup>3</sup>He anomalies in the Icelandic crust. *Chem. Geol.* 88, 53-67.
- Hilton, D. R., Hammerschmidt, K., Loock, G., and Friedrichsen, H., 1993. Helium and argon isotope systematics of the central Lau Basin and Valu Fe Ridge: evidence of crust\mantle interactions in a back-arc basin. *Geochim. Cosmochim. Acta* 57, 2819-2841.
- Hilton, D. R., Hoogewerff, J.A., Van Bergen, M.J., and Hammerschmidt, K., 1992. Mapping magma sources in the east Sunda-Banda arcs, Indonesia: constraints from helium isotopes. *Geochim. Cosmochim. Acta* 56, 851-859.
- Hirschi, H. and Heim, A. 1938. Zur geologie und petrographie von Nord-Siam. *Schweizer mineralogisch petrographisch Mitteilungen* 18 (2).
- Ho, C. S. 1986. A synthesis of the geological evolution of Taiwan. *Tectonophysics* 125, 7-16.
- Ho, C. S. and Lee Chinnan, 1953. Chapters on geology of Taiwan. *Economic minerals of Taiwan* 1, pp. 3-57. Geological Survey of Taiwan.
- Hoke, L., 1997. Origin of CO<sub>2</sub> in Hydrothermal Spings in Vietnam and Thailand. *Proc. Intern. Conf. on Stratigraphy and Tectonic Evolution of Southeast Asia and*

- Associated Meetings of IGCP 359 and IGP 383. Siam City Hotel, Bangkok, August 1997. Vol. 1 & 2 : 726.
- Hoke, L. and H. J. Campbell, 1995 Active Mantle Melting Beneath Thailand? International Conference on Geology. Geotechnology and Mineral Resources of Indochina (Geo-Indo'95), 22-25 November 1995, KhonKaen, Thailand.
- Hoke, L. and Hoang Huu Quy, 1995. Active mantle processes beneath Vietnam. Journal of Geology, special issue, Series B, no. 5-6 (1995), 309-310
- Hoke, L., Hilton, D.R., Lamb, S.H., Hammerschmidt, K. and Friedrichsen, H., 1994. "The evidence for a wide zone of active mantle melting beneath the central Andes. Earth Planet. Sci. Lett. 128, 341-355.
- Holcombe, C. J. 1977a. How rigid are the lithospheric plates? Fault and shear rotations in Southeast Asia. Journal of the Geological Society London 134, 325-42.
- Holcombe, C. J. 1977b. Earthquake foci distribution in the Sunda arc and the rotation of the back arc area. Tectonophysics 43, 1 69-80.
- Holloway, N. H. 1981. The North Palawan Block, Philippines: its relation to the Asian mainland and its role in the evolution of the South China Sea. Geological Society of Malaysia Bulletin 14, 79-58.
- Hou, Y., Potts, R., Baoyin, Y., Zhengtang, G., Deino, A., Wang, W., Clark, J., Guangmao, X., & Weiwen, H., 2000. Mid-Pleistocene Acheulean-Like Stone Technology of the Bose Basin, South China. Science, 287, 1622-1626.
- Howard, K. T. 1999: Distal environmental effects of the 770ka impact event. Unpublished Hons. Thesis at the University of Tasmania, Australia. 175 pp.
- Howard, K. T. 2004: Origin of Darwin Glass. Unpublished Ph. D. Thesis at the University of Tasmania, Australia. 362 pp.
- Howard, K.T., 2011. Tektites in: Ridd, M.F., Barber, A.J., Crow, M.J. (Eds.), The Geology of Thailand, The Geological Society of London, pp. 574-591.. DOI:https://doi.org/10.1144/GOTH.21
- Howard, K. T., S. Bunopas, F. Burrett, P.W. Haines, M.D. Norman, The 770 ka tektite producing impact event: evidence for distal environmental effects in NE Thailand, Lunar Planet., Sci. Conf. 31 (Abstract no. 1308).
- Howard, K. and Maloney, 2000. Pollen sum, total pollen, and pteridophyte spore graminiae and sample from section 1 and 2 at Ban Tha Chang. Newsletter 2000, Geological Society of Thailand
- Howard, K. T., P.W. Haines. C. F. Burrett, J. R. Ali, S. Bunopas, 2003. Sedimentology of 0.8 Ma log-bearing flood deposits in northeast Thailand and mechanisms for pre-flood deforestation, Proceedings, 8th International Congress on Pacific Neogene Stratigraphy, Chang Mai, Thailand, 2003, pp. 49 -67.
- Hsü, K., J., 1980 Terrestrial catastrophe caused by cometary impact at end of Cretaceous. Nature, London, 285, 201-203.
- Hsü, K., J. 1981 Origin of geochemical anomalies at Cretaceous-Tertiary boundary: Asteroid or cometary impact? Oceanologica Acta, no. SP, 129-133
- Hsü, K., J. 1983a. Actualistic catastrophism; Address of the retiring president of the International Association of Sedimentologists. Sedimentology, 30, 309.
- Hsü, K., J., 1983b. Geochemical markers of impacts and of their effects on environments. In H. D. Holland et al. "Patterns of Change in Earth Evolution", Springer-Verlag, New York, 63-74.
- Hsü, K., J., 1986. Three mistakes of Darwinism. Geology, 14.
- Hsü, K. J. Mckenzie, J. A. and He, Q. H., 1982. Terminal Cretaceous environmental and evolutionary changes. Geol. Soc. Amer., Spec. Pap., 190, 317-323.
- Hsü, K. J. & Mckenzie, J. A 1985 A "Strangelove" ocean in the earliest Tertiary. in E. T. Sundquist and W. S. Broecker (eds.), "The Carbon Cycle and Atmospheric CO<sub>2</sub>: Natural Variations from Archean to Present", Am. Geoph. Union, Geoph. Monograph, 32, 487-402.
- Hsü, K. J. & Mckenzie, J. A. 1986 Rare events in geology discussed at meeting. Geotimes (March), 11-12.
- Hsü, K. J., Obrhansli, H. & Gao, J. Y., 1985. A "Strangelove Ocean" before the Cambrian Explosion. Nature, London, 316, 809-811.
- Huchon, P., X. Le Pichon, and C. Rangin, 1994. Indochina peninsula and the collision of India and Eurasia, Geology, 22, 27-30, 1994.
- Huchon, P., Barrier, E., De Bremaecker, J. C., and Angelier, J. 1985. Collision and stress trajectories in Taiwan: a finite element model. Tectonophysics 125, 179-91.
- Hughes, D.W. . 1985.. Quart. J. Rov. Astron. Soc. 16, 410-427
- Hut. P., Alvarez, W., Elder, W., Hansen, T. A., Kauffman, E. O., Keller, O., Shoemaker E. M. and Weissman, P., 1987). Comet showers as a cause of mass extinction. Nature, London, 329, 118-126.
- Hutchison, C.S. 1972. Alpine-type chromite in North Borneo, with special reference to Darvel Bay. American mineralogist 57, 83 5-5 6.
- Hutchison, C. S. 1973a. Plutonic activity. Geology of the Malay Peninsula (ed. D. J. Gobbett and C. S. Hutchison), pp. 215-52. John Wiley-Interscience, New York.
- Hutchison, C. S. 1973b. Metamorphism. Geology of the Malay Peninsula (ed. D. J. Gobbett and C. S. Hutchison), pp. 253-303. John Wiley-Interscience, New York.
- Hutchison, C.S. 1973c. Tectonic evolution of Sundaland: a Phanerozoic synthesis. Geological Society of Malaysia Bulletin 5, 61-86.
- Hutchison, C. S. 1975a. Ophiolite in Southeast Asia. Geological Society of America Bulletin 86, 797-806.
- Hutchison, C. S. 1975b. Correlation of Indonesian active volcanic geochemistry with Benioff zone depth. Geologie en Mijnbouw 54, 157-68.
- Hutchison, C. S. 1975. Indonesian active volcanic arc: K, Sr and Rb variation with depth to the Benioff zone. Geology 4, 407-8.
- Hutchison, C. S. 1977. Granite emplacement and tectonic subdivision of Peninsular Malaysia. Geological Society of Malaysia Bulletin 9, 187-207 .
- Hutchison, C. S. 1978a. Southeast Asian tin granitoids of contrasting tectonic setting. Journal of the Physics of the Earth 26, supplement, S221-32.
- Hutchison, C. S. 1978. Ophiolite metamorphism in northeast Borneo. Lithos 17, 195-208.
- Hutchison, C. S. 1982a. Southeast Asia. The Ocean basins and margins: The Indian Ocean (ed. A. E. M. Nairn and F. G. Stehli), 6, pp. 451-512. Plenum Press, New York.
- Hutchison, C. S. 1977. Granite emplacement and tectonic subdivision of Peninsular Malaysia. Geological Society of Malaysia Bulletin 9, 187-207.
- Hutchison, C. S. 1985. A re-assessment of the evidence that the S.E. Asian shear basins resulted from the India-Tibet collision. Geological Society of Malaysia

- Petroleum and Geology seminar '85, Kuala Lumpur, Dec. 1985.
- Hutchison, C.S., 1989. Geological evolution of South-East Asia. Oxford Monographs on Geology and Geophysics, No. 13, Clarendon Press, Oxford, 368pp.
- Hutchison, C. S. 1996. Geological Evolution of South-East Asia. Geological Society of Malaysia. 368 pp., Geological Society of Malaysia, c/o Department of Geology, University of Malaya, 50603 Kuala Lumpur, Malaysia.
- Hutchison, C. S and Jezek, P. A. 1978. Banda arc of eastern Indonesia: petrography, mineralogy and chemistry of the volcanic rocks. Proceedings of the third regional conference on geology and mineral resources of S.E. Asia (ed. Prinya Nutalaya), PP. 607-19. Asian Institute of Technology, Bangkok
- Institute of Atomic Energy, Academua Sinica et al., 1976. Fission Tract Dating of Muscovite and Tektites. Peking, no.2, 148-156.
- IHEG (Institute of Hydrogeology and Engineering Geology, Chinese Academy of Geological Sciences) 1959. Loesses and loessy rocks in China. Publishing House of Geology, Beijing.
- Itihara, M., et al. 1985. Concluding Remarks. In Watanabe, N., Kadar, D., eds. 1985. Quaternary Geology of the Hominid Fossil Bearing in Java. Geol. Res. Devel. Centre Sp. Paper No. 4. 378p. Bundung.
- Izett, G. A.; Dalrymple, G. B. and Snee, L. W., 1991. 40Ar / 39Ar Age of Cretaceous-Tertiary boundary tektites from Haiti. Science, 252, 1539-1542.
- Izett, G. A. and Obradovich J. D. 1992: Laser-fusion 40Ar/39Ar ages of Australian tektites (abstract). Luna Planet. Sci. 23: 593-594.
- Izett, Glen A., et al. "Laser-fusion 40Ar-39Ar ages of Australasian tektite cores and f1anges." Oct. 27, 1992. (EOS, TRANSACTIONS, AMERICAN GEOPHYSICAL LINION ; Vol. 73, No. 43, Suppl., p. 328)
- Izokh, E. P., 1985. Age paradox of tektites and the field where they fell. Meteoritika, 44. 127-134.
- Izokh, E. P. and An, L. D., 1983. Tektites of Vietnam: Tektites delivered by a comet - A hypothesis (in Russian). Meteoritika 42, 158-170.
- Izokh, E..P. (1985): The age-patadox of tektites and their strewn-fields.-Meteoritika, 44,127-134, (In Russian).
- Izokh, E..P. (1987): The petrochemical properties of rhe target-rocks, impactites and tektites-of zhamanshine -astroblerm. - In: Kosmicheskoe veshchestwo i Zemlya, Novosibirsk, "Nauka". 159-203. (In Russian).
- Izokh, E. P. (1988), Age-paradox and the origin of tektites, in Proceedings of the Second International conference on Natural Glasses, edited by V. Bouska. pp. 379-384. Charles Univ. Press, Prague.
- Izokh, E. P., Izokh, Ye. P. "Zhamanshin impact crater and tektite problem." 1991. (Soviet Geology And Geophysics ; Vol. 32, No.4, p. 1-10)
- Izokh, Emil. "Zhamanshin impact crater and the tektite puzzle." Dec. 1991. (METEORITICS ; Vol. 26. No. 4, p. 350)
- Izokh, E. P. "Relationship between Austral-Asian tektite strewn field and ZhamanshinCrater; clue to the origin of tektites." Wasson, John T. Dec. 1989. (METEORITICS; Vol.24, No.4. p. 280)
- Izokh, E. P. and AN, L. D. 1988. The geological position of the tektites and their significance for the Quaternary geology and geomorphology of Vietnam. Aktual' nie Voprosi Meteoritiki v Sibiri, Novosibirsk, "Nauka", 1988., Translated on behalf of J. O 'Keefe by SCITRAN,, Santa Babara CA 93150., corrected by E. Izokh : pp. 31-57.
- Jacks, G. V., 1963. Soil. Tomas Nelson & Sons Ltd. London. 221 pp.
- Jezek, P.A. and Hutchison, C. S. 1973. Banda arc of eastern Indonesia: petrography and geochemistry of the volcanic rocks. Bulletin of Volcanology'41, 584608.
- Jezek, P. A., Whitford, D. J., and Gill, J. B. 1931. Geochemistry of Recent lavas from the Sangihe-Sulawesi arc, Indonesia. The geology and tectonics of eastern Indonesia (ed. A. J. Barber and S Wiryosulino), pp. 383-9. Geological research and development centre, Bandung.
- Jhakkaphak, P., and Veerabarus, M., 1982, Geoiogic map of Thailand: Bangkok, Deparknent of Mineral Resources, scale 1:100,000.
- Jin, Chengwei and Xu Ronghua. 1984. Les granitoids de la partie centrale de l'Himalaya et du Gindise au Xizang (Tibet) meridional. Mission Franco-Chinoise au Tibet 1980 (ed. J. L. Mercier and Li Guangcen), pp. 289-308. Editions du centre national de la recherche scientifique, Paris,
- Jiongxin,X., 1991. Astudy of river channel pattern in formation recorded by grainsize parameters of fluvial sediment: Earth Surface Processes and Landforms, v. 16, p.129-142.
- Johnson, B. D., Powell, C. McA., and Veevers, J. J. 1976. Spreading history of the eastern Indian Ocean and greater India's northward flight from Antarctica and Australi a. Geological Society of America Bulletin 87, 1550-6.
- Johnston, C. R. 1981. A review of Timor tectonics with implications for the development of the Banda arc. The geology and tectonics of eastern Indonesia (ed. A. J. Barber and S. Wiryosujono), pp. 199-216. Geological Research and Development Cenrre, Bandung.
- Jones, K. M., Bacon .M. and Hastings. D.A.. 1981. The Lake Bosurntwi impact crater. Ghana. Geol. Soc. Am. Bull. 92. 342-349
- Jones, W. B. 1985. The origin of the Bosumtwi crater, Ghana-an historical review, Proc. Geol. ,Assoc. 96(3), 275-284, 1985.
- Johnson, B. D., Powell, C. McA., and Veevers, J. J. (1976). Spreading history of the eastern Indian Ocean and greater India's northward flight from Antarctica and Australia. Geological Society of America Bulletin 87, 1560-6.
- Jourdan F., Nomande S., Wingate M. T. D., Eroglu E., and Deino A. 2019. Ultraprecise age and formation temperature of the Australasian tektites constrained by 40Ar/39Ar analyses. Meteoritics & Planetary Science 54, 2573–2591. <https://doi.org/10.1111/maps.13305>
- Kaewyana, W., 1980. Late Quaternary Alluvial Terraces And Their Cover Bed Stratigraphy, Eketahuna And Pahitatu District, New Zealand. Thesis submitted for the degree of M. Sc., Victoria University of Wellington, New Zealand. 398 p. (copies lodged at Victoria University Library, and at DMR., Bangkok.)
- Kallemeyan, G. W. and Wasson, J. T., 1981. The atu classification of chondrites-I. The carbonaceous chodritic group. Geochim. Cosmochim. Acta 45, 1217-1230.
- Kamineni, D. C. and Chakraborty, K. R. 1975. Traceelement characteristics of Segamat volcanics. Neusletter of the geological society of Malaysia 4, tt5-22.
- Kamiya, H. and K. Kato, 2003. Comparison of Enamel

- Texture of Fossil Elephant Molars from Japan and its Neighbors, with Special Notes on Modern Elephants. P. 121-125.
- Karig, D. E. 1971. Origin and development of marginal basins in the western Pacific. *Journal of geophysical research* 76, 25 42-61.
- Karig, D. E. 1972. Remnant arcs. *Geological society of America bulletin* 83, 1057-58.
- Karig, D. E. (1973). Plate convergence between the Philippines and the Ryukyu islands. *Marine geology* 14, 153-58.
- Karig, D. E., Lawrence, M. B., Moore, G. F., and Curray, J. R. (1980). Structural framework of the fore-arc basin, N.W. Sumatra. *Journal of the geological society London* 137,77-91.
- Kashkarov, L. L., Genaeva, L. I. And Izokh, E.P., 1987. Fission track dating of glasses from Zhamanshin astrobleme. 2nd Intern. Conf. On Natural Glasses. Prague. 199-202.
- Katili, J. A. (1974a). Sumatra. Mesozoic-Cenozoic orogenic belts, Geological Society of London special publication 4 (ed. A. M. Spencer), 317-31.
- Katili, J. A. (1974b). Volcanism and plate tectonics in Indonesian island arc. *Tectonophysics* 26, 155pp.
- Katili, J. A. (1978). Past and present tectonic position of Sulawesi, Indonesia. *Tectonophysics* 45, 289-322.
- Katili, J. A. (1984). Tectonic framework, resources, occurrences and related problems in Southeast Asia. Southeast Asia: tectonic framework, earth resources and regional geological programs, International Union geological sciences publication 13 (ed. I. A. Katili and J. A. Reinemundj), 1g-5S.
- Kawamura, Y., 1998. Immigration of Mammals in Japanese Islands during the Quaternary. *The Quat. Res.*, 37 (3): 25 1-257.
- Kauffman, E. G. 1984. Toward a synthetic theory of mass extinction. *Soc. Amer. Ast. Progr.* 16, 555.
- Kear, D. and Waterhouse, B. C., 1961. Quaternary surfaces of sediment at Waihi Beach. *N. Z. J. Geol. Geophys.* 4, 434-45.
- Keller, G., D'Hondt, S.L. and Valler, T. L., 1983. Multiple microtektite horizons in Upper Eocene marine sediments: No evidence for mass extinction. *Science*, 221, 150-151.
- Keller, G., D'Hondt, S.L., Orth, C.J., Gilmore, S. S., Oliver, P. Q., Shoemaker, E.M., and Molina, E., 1987. Late Eocene impact microspherules: Stratigraphy, age and geochemistry: *Meteoritics*, v. 22, p. 25-60.
- Kent, D. V., 1973. 'Post-depositional remanent magnetisation in deep-sea sediment', *Nature*, 246, 32-34.
- Kes, A.S., 1959. The question of the origin of loess in Northern China, in: *Loess of North China* (eds. Popob, et al., 85-105, Israel Prog. Sd. Trans.
- Khin Zaw and C. F. Burrett, 1997. Cambrian volcanic hosted massive sulphide mineralisations in Australia and Southeast Asia: Are they related to rifting of northern Gondwanaland. *The International Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific Bangkok, Thailand*, 19-24 August 1997
- Kieffer/ S.W. & Simonds, C.H. 1980. *Proc. Lunar Planet in Sci. Conf.* 18th, L43-181.
- King, E. A., 1977. Terrestrial origin of tektites. *American Scientist* 64, 212-218.
- Kiang, Sia Hok, 1978. Tektites found on beach near Endau, Johore, peninsular Malaysia. *Warta Geologi: Newsletter of the Geol. Soc. Malaysia*; 4 (5), 153- 155.
- Kilham, P., 1971. A hypothesis concerning silica and the fresh water planktonic. *16(1)*, 10-18.
- Kirk, H. J. C. 1957. The geology and mineral resources of the Upper Rajang and adjacent areas. Geological Survey Department for the British Territories in Borneo, Memoir 8.
- Kirk, H. J. C. 1958. The igneous rocks of Sarawak and Sabah. Geological survey of Borneo region Malaysia bulletin 5.
- Kita, I., Nagao, K., Takashima, I., Honda, S., and Ratanasthien, B., 1990. A preliminary study on the chemical and isotopic characteristics of geothermal waters and gases in northern Thailand. *Mass Spectro.* 38, 295-305.
- Kizaki, K. 1978. Tectonics of the Ryukyu island arc. *Journal of Physics of the Earth* 26 Supplement, s301-7.
- Kizaki, K. 1985. Geology and tectonics of the Ryukyu islands. *Tectonophysics* 125, 193--207
- Klompé, T. H. F. 1962. Igneous and structural features of Thailand. *Geologie en mijnbouw* 41, 290\_302.
- Klootwijk, C. T. and Radhakrishnamurthy, C. 1981. Phanerozoic palaeomagnetism of the Indian Plate and the India-Asia collision. *Palaeoreconstruction of the continents, Geodynamics Series* 2,93-t05.
- Klongsara, N., 2002. Thermoluminescence dating of tektites from Ban Mai, Tambon Ta Chang, Amphoe Chalerm Phrakiet, Changwat Nakhon Ratchasima. An unpublished B.Sc. report, Department of Geology, Chulalongkorn University, 62 p.
- Knight, M.D., Walker, G. L. P., Ellwood, B. B., and Diehl, J. F. (1986). Stratigraphy, paleomagnetism, and magnetic fabric of the Toba Tuffs, constraints on the sources and eruptive styles. *Journal Geophysical Research* 91, 10355-82.
- Knittel, U. (1979). Petrography and origin of the syenite from Cordón, Isabela. *Geological Society of the Philippines Journal* 33, 3H3.
- Koeberl C., 1986. Geochemistry of tektites and impact glasses. *Annu. Rev. Earth Planet. Sci.* 14,323-350.
- Koeberl, C. 1988. *Proc. Nat. Inst. Polar Res. Svmp. L*, 261-290.
- Koeberl C., 1992. Geochemistry and origin of Muong Nong-type tektites. *Geochim.Cosmochim.Acta*, 56, 1033-1064.
- Koeberl C., 1993. Extraterrestrial component associated with Australasian microtektites in a core from ODP site 758B. *Earth Planet. Sci. Lett.* 119, 453-458.
- Koeberl C., 1994. Tektite origin by hypervelocity asteroid or cometary impact: Target rocks, source craters and mechanism. In *Large Meteorite Impact and Planetary Evolution* (eds. B. O. Dressler, R. A. F. Grieve and V. L. Sharpton), pp. 133-151. *Geol. Soc. Am. Spec. Pap.* 293, Boulder, Colorado.
- Koeberl. C. et al. 1984 "Geochemistry of Muong-Nong type tektites; V, Unusual ferric/ferrous ratios." *METEORITICS : Vol. 19, No. 4*, p.253-254)
- Koeberl C. And Beran A. (1988) Water content of tektites and impact glasses and related chemical studies. *Proc. Lunar Planet Sd Coat* 18th, 403-408.
- Koeberl, C, Bottomley, R. J., Glass, B. P., Stozler, D. and York, D., 1989. Geochemistry and age of Ivory Coast tektites (Abstract). *Meteoritics* 24, 287.
- Koeberl, C.; Poag, C. W.; Reimold, W. U.; Brandt, D. 1996. "Impact Origin of the Chesapeake Bay Structure and the Source of the North American Tektites". *Science*.

- 271 (5253):1263-1266. doi:10.1126/science.271.5253.1263. S2CID 128672140.
- Koerberl, C., B. P. Glass, 2000. Tektites and the age paradox in mid-Pleistocene China, *Science* 289 (2000) 507.
- Koerberl, C., and Glass, B. P., 1988, Chemical composition of North American microtektites and tektite fragments from Barbados and DSDP Site 612 on the continental slope off New Jersey: *Earth and Planetary Science Letters*, v. 87, p. 286-296
- Koenigswald, G. H. R. Von. Tektite studies-(pt.) 10, the relationship of shape, size and texture in asiatic tektites. (Konfnkl. NEDERLANDSE AKAD. WETENSCH. PROC. 71. NO. 1. P I-9 1968)
- Koenigswald, G. H. R. von. Tektite studies, X; The relationship of shape, size and texture in Asiatic tektites. [Ned. Akad. Wetensch., Proc., Ser. B' Vol.71' No' I, p' I-9, ilius. 1968)
- Koenigswald, G. H. R. Von. Tektite studies-(pt.) 10, the relationship of shape, size and texture in asiatic tektites. (Konfnkl. Nederlandse Akad. Wetensch. Proc.' V. 71. NO. 1. P I-9 1968)
- Koenigswald, G. H. R. von. Tektite studies, X; The relationship of shape, size and texture in Asiatic tektites. [Ned. Akad. Wetensch., Proc., Ser. B' Vol.71' No' I, p' I-9, ilius. 1968)
- Koenigswald, G. H..R. Von, 1940. Neue Pithecanthropus -Funde 1936-1938. Ein Beitrag zu Kenntniss de Praehominiden. Dienst. Mijinb. Ned. Ind. 28 232p.
- Koenigswald, G. H..R. Von, 1959. A Mastodon and other fossils from Thailand. Rept. of the Investigation 2. Roy. Dept. Mines, Bangkok. P. 25-28, pls. 1-2, 5 and 7.
- Koenigswald, G. H..R. Von, 1962. Das Absolute Alter des Homo erectus erectus Dubois. In G., Kurth (ed.). *Evolution und Hominisation*. Gustav Fischer, Stuttgart, p. 112-119.
- Koenigswald G. H. R. v'on 1963. Tektite studies, VI. Rims, flow ridges, and flanges in Java tektites. Proc. K. Ned Akad. Wetens., -Amsterdam, B66, 198-205.
- Koesoemadinata, R. P. (1978). Sedimentary frame-work of Tertiary coal basins of Indonesia. Third regional conference on geology and mineral resources of S.E. Asia (ed. Prinya Nutalaya), pp. 62140. Asian Institute of Technology, Bangkok.
- Kolbe, P., W.H. Pinson, J.M. Saul and E. W. Miller, 1967. Rb-Sr study on country rocks of the Bosumtwi crater, Ghana, *Geochim. Cosmochim. Acta* 31, 869-875, 1967.
- Korotkova, N. N., Kashkarov, L. L., Kaskarova, V. G., Lavukhina, A. K. and Izokh, E. P., 1987. Study of tektite surface heating by track and thermoluminescence methods. 2nd Intern. Conf. on Natural Glasses, Prague. 331-337.
- Khositanont, S., C. Burrett, P. Charusiri, P. Chaodumrong, K. Howard; T. Suphajanya and S. Bunopas, 2008. Australasian Cometary Impact terrestrially from near the North Pole at 0.8 Ma. Best Dated and Global Correlation of Pre-late Lower Quaternary Alpine-Himalayan Epeirogenesis-Inthanon Uplift continuing to New Zealand-Pacific Orogenesis; also Euraustralasian Quaternary Geological Update. GREAT 2008,
- Kring, D. A., 1997, Air blast produced by the meteor crater impact event and a reconstruction of the affected environment: *Meteoritics and Planetary Science*, v.32, p. 5 17-530.
- Kring, D. A., 2003. Environmental consequences of impact cratering events as a function of ambient conditions on Earth, *Astrobiology* 3 (2003) 133-152.
- Krinov, E. L., 1966, *Giant meteorites*: London, Pergamon Press.
- Kroenke, I. W., and Rodda, P. 1984. Cenozoic tectonic development of the southwest Pacific. U.N. ESCAP, CCOP/SOPAC Technical bulletin No. 6. -
- Kunz, J., Bollinger, K., and Jessberger, E. K., 1995, Ages of Australasian tektites: Lunar and Planetary Science Conference, 266, P roceedings : p, 809- 810.
- Kudrass, H. R. and Schlüter, H. U. 1983. Report on geophysical and geological offshore surveys in the Malacca Strait (1976). Exploration for offshore placers and construction materials CCOP/TP 12, Bangkok, 115-154.
- Kukla, G., and An, Z., 1989. Loess Stratigraphy in Central China. *Palaeogeog. Palaeoclim. Palaeoecol.* 72, 203-225.
- Kumai, H. et al., 1985. Geology and stratigraphy of the Mokojeti Area. In Watanabe, N., Kadar, D., eds. 1985. *Quaternary Geology of the Hominid Fossil Bearin in Jave*. Geol. Res. Devel. Centre Sp. Paper No. 4. 378p. Bundung.
- Kunimatsu, Y., B. Ratanasthien, H. Nakaya, H. Saegusa, and S. Nagaoka. 2003. The Potential of Thailand for Understanding the Hominoid Evolution in Southeast Asia. RCNPNS - 8th Int. Congress on Pacific Neogene Stratigraphy. Chiang Mai, Febr. 2003. p. 83-89.
- Kuno, H. (1962). Catalogue of the active volcanoes of the world including solfatara fields: Japan, Taiwan and the Marianas. 11. International Association of Volcanology, Rome.
- Kuno, H. (19 66). Lateral variation of the basalt magma type across continental margins and island arcs. *Bulletin of Volcanology* 29, 195:222.
- Kunz, J., Bollinger, K., Jessberger, E.K., 1995. Ages of Australasian Tektites. Abstracts of the Lunar and Planetary Science Conference, 26th, p. 809-810.
- Ky, Hoang Ngoc, 1986. The Phuduc Loess Formation in Vietnam and loess distribution the World over. In: Abstracts of Papers, First Conference on Geology of Indochina. 5-7 Dec. 1986. Ho Chi Minh City. P. 42-43.
- Kyte, F. T., 1980 Siderophile-enriched sediments from the Cretaceous-Tertiary boundary. *Nature*, London, 288, 651-656.
- Kyte, F. T., 1981 Extraterrestrial siderophiles in abyssal sediments. *Meteoritics*, 16, 345.
- Kyte, F. T. and Wasson J. T., 1982. Geochemical constraints on the nature of large accretionary events. In *Geological Implications of Impacts at Large Asteroids and Comets on the Earth* (ed. L. T. Silver): GSA Spec. Paper 190, 235-242.
- Kyte, F. T. and Wasson J. T., 1982. Search for iridium anomalies. Abstracts 13th Lunar Planet. Sci. Conference, 412.
- Kyte, F. T. and Wasson, J. T., 1986. Accretion rate of extraterrestrial matter: Iridium deposited 33 to 67 million years ago. *Science* 232, 1225-1229.
- Kyte, F. T., Zhou, Z. and Wasson J. T., 1980. Siderophile-enriched sediments at the Cretaceous-Tertiary boundary. *Nature* 288, 651-656.
- Kyte, F. T., Zhou, Z., and Wasson J. T., 1981. High noble metal concentrations in a late Pliocenc sediment. *Nature* 292, 4 I 7-420.
- Kyte F. T., Smit, J., and Wasson. J. T., 1985 Siderophile interelement variations in the Cretaceous-Tertiary boundary sediments from Caravaca. Spain. *Earth*

- Planet. Sci. Lett. 73, 183-195.
- Kyte F. T., Zhou L., and Wasson J. T., 1988. New evidence on the size and possible effects of a late Pliocene oceanic asteroid impact. *Science* 241, 63-65.
- Kyte F. T., Zhou L., and Lowe D., 1992. Noble metal abundances in an Early Archean impact deposit. (*Geochim. Cosmochim. Acta* 56, 1365-1372.
- Kyte F. T., Leinen M., Heath R., and Zhou L. 1993. Cenozoic sedimentation history of the central north Pacific: Inferences from the elemental geochemistry of core LL44-GPC3. *Geochim. Cosmochim. Acta* 57, 1719-1740.
- Lacassin, R., U. Scharer, P. H. Leloup, N. Arnaud, P. Tapponnier, Liu Xiaohan, and Zhang Lian-Sheng, 1996. Tertiary deformation and metamorphism SE of Tibet: The folded Tiger-leap décollement of NW Yunnan, China, *Tectonics*, in press, 1996.
- Lacassin, R., Maluski, H., Leloup, P. H., Tapponnier, P., Chaiyan Hinthong, Kanchit Siribhakdi, Saengathit Chuaviroj, Adul Charoenravat, 1997. Tertiary extrusion and deformation of Indochina western margin: structural and  $^{40}\text{Ar}/^{39}\text{Ar}$  evidence from NW Thailand. *J. Geophysical Res.*
- Lacroix, A. 1934. Sur la découverte de tectites a la Côte d'Ivoire, *Comptes Rendus de l'Académie des Sciences*. 199. 1539—1542.
- Lacroix, A. 1935. Alfred Lacroix Les tectites sans formes figurées de l'Indochine. *Comptes Rendus de l'Académie des Sciences*, 200, 2129-2132.
- Lacroix, A., 1935. Les tectites de l'Indochine des ces abords et celles de la Cote d' Ivoire. *Archives du Muséum national d'histoire naturelle (Paris)*, 6th ser. 12, 151-170.
- Ladd, H. S. and R. W. Brown, 1956. Fossils Lift the Veil of Time. *The Nat. Geographic Magazine*. Vol. CIX, No. 3. March, 1956, 363-386.
- Lam, S. K., 1983. Tektite found in Sarawak. *Warta Geologi: Newsletter of the Geol. Soc. Malaysia*. 9 (6), 273-275.
- LaMarche P. H., Rauch F. And Lanford W. A. 1984. Reaction between water and tektite glass. *J. Non-Cryst. Solids* 67, 361-369
- Langford-Smith, T. 1960. The dead river systems of the Murrumbidgee. *Geogr. Rev.* 5. 368-89.
- Langford-Smith, T., and Dury, G. H. (1965). Distribution, character, and attitude of the duricrust in the northwest of New South Wales and the adjacent areas of Queensland. *Am. J. Sci.* 263, 170-90.
- Latin, O., and White, N., 1990. Generating melt during lithospheric extension: pure shear vs. simple shear. *Geology* 18, 327-331.
- Lauprasert, K., Watchajittaphan, P., Juanggam, S., & Bhuttarach, S. 2019. Freshwater crocodile, *Crocodylus siamensis* Schneider, 1801, from the Middle Pleistocene deposits in Chaloeam Phrakiat District, Nakhon Ratchasima, Thailand. *Annales de Paléontologie*. doi:10.1016/j.annpal.2019.05.002
- Le Dain, A. Y., P. Tapponnier, and P. Molnar, 1984. Active faults and tectonics of Burma and surrounding regions, *J. Geophys. Res.*, 89, 453-472, 1984.
- Leakey, L. S. B. R. Protsch and R. Berger. 1968. Age of Bed V, Oldavai Gorge, Tanzania. *Science*, v 168, p. 559.
- Lee, M.-Y., K.-Y. Wei, 2000. Australasian microtektites in the South China Sea and the West Philippine Sea: implications for the age, size, and location of the impact crater, *Meteorit. Planet. Sci.* 35 (2000) 1151-1155.
- Lee, Y.-T., J.-C. Chen, K-S Ho, and W-S. Juang (2004), Geochemical studies of tektites from east Asia, *Geochem. J.*, 38, 1 - 17.
- Lekakul, B. 1949. "The exclamation of the fossils of Hipopotamus, Stegodon insignis at Nakorn Sawan". Collected papers at the Science Conference 1948, 40-44.
- Le Loeuff, J.; Saenyamoon, T.; Souillat, C.; Suteethorn, V. and Eric Buffetaut. 2007. Triassic Trackways from Thailand. (GEOTHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. Geothai 2007, pp. 362-363. 21-22 November. Imeral Hotel, Bangkok, Thailand
- Leloup, P. H., T. M. Harrison, F. J. Ryerson, Chen Wenji, L. Qi, P. Tapponnier, and R. Lacassin, 1993. Structural, petrological and thermal evolution of a Tertiary ductile strike-slip shear zone, Diancang Shan, Yunnan. *J. Geophys. Res.*, 98, 67 15-6743, 1993.
- Leloup, P. H., R. Lacassin, P. Tapponnier, Zhong Dalai, Liu Xiaohan, Zhang Lianshang, Ji Shaocheng, and Phan Trong Trinh, 1995. The Ailao Shan-Red River shear zone (Yunnan, China), Tertiary transform boundary of Indochina, *Tectonophysics*, in press, 1995.
- Letouzey, J. and Kimura, M. 1985. The Okinawa Trough: genesis of a back-arc basin developing along a continental margin. *Tectonophysics* 125, 129-30.
- Levell, B. K. and Kasumajaya, Awang. 1985. Slumping at the Late Miocene shelf-edge offshore west Sabah: A view of a turbidite basin margin. *Geological Society Malaysia Bulletin* 18, 1-29.
- Lewis, S. D. and Hayes, D. E. 1983. The tectonics of northward propagating subduction along eastern Luzon, Philippine islands. Hayes, D. E. (ed.) *The tectonic and geologic evolution of Southeast Asian seas and islands*, part 2. Geophysical monograph 27 (ed. D. E. Hayes), pp.57-78. American Geophysical Union, Washington.
- Lewis, J. S., Wtkin, K. H., Hartman., and Prinn, R. G., 1982. Chemical consequences of major impact events on Earth. *Geol. Soc. Amer. Spec. Pap.* 190, 215-222.
- Li, H.M., An, Z.S. & Wang, J. D., 1974. Preliminary paleomagnetic study of loess from the Wucheng section, Northern China. *Geochimica* 2, 93-104. (in Chinese with English abstract)
- Li, C.; Ouyang, Z.; Liu, D. and An, Z., 1991. Microtektites and glass microspherules in Loess; Their discovery and implications (abstract). *INQUA Conference Beijing*, p. 132.
- Li, C.; Quyang, Z. 1992. The first discovery of Australasian-Asian microtektites in loess. 29th International Geological Congress. Abstract, vol. 3(3). Kyoto, Japan. Aug.-3 Sept., 1992. II-12-4 O-9, p.650.
- Li Desheng 1984. Geologic evolution of petroliferous basins on continental shelf of China. *American association of petroleum Geologists bulletin* 68, 993-1003.
- Lin Wenzhu. "Source crater parent rocks of the Australasian strewnfield tektite." 1990. (*METEORITICS* ; Vol. 25, No. 4. p. a19-a20)
- Lin, J. L., Fuller, .M., and Zhang, W. Y. 1985. Preliminary Phanerozoic polar wander paths for the north and south China blocks. *Nature* 313, 444-9.
- Liu, T.S. 1988. Loess in China. Springer-Verlag, Berlin & Heidelberg.
- Liu, X. M., Liu, T .S., Xu, T. C., Liu, C. & Chen, M.Y., 1989.

- The Chinese loess in Xifeng, I. The primary study on magnetostratigraphy of a loess profile in Xifeng area, Gansu Province. *Geophys.* 1. 92, 345-348
- Liu, T.S. & Chang T.H., 1962. The Huangtu (loess) of China. *Acta Geologica Sinica* 42, 1-14.
- Liu, T. S. & Ding, Z. L., 1993. Stepwise coupling of monsoon circulations to global ice volume variations during the late Cenozoic. *Global and Planetary Change* 7, 119-130.
- Loofs H. H. E., 1977, Tektites in Thai prehistory. *Asian Perpect.* 20, 113-129.
- Lovering J. F., Mason B., William G. E., and Mcl-oll D. H. (1972): Stratigraphical .oid..r for the terrestrial age of australites. - *Journ. of the Geol. Soc. of Ausralia*, tg, 4i9-4t8.
- Lowe, D. R., 1976, Grainflow and grainflow deposits: *Sedimentary petrology Journal*, v. 46, p. 188-199.
- Luft, E., 1982. Zur Bildung der Moldavite beim Reissimpaki aus Tertiaren Sedimenten. Ph.D. Dissert.. Eberhard - Karis -Universitat.. Tubingen. 202. pp.
- Lyne, J. E., Tauber, M., 1995. Origin of the Tunguska event. *Nature*, 375, 638-639.
- Ma, P., et al. (2004). Beryllium-10 in Australasian tektites: Location of the source crater, *Geochim. Cosmochim. Acta*, 68, 3883-3896.
- Maloney, B. K., Howard, K. T., 2000. Tha Chang, Khorat Plateau, Thailand: The stratigraphy and pollen content of sandpits 1 and 2. *Geological Society of Thailand Newsletter*, April, 2-8,.
- Mankinen, E. A.; J. M. Donnelly-Nolan. C. S. Grommé and B.C. Hearn, 1980. Paleomagnetism of the Clear Lake volcanics and new limits on the age of the Jaramillo normal-polarity event. *U.S. Geol. Surv. Prof. Pap.* 1141. 67-82. 1980.
- Maranate, S., 1982, Paleomagnetism of the Khorat Group in Northeast Thailand: Thesis submitted for the degree of M. Sc., Victoria University of Wellington, New Zealand. 398 p. (copies lodged at Victoria University Library, and at DMR., Bangkok.)
- Maranate, S., 1983, Paleomagnetic core collection and field data of the Huai Hin Lat Formation in Northeast Thailand: *GEOMAG Pub. no. 10*, Geological Survey Division, Department of Mineral Resources, 25 p
- Maranate, S., 1984. Magnetostratigraphic correlation for dating in the Khorat Group, Northeastern Thailand. *Conf. on Application in Geol. and Nat. Development*. Chulalongkorn University, Bangkok, 19- 22, Nov., 1984.
- Maranate, S., and Vella, P. 1986. Paleomagnetism of the Khorat Group, Mesozoic, Northeast Thailand. *J. of Southeast Asia Earth Sciences*, 1, 23-31.
- Martin, J.E., Buffetaut, E., Naksri, W., Lauprasert, K., and Claude, J. 2012. Gavialis from the Pleistocene of Thailand and its relevance for drainage connections from India to Java. *PLoS ONE*, 7(9):e44541. <https://doi.org/10.1371/journal.pone.0044541>
- Masaytis, V. L.. el al. 1989 "Zapadno-Sibirskiy areal rasseyaniya tektitopodobnykh stekol." *Doklady Akademii Nauk Sssr*, Vol. 304, No. 6, p. 1419-1423
- Matsuda, J. -I., Matsubara, K. and Koeberl C., 1993. Origin of tektites: Constraints from heavy noble gas concentrations. *Meteoritics* 228, 586-599.
- Matsu'ura, S. 1985. A consideration of the stratigraphic horizons of hominid finds from Sangiran by the fluorine method. In (N. Watanabe & D. Kadar Eds.) *Quaternary Geology of the Hominid Fossil Bearing Formations in Java: Report of the Indonesia-Japan Joint Research Project CTA-41, 1976-1979. Special Bulletin Number. 4.* Bandung, Indonesia: Geological Research and Development Centre, pp. 359-366.
- McCaffrey, R. 1989. Seismological constraints and speculations on Banda Arc tectonics. *Netherlands Journal of Sea Research* 24, 13 I 52.
- McGetchin T. R., Settle M., and Head J. W. 1973 Radial thickness variation in impact crater ejecta: Implications for lunar basin deposits. *Earth Planet Sci. Lett.* 20, 226-236.
- McHone, J. F., Jr. and Greeley, R., 1987. Talemzane: Algelian impact crater detected on SIR-A orbital imaging radar. *Meteoritics* 22, 253-279.
- McHone, J. F., Pitakpaivan, K., Bunopas, S., Angsuwatana, P., Supajanya, T. and Wasson, J. T. 1994: A search for tektite related structure: An examination on SPOT satellite images. *New Development Regarding KT Event and Other Catastrophe in Earth History*, 80-81. Lunar and Planetary Institute (LPI) Contribution No. 825, p. 80-81.
- McKenzie, OP., and Bickle, M.J., 1988. The volume and composition of melt generated by extension of the lithosphere. *J. Petrol.* 26, 625-679
- McLaren, D. J. 1985a. Mass extinction an Iridium anomaly in the Upper Devonian of Australia. *A commentary Geology*, 13, 170-172.
- McLaren, D. J. 1970. Time, life and boundary. *J. Paleont.* 44 801 815
- McLaren, D. J. 1985b. Paleontology; ammonoids and extinction. *Nature*, London 313, 12-13.
- McLaren, D. J. 1985c. Rare event in geology. *Epsodes*, 8
- McLennan, S. M., 1989. Rare earth elements in sedimentary rocks: influence of provenance and sedimentary processes. *Rev. Mineral.* 21, 169-200.
- Melosh, H. J., 1988. *Impact cratering: A Geologic Process.* 245 pp., Oxford University Press. New York. N. Y..
- Melosh, H. J., 2000. Can impacts induce volcanic eruptions? *Lunar and Planetary Science Institute Contribution no. 1053*, 141-142.
- Melosh, H. J., Schneider, N. M., Zahnle, K. J., and Lathan, D., 1990, Ignition of global wildfires at the Cretaceous/Tertiary boundary: *Nature*, v. 343, p. 251-254.
- Metcalfe, I., 1988, Origin and assembly of south-east Asian continental terranes, in Audley-Charles, M., and Hallam, A., eds., *Gondwana and Tethys: Geological Society of London Special Publication* 37, p. 101-118.
- Meyer, B., Mécanismes des grands tremblements de terre et du raccourcissement crustal oblique au bord Nord-Est du Tibet, Thesis, 129 pp., Doctorat de l'Université Paris 6, Paris, 1991.
- Michaux, B. 1989. Generalized tracks and geology. *Systematic Zoology* 38, 390-8.
- Miller, K.G., Berggren, W. A., Zhang, J., and Palmer-Julson, A. A., 1991, Biostratigraphy and isotope stratigraphy of upper Eocene microtektites at Site 612: How many impacts?: *Palaeonts*, v. 6, p. 17-38.
- Mitchell, A. H. G., Hernandez, F., and Dela Cruz, A. P. (1986). Cenozoic evolution of the Philippine Archipelago. *Journal of Southeast Asian Earth Sciences* 7, 3-22.
- Molnar, P. and Tapponier, P. 1975. Cenozoic tectonics of Asia: effects of a continental collision. *Science* 189,419-26'
- Moore, G. F. and Karig' D' E. 1980. Structural geology of Nias island, Indonesia: implications for subduction and tectonics' *American Journal of Science* 280,793-

- 223.
- Moore, G. F. and Silver, E. A. 1983. Collision processes in the northern Molucca Sea. The tectonic and geologic evolution of Southeast Asian seas and islands, part 2, Geophysical monograph 27 (ed' D. E. Hayes), pp. 360-72. American Geophysical Union, Washington.
- Moore, G. F., Curray, J. R., Moore, D' G., and Karig, D. E. (1980). Variations in geologic structure along the Sunda fore-arc, northeastern Indian Ocean. The tectonic and geologic evolution of Southeast Asian seas and islands, Geophysical monograph 23 (ed. D. E. Hayes), pp. 45-60. Geophysical Union, Washington.
- Molnar, P., and England, P.C., 1990. Late Cenozoic uplift of mountain ranges and global climate change - chicken or egg? *Nature* 346, 29-34.
- Molnar, P., England, P.C., and Martinod, M., 1993. Mantle dynamics, the uplift of the Tibetan Plateau, and the Indian monsoon. *Rev. Geophys.* 31, 357-396.
- Montanari, A., C. Koeberl, 2000. Impact stratigraphy: the Italian Record, *Lect. Notes Earth Sci.* vol. 93, Springer, Berlin, 2000, 364 pp.
- Morris, J. D., Jezek, P. A., Hart, S. R., and Gill, J. B., 1983. The Halmahera island arc, Molucca Sea collision zone, Indonesia: a geochemical survey. The tectonic and geologic evolution of Southeast Asian seas and islands, part 2. Geophysical monograph 27 (ed. D. E. Hayes), pp. 373-87. American Geophysical Union, Washington.
- Muller, O., and W. Gentner, 1965. Gas content in bubbles of tektites and other natural glasses, *Earth Planet. Sci. Lett.*, 4, 406-410, 1965.
- Muller, O. and Genter, W., 1973. Enrichment of volatile elements in Muong Nong type tektites: clue for their formation history (abstr.). *Meteorites* 8, 414-415
- Muller, R. A. and Morris, D.. 1986. Geomagnetic reversals driven by abrupt sea level changes. *Geophys. Res. Lett.* 13: 1177-1180.
- Muller, R. A. and D.E. Morris, 1989. Geomagnetic reversals driven by sudden climate changes, *Trans. Am. Geophys. Union* 70, 276, 1989.
- Murali, A. V., et al. "Tektite-like bodies at Lonar Crater, India: implications for the origin of tektites." *Ryder, Graham, et al. March 30, 1987. (JOURNAL OF GEOPHYSICAL RESEARCH, B, SOLID EARTH AND PLANETS ; Vol. 92, No. 4, p. E729-E735)*
- Murali, A. V., et al. "Impactite and tektite glasses from Lonar Crater, India." *Nov. 12, 1985. (EOS, TRANSACTIONS, AMERICAN GEOPHYSICAL UNION ; Vol. 66, No. 46, p.948 )*
- Murali, A.V., et al. "impactite and tektite glasses from Lonar Crater, India." *May 1986. (JOURNAL OF THE, GEOLOGICAL SOCIETY OF INDIA. Vol.27, No. 5, p.473)*
- Muraoka, J., Poom-Im, S., Dinsohita, K., 1997. The Tertiary deposits- Phae Formation- in the Phrae Basin. *Proc. of the national conf. on stratigraphy and tectonics evolution of Southeast Asia and the South Pacific.* 19-24 Augt. 1997, Bangkok, Thailand, 587-595.
- Myint, T., D. Bannert, D. D. Helmcke, and S. Ritzkowski. 2003. On the geology of Ramree and Cheduba islands, Arakan Coast, Union of Myanmar. *RCNPS 46 - 8th International Congress on Pacific Neogene Stratigraphy.* Chiang Mai, Thailand, pp 1-9. - February 2003.
- Nagaoka, S., Y Suganuma, and B. Ratanasthien, 2003. Tectonics and paleomagnetism of Tertiary strata with mammalian fossils in northern Thailand. *RCNPS - 8th Int. Congress on Pacific Neogene Stratigraphy.* Chiang Mai, Febr. 2003. p. 90. P. 285-285.
- Naksri, W., Tong, H., Lauprasert, K., Jintasakul, P., Suteethorn, V., Vidthayanon, C., & Claude, J. 2019. Plio-Pleistocene giant tortoises from Tha Chang sandpits, Nakhon Ratchasima Province, Thailand. *Annales de Paléontologie.* doi:10.1016/j.annpal.2019.06.001
- National Geographic (Thai Edition), 2006. Attached National Geographic Map on Earthquakes Rings of Fires, and Stunami in South Thailand. March 2006, Bangkok.
- Nayaka, H., H. Saekusa, B. Ratanasathien, Y Kunimatsu, T Tsubamoto, S. Nagaoka, Y Suganuma, P Chintaskul, Y Thasod. 2003. Late Cenozoic Mammalian Faunas of Thailand. *RCNPS - 8th Int. Congress on Pacific Neogene Stratigraphy.* Chiang Mai, Febr. 2003. p. 90.
- Nesbitt, H. w., and Young, G.M., 1982, Early Proterozoic climates and plate motions inferred from major element chemistry of lutites : *Nature*, v. 299, p. 7 | 5 -7 17 .
- Neumann van Padang, M. 1951. Catalogue of the active volcanoes of the world including solfatar fields, part I, Indonesia. *International Volcanology Association, Naples.*
- Neumann van Padang, M. 1953. Catalogue of the active volcanoes of the world including solfatar fields, part II, Philippine islands and Cochin China. *International Volcanological Association, Naples.*
- Nguyen Thi Kim Thoa. 1995. Investigations on some Paleozoic sediments from North Vietnam. *Journal of Geology, special issue, Series B, no. 5-6 (1995), Pp. 380-388.*
- Nguyen Thanh Giang 1982. Palaeomagnetic studies of Cenozoic basalts in Vietnam. *Palaeomagnetic Research in Southeast and East Asia, CCOP/TP 13'58-63.*
- Nicolas, A., et al. 1981. The Xigaze ophiolite (Tibet): a peculiar oceanic lithosphere. *Nature* 294, 414-17.
- Ninkovich, D. 1975. Late Cenozoic clockwise rotation of Sumatra. *Earth and Planetary science letters* 29, 269-75 .
- Nicolaysen, L.O.. 1987. Tektites: ejecta from massive cratering events, caused by periodic escape and detonation of deepmantle fluids. In: *mt. Workshop Cryptoexplosions and Catastrophes in the Geological Record (Pars). Sect. N3. 15 pp.*
- Nicholes, D. J., Jarzen, D. M., Orth, C. J. and Oliver, P. Q., Palynological and Iridium anomalies at South-Central Cretaceous - Tertiary boundary Saskatchewan. *Science*, 231, 714-717.
- Nie, S., Rowley, D. B. and Ziegler, A. M. 1990. Constraints on the locations of Asian microcontinents in Paleotethys during the Late Paleozoic In 'Palaeozoic Palaeogeography and Biogeography, Geological Society Memoir No. 12'. (Eds W. S. McKerrow. and C. R. Scotese.) 397-409. (Geological Society of London.)
- Niitusuma, N. 1977. Zone magnetization model and depth lag of NRM in deep-sea sediments, *Rock Magn. Paleogeophys.* 4. 65 -71. 1977.
- Nilson, T., 1983. Pleistocene Geology and life in the Quaternary Ice Age. 651 pp., Stuttgart.
- Nininger H. H. & Huss G. I. 1967. Tektites that were partially plastic after completion of surface sculpturing. *Science* 157, 6 1-62.

- Ninkovich, D. 1976. Late Cenozoic clockwise rotation of Sumatra. *Earth and Planetary science letters* 29, 269-75.
- Ninkovich, D., Shackelton, N.J., Abel-Monem, A. A. Obradovich, J. D., and Izett, G. 1978. K:Ar age of the Late Pleistocene eruption of Toba, north Sumatra. *Nature* 276, 574-77.
- Nishimura, S. and Suparka, S. 1985. Tectonic development of east Indonesia. *Journal of Southeast Asian Earth Sciences* 1,45-57.
- Nishimura, S., Abe, E., Yokoyama, T., Sugiyarta, and Dharma, A. 1978a. Danau Toba-the outline of Lake Toba, north Srrmatara, Indonesia. *Studies of physical geology on the Sunda island arc* (ed. S. Sasajima), pp. 54-62. University Press, Kyoto.
- Nishimura, S., Sasajima, S., Hirooka, K., Thio, K. H., and Hehuwat, F. 1,978b. Radiometric ages of volcanic products in Sunda arc. *Studies of physical geology on the Sunda island arc* (ed. S. Sasajima), pp. 34-7. University Press, Kyoto.
- Nishimura, S., et al. 1981. Physical geology of the Sumba, Sumbawa and Flores islands. *The geology and tectonics of eastern Indonesia, Geological research and development centre, Bandung, Special Publication 2* (ed. A. J. Barber and S. I Viryosujono), pp. 105-113.
- Nur, A. and Ben Avraham, Zvi. 1977. Lost Pacifica continent. *Nature* 270, 41-3.
- Nutalaya, Prinya (ed.) 1978. *Geology and mineral resources of Southeast Asia: GEOSEA III*. Asian Institute of Technology, Bangkok.
- Nutalaya, P. (ed.) 1983. *Stratigraphic correlation of Thailand and Malaysia 1, technical papers*. Geological Society of Thailand, Bangkok/Geological Society of Malaysia, Kuala Lumpur.
- Nutalaya, P., Petchvarun, V., and Saengsuwan, C. 1978. *Bibliography of geology and mineral resources of Thailand (Thai language)*. Prinya Nutalaya (ed.), *Geology and mineral resources of Southeast Asia* (ed. Prinya Nutalaya), pp. 2343. Asian Institute of Technology, Bangkok.
- Nutalaya, P. and Rau, J. L. 1981. Bangkok: the sinking metropolis. *Episodes* 4, 3-8.
- Nutalaya, P., P. Vella, S. Bunopas and V. Kaewyana, 1986. Quaternary processes in Thailand. In: Thiramongkol (ed.), *Proceeding of the Workshop on Economic, Geology, Tectonics Sedimentary Processes and Environmental of the Quaternary of Southeast Asia*. Pp. 23-33, Bangkok.
- Oba, N., Tomita, K., Yamamoto, M., Istidjab, M., Sudradjat, A., and Suhandi, T. 1985. Geological significance of granitic fragments found in pumice flow 1883 eruption at the Krakatau Group, Indonesia. *Geological society of Malaysia bulletin* 19,51-58.
- Officer, C. B., and Drake, C. L., *Terminal Cretaceous Environmental Events*. *Science*. 227, 1161-1167.
- Officer, C. B., Hallam, A., Drake, C. L. Devine, J. D., 1987. Late Cretaceous and Paroxysmal Cretaceous / Tertiary extinctions. *Nature*, London, 326, 143-149.
- O'Keefe, J. A., P. D. Lowman, and K. L. Dunning, 1962. Gas in tektite bubbles, *Science*, 137, 228.
- O'Keefe, J. A. 1967. Tektite sculpturing. *Geochimica et Cosmochimica Acta* 31, 193 1-1933.
- O'Keefe, J. A., 1969, The microtektite data: Implications for the hypothesis of the lunar origin of tektites: *Journal of Geophysical Research*, v. 74, p.6795-6804.
- O'Keefe, J. A., 1976. *Tektites and Their Origin*. 354 pp., Elsevier. Amsterdam.
- O'Keefe, J. A., 1984. *Natural Gases.* ; In; L. D. Pye, J. A. O'Keefe and V. D. Frechette (eds.), *Natural Glasses*. North Holland Phys. Publ. Amsterdam, pp. 1-17.
- O'Keefe, J. A., 1987. Zhamanshin and Auelloul: craters produced by impact of tektite-like glasses? *Meteorites*, 22, 219-228.
- O'Keefe, J. A., and Allen, T. J., 1982. The interaction of the Cretaceous/Tertiary extinction bolide with the atmosphere, ocean, and solid earth. *Geol. Soc. Am. Spec. Pap.*, 190. 103-120.
- Olson, E. C., 1982. Extinction of Permian and Triassic nonmarine vertebrate. *Geol. Soc. Amer.*, Special paper. 190, 501- 511.
- Omez, I., Finnegan, D. L., and Zoller, W. H., 1986. Iridium emissions from Kilauea Volcano. *J. Geophys. Res.*, 91, 653.
- Orth, C. J., Gilmore, J. S., Knight, J. D., Pilmore, C. L., Tschudy and Fassett, J. E. 1981. An Iridium abundance anomaly at the palynological Cretaceous -Tertiary boundary in Northern New Mexico. *Science*, 214, 1341-1343.
- O'Sullivan, T., Pegum, D., and Tarigan, J. 1985. *Seram oil search, past discoveries and future oil potential*. *Proceedings of the Indonesian Petroleum Association 74th annual convention, Jakarta I*, 3-20.
- Otofuji, Y., Sasajima, S., Nishimura, S., Dharma, A., and Hehuwat, F. (1982). paleomagnetic evidence for clockwise rotation of the northern arm of Sulawesi, Indorresia. (abstract). C.C.O.p. paleomagnetism workshop, Kuala Lumpur, March 1982.
- Ouyang, Z., Zhong, P., Yi, W., 1976. Trace elements in tektite from Hainan Island, China. *Geochimica, Peking* no. 2, 144-147
- Ozima, M., Kaneoka, I. and Ujiej, H. 1977. <sup>40</sup>Ar-<sup>39</sup>Ar age of rocks, and the development mode of the Philippine Sea. *Nature* 267, 816-18.
- Parker, E. S. and Gealey, Jf. K. (1985). Plate tectonic evolution of the western Pacific-Indian Ocean region. *Energy* 1A, 249-51.
- Parsons, B. and Sclater, J. G. (1977). An analysis of the variation of ocean floor bathymetry and heat flow with age. *Journal of Geophysical Research* 82, 803-27.
- Peccherillo, A. and Taylor, S. R. (1975). Geochemistry of Eocene calc-alkaline volcanic rocks from the Kastamonu area, northern Turkey. *Contributions to mineralogy and petrology* 58, 53-81.
- Packham, G., 1996, *Cenozoic Southeast Asia: reconstructing its aggregation and reorganization*,
- Page, B. G. N., et al. 1979. A review of the main structural and magmatic features of northern Sumatra. *Journal of the geological society London* 135,569-79.
- Page, B. N. and Suppe, J. 1981. The Pliocene Lichi mélange of Taiwan: its plate-tectonic anolistrostomal origin. *American Journal of Science* 281,193-227.
- Pal, D.K., Tuniz, C., Moniot, R.K., Druse, T.H., and Herzog, G.F. 1982. Beryllium-10 in Australasian tektites: evidence for a sedimentary precursor. *Science*, 218: 787-789.
- Parry, J. T., 1996: The high terrace gravels, Northeast Thailand- a revaluation and integrated theory of their origin. *Z. Geomorph. N. I.* 42 (2): 145-175.
- Parsons, B. and Sclater, I. G. 1977. An analysis of the variation of ocean floor bathymetry and heat flow with age. *Journal of Geophysical Research* 82, 803-27.
- Paul, D. D. and Lian, H. M. 1975. *Offshore Tertiary-basins*

- of Southeast Asia, Bay of Bengal to Siam China Sea. Ninth World petroleum congress proceedings 3, Exploration and transportation. w. 107-21. Applied Science publishers, London.
- Peccerillo, A. and Taylor, S. R. 1975. Geochemistry of Eocene calc-alkaline volcanic rocks from the Kastamonu area, northern Turkey. Contributions to mineralogy and petrology 58, 53-81.
- Petitot, A. 1895. Geologie de L'Indochine. Imprimerie Nationale, Paris
- Peltzer, G., and P. Tapponnier, 1988. Formation and evolution of strike-slip faults, rifts, and basins during India-Asia collision an experimental approach, J. Geophys. Res., 93, 15,085-15,117, 1988.
- Peng, H., K. Zhao, and S. Chen. 1982. The evidence of 'CHOI' in sediment core samples of Pacific Ocean collected among FGGE, J. Geoph.t:c. Res., 87, 5563-5565.
- Piccoli, G., Robba, E., Chonglakmani, C., Dheeradilok, P., Ingavat, R., 1984. Paleogene Tethyan and Neogene Indo-Pacific shallow benthic molluscs: some thoughts. Jour. Geol. Soc. Thailand. v. 7 (1), 13-18.
- Pierce, K. L., 1986. Dating Methods. In: Active Tectonics. p. 195-214. National Academy Press, U.S.A.
- Pieters, P. E., and Supriatna, S. 1990. Late Cretaceous-Early Tertiary continent collision in Borneo. In: Terrane Analysis of China and the Pacific Rim'. (Eds. T. J. Wiley, D. O. Howell and F. L. Wong.) 193-4. (Circum-Pacific Council for Energy and Mineral Resources: Houston, Texas.)
- Pigram, C. J., and Davies, H. I., 1987. Terraines and the accretion history of the New Guinea orogen. BMR Journal of Australian Geology and Geophysics 10. 193-211.
- Pigram, C. J. and Panggahean, H. 1984. Rifting of the northern margin of the Australian continent and the origin of some Microcontinents in eastern Indonesia. Tectonophysics 107, 331-53.
- Pigram, C.J., Challinor, A. B., Hasibuan, F., Rusman E., and Hartono, U. 1982a. Lithostratigraphy of the Misool Archipelago, Irian Jaya, Indonesia. Geologie en mijnbouw 61,265-79.
- Pigram, C. J., Robinson, G. P., and Tobing, S. L. 1982b. Late Cainozoic origin for the Bintuni Basin and adjacent Lengguru fold belt, Irian Jaya. Proceedings of the Indonesian Petroleum Association 77th annual convention, Jakarta, 109-26.
- Pimm, A. C. 1955. Serian area, west Sarawak Malaysia. Geological survey of the Borneo Region Malaysia report 3.
- Pimm, A. C. 1967. Bau mining district, west Sarawak, East Malaysia. Part II: Kronkong. Geological survey of the Borneo Region Malaysia bulletin 7.
- Philippines, 1981. Proceedings, fourth regional conference on the geology of Southeast Asia. Geological society of the Philippines, Manila.
- Pike R. J. 1974. Craters on Earth, Moon and Mars. Multivariate classification and mode of origin. Earth and Planetary Science Letters 22, 245-255.
- Pohl, J., 1977. Palaeomagnetische und gesteinsmagnetische Untersuchungen an den Kernen der Forschungshohrung Noerdlingen 1973, Geol. Bas. 75, 329-348. 1977.
- Pohl, J. 1978. Evidence for the coincidence of a geomagnetic reversal with the Ries impact event. Meteoritics. 13: 600 (Abstr.).
- Polachan, S., S. Praditjan, C. Tongtaow, S. Janmaha, K. Intarawijitr, C. Sungsuwan. 1991. Development of Cenozoic basins in Thailand. Mar. Petrol. Geol. 8: 84-97
- Poreda, R., and Craig, H., 1989. Helium isotope ratios in circum-Pacific volcanic arcs. Nature 338, 473-475.
- Pramojanee, P., Liengsakul, M. and Engkagul, V. 1985: Grain size analysis of some sand rises and stream sediments in the Northeast of Thailand in order to indicate depositional environment. Conf. on Geol. and Min. Res. Devel. of the Northeast, Thailand. Khon Kaen University. 26-29 Nov. 1985.
- Prakash, U., 1979. Fossil dicotyledonous woods from the Tertiary of Thailand. The Palaeobotanist, 26, 50-62.
- Prasad, M. S. 1994a. New occurrences of Australasian microtektites in the central Indian basin, Meteoritics. 29, 66-69.
- Prasad, M. S. 1994. Australasian microtektites in a substrate: A new constraint on ferromanganese crust accumulation rates, Ma.: Geol., 116, 259-266.
- Prasad, M. S., and V. D. Khedekar. 2003. Impact microcrater morphology on Australasian microtektites, Meteoritics. Planet. Sci., 38, 1351 - 1357.
- Prasad, M. and P. S. Rao 1990. Tektites far and wide. Nature. 347, 340. Prasad, M. S., and M. Sudhakar. 1996a. Impact microcraters on an Australasian microtektite. Meteorit. Planet. Sci., 31, 46-49.
- Prasad, M. S., and M. Sudhakar. 1996b. Australasian microtektites from the central Indian basin: Implications for ejecta distribution patterns, Curr Sci., .70, 74-77.
- Prasad, M. S., and M. Sudhakar. 1998. Microimpact phenomena on Australasian microtektites: Implications for ejecta plume characteristics and lunar surface processes, Meteorit. Planet. Sci., 33, 1271-1279.
- Prasad, M. S., and M. Sudhakar. 1999. Australasian microtektites discovered in the Indian Ocean. Meteorit. Planet. Sci., 34, 179-184.
- Prasad, M. S., P. Ci. Mislankar, S. N. Charan, and K. Prasad. 1998. An Australasian tektite fragment discovered in the central Indian Ocean, Curr. Sri., 74, 405-408.
- Prasad, M. S., S. M. Gupta, and V. N. Kodagati. 2003. Two layers of Australasian impact ejecta in the Indian Ocean?, Meteorit. Planet. Sci., 38, 1373-1381.
- Prasad, M.S., Mahale, V.P., and Kodagali, V.N., 2007. New sites of Australasian microtektites in the central Indian Ocean: Implications for the location and size of source crater: Journal of Geophysical Research, v. 112. p. E06007, doi: 10.1029/2006J8002857.
- Priem, H. N. A., et al, (1978). Isotopic evidence for a Middle to Late Pliocene age of the cordierite granite on Ambon, Indonesia. Geologie en mijnbouw 57, 441-3.
- Priem, H. N. A., Boelrijk, N. A. I. M., Bon, E. H., Hebeda, E. H., Verdurman, E. A. T., and Verschure, R. H. (1975). Isotope geochronology in the Indonesian tin belt. Geologie en mijnbouw 54, 61-70.
- Prothero, D. R., and Schwab, F., 1996, Sedimentary Geology: New York, W. H. Freeman.
- Punjasutharos, S., 1995. Artesian Flowing Wells in Unconsolidated Rock Aquifer along the Mun River. International Conference on Geology, Geotechnology and Mineral Resources of Indochina, Royal Thai Department of Mineral Resources, Bangkok, 521-528.
- Punpate, N.; S. Pailoplee; I. Takashima and P. Charusiri, 2005. Ages of layered Tektites and tektites bearing sediments in Buntharik area, Ubonrathchathani, Northeast Thailand. Proc. of the Int. Conference on Geology, Geotechnology and Mineral Resources of

- Indochina. GEOINDO 2005, Kosa Hotel, Khon Kaen, 28-30 Nov. 2005.
- Qian Ding-yu, Zhang Shu-min & Gu Qing-ge. 1982. Some new data on the age of the Namring Formation of Xigaze group in the Southern Xizang (Tibet), China.
- Qui, Y., Xia, Y., Yang, S. 1993. The contorted sutured belt of East Tethys. Gondwana Dispersal and Asian Accretion. IGCP 321 Third Symposium and Field Excursion, 23-28 August 1993. Kuala Lumpur, Malaysia.
- Raeside, J. D., 1964. Loess deposits of the South Island, New Zealand, and soil formed on them. *N. Z. J. Geol. Geophys.* 7, 811-88.
- Raisbeck, G.M., F. Yiou, D. Bourles and D.V. Kent, Evidence for an increase in cosmogenic  $^{10}\text{Be}$  during a geomagnetic reversal, *Nature* 315, 315-317. 1985.
- Raj, J. K., 1983. Negative lineaments in the granitic bedrock areas of N. W. Peninsular Malaysia. *Geological society of Malaysia bulletin* 16, 61-70.
- Raisbeck, G.M., Yiou, F., Bourles, D., and Kent, D.V., 1985. 'Evidence for an increase in cosmogenic  $^{10}\text{Be}$  during a geomagnetic reversal', *Nature*, 315, 315-317.
- Rampino, M. R., and Stothers, R. B., 1984. Terrestrial mass extinctions, cometary impact and the Sun's motion perpendicular to the galactic plane. *Nature*, London, 308, 709-712.
- Rangin, C. et al., 1990. Paleogeodynamic maps of the West Pacific. *Bull. Soc. Géol. France*, Paris, 1990, no 6.
- Rangin, C., Jolivet, L., Pubellier, M. and the Tethys working group. 1990. A simple model for the tectonic evolution of Southeast Asia and Indonesia region for the past 43 Myr. *Bull. Soc. Geol. France* (8) VI, no. 6, 889-905.
- Rankama, K. 1965. *Nature* 207, 1383.
- Rau, J., and Nutalaya, P. 1983: Geology of the Bangkok Clay. *Bull. Geol. Soc. Malaysia*, No 16, pp 99-116.
- Raup, D. M. and Sepkoski, J. J. Jr., 1984. Periodicity in extinction in the geological past, *Proc. Natl. Acad. Sci., USA*, 81, 301-305.
- Raup, D. M. & Sepkoski, J. J. Jr., 1986. Period extinction of families and genera. *Science*, 231, 833-836.
- Remus, D., M. Webster and K. Kaewkan, 1993. Rift architecture and sedimentology of the Phechabun Intermontane Basin, central Thailand. *J. Southeast Asian Earth Sci.*, 8, 421-432.
- Reeran, A. and Koeberl, C. 1996. Water in tektites and impact glasses by courier-transformed infrared spectrometry. *Meteorit. Planet. Sci.* 32, 211-216.
- Reynolds, J. H., 1960. Rare gases in tektites. *Geochemica et Cosmochemica Acta* 20, 101-114.
- Rhodes, B. P., Conejo, R., Benjawan, T., Titas, S., Lawson, R. 2003. Paleocurrent and Provenance of the Mae Rim Formation, Northern Thailand: Implication for Tectonic Evolution of the Chiang Mai Basin. RCPNS Proceedings, 8th International Congress on Pacific Neogene Stratigraphy, Chang Mai, Thailand, 2003, pp. 311-331.
- Richter, B., Fuller, M. Schmidtke, E., Myint, U. T., Ngwe, U. T., Win, U. M., and Bunopas, S., 1993. Paleomagnetic results from Thailand and Myanmar: Implications for the interpretation of tectonic rotations in southeast Asia *Journal of Southeast Asian Earth Sciences*, 8, 247-255.
- Richtofen, F. von, 1882. On the mode of origin of the loess. *Geol. Mag.* 9, ser 2, 293-305.
- Ridd, M. F. 1971. South-East Asia as a part of Gondwanaland. *Nature* 234, 531-3.
- Ridd, M. F. 1978. Thailand. The Phanerozoic geology of the world, II, The Mesozoic (ed. M. Moullande and A. E. M. Nairn), pp. 145-53. Elsevier, Amsterdam.
- Ridd, M. F. 1980. Possible Palaeozoic drift of SE Asia and Triassic collision with China. *Journal of the geological society London* 137, 53-54.
- Rigby, J. F. and Shah, S. C. 1981. The flora from the Permian nonmarine sequences in India and Australia: a comparison. *Gondwana Flize* (ed. M. tU7. Crosswell and P. Vella), pp. 394-41. A. A. Balkema, Rotterdam.
- Rison, W., and Craig, H., 1983. Helium isotopes and mantle volatiles in Loihi Seamount and Hawaiian Island basalts and xenoliths. *Earth Planet. Sci. Lett.* 66, 407-426.
- Robbins, J.A., 1986. 'A model for particle-selective transport of tracers in sediments with conveyor belt deposit feeders', *J. Geophys. Res.*, 91, 8542-8558.
- Roberts, T. R. and J. Jumnonthai. 1999. Miocene fishes from lake, Phetchabun in North Central Thailand, with description of new taxa of Cyprinidae, Pangasiidae and Chandidae. *Nat. Hist. Bull. Siam Soc.* 47: 153-189.
- Rock, N. M. S., Syah, H. H., Davis, A. E., Hutchison, D., Styles, M. T., and Rahayu, Lena. 1982. Permian to Recent volcanism in northern Sumatra, Indonesia: a preliminary study of its distribution, chemistry and peculiarities. *Bulletin of Volcanology* 45, 127-52.
- Roger, F. S. Calassou, J. Lancelot, J. Malavieille, M. Mattauer, Xu Zhiqin, Hao Ziwen, and Hou Liwei, Miocene emplacement and deformation of the Konga Shan granite (Xiashui He Fault zone, west Sichuan, China): Geodynamical implications., *Earth Planet. Sci. Lett.*, in press, 1995.
- Rose, R. 1983. Miocene carbonate rocks of Sibolga basin, Northwest Sumatra. *Proceedings of the Indonesian Petroleum Association 72th annual convention*, Jakarta, 107-25.
- Rose, R. and Hartono, P. 1978. Geological evolution of the Tertiary Kutei-Melawi basin, Kalimantan, Indonesia. *Proceedings of the seventh annual convention of the Indonesian petroleum association*, Jakarta, 225-37.
- Ruddiman, W.F. 1971. Pleistocene sedimentation in the equatorial Atlantic: stratigraphy and faunal paleoclimatology. *Geol. Soc. Am. Bull.* 82, 283-302, 1971.
- Ruddiman, W.F. and Clover, L.K., 1972. 'Vertical mixing of ice-rafted debris', 1965. *Quaternary paleopedology*. in: *The Quaternary of the United States* (eds. Wright Jr., H.E. & Frey, D.G.), 755-764, Princeton University Press, Princeton.
- Russel, D. A. 1979. The enigma of the extinction of the dinosaurs. *Ann. Rev. Earth Planet Sci.*, 7, 163-182.
- Russel, D. A. and Rice, G. (eds.), 1982. Cretaceous-Tertiary extinction and possible terrestrial and extraterrestrial cause. *Sylogues*, 1-51.
- Sanfilippo, A., Riedel, W. R., Glass, B. P. and Kyte, F. T., 1985. Late Miocene microtektites and radiolarian extinction on Barbados. *Nature* 314, 613-615
- Sangsuwan, Ch. et al. 1987: A palynological study of the Bangkok Clay at Senavate Pit, Bangkok Metropolis. In Wezel, F.W., Rau, J. L. (eds.) *Progress in Quaternary geology of East and Southeast Asia*, pp 197-205, CCOP, Technical Publication No. 18, Bangkok.
- Santosh, M., Biju-Sekhar, S., and K. P. Shabeeer. Special Issue on Rodinia, Gondwana and Asia; *International Geoscience Journal, Gondwana Research*. Editors. 2001. *Gondwana Research* vol.4, no. 4. October 2001.

- Salyapongse, S., Pitakpaivan, K., 1999. Older eolian evidences on the Khorat Plateau. *Journal of the Geological Society of Thailand*, 1, 18-26.
- Sasajima, S., Wang, Y.Y. (eds.), 1984: Recent research on loess in China pp 201 -232, Kyoto University.
- Sasajima, S., Otofujii, Y., Hirooka, Supraka, Suwijanto, and Hehuwat, F. 1978. Paleomagnetic studies on Sumatra island on the possibility of Sumatra being part of Gondwanaland. *Rock magnetism and paleogeophysics* 5,104-10.
- Sasajima, S., Nishida, J., Maenaka, K., Chak, L. S., and Yokoyama, T. (in press). Paleomagnetic research of Jurassic acid igneous rocks and Cretaceous red bed in Hong Kong. C.C.O.P. Palaeomagnetism Conference, Kuala Lumpur, March 1982.
- Sato, Y.. 2002. Preliminary report on the occurrence of fossil mammals in Nakhon Ratchasirna, northeast Thailand, in: N. Mantajit (Ed.). *Proceedings of the Symposium on Geology of Thailand*, Department of Mineral Resources. Bangkok, 2002, pp. 230-232.
- Sattayarak, N., 1985. Review on geology of the Khorat plateau (in Thai). *Proc. Conf. In Geology and Mineral Resources development of the northeast -Thailand*. Khon Kaen. Royal Thai Department of Mineral Resources, Bangkok, 23-30.
- Saul, J.M. 1969. Field investigations at Lake Bosumtwi (Ghana) and in the Ivory Coast Tektite Strewn Field, *Nat. Geogr. Soc. Res. Rep.* 1964 Proj. 201-212, 1969.
- Saunders, J.A., M.A. Pritchett, R.B. Cook, 1997. Geochemistry of biogenic pyrite and ferromanganese coatings from a small watershed: a bacterial connection? *Geomicrobiol. J.* 4 (1997) 203 -217.
- Saurin, E., 1935. Sur quelques gisement de tektites de l'Indochine du Sud. *C. r. Acad. Sci. Paris* 1935, v. 200, p. 246-248
- Saykawlad, S, P. Sithiyos and S. Rodjanapo. 2003. Evolution of the Yom River and land use change in Srisatchanala, Sukhothai Province, Northern Thailand. *Proceedings, 8th International Congress on Pacific Neogene Stratigraphy*, Chang Mai, Thailand, 2003, pp. 203-211.
- Savage, R. J. G. and Long, M. R. 1986. *Mammal Evolution an illustrated guide*. British Museum (Natural History), 259 pp.
- Sawata, 1973. Note on a young deposit near Lan Sang Village, Tak Province, northern Thailand. *Proc. Geol. Conf. Chiangmai Univ.*, Chiangmai.
- Scharer, U., P. Tapponnier, R. Lacassin, P. H. Leloup, Zhong Dalai, and Ji Shaocheng, 1990. Intraplate tectonics in Asia: a precise age for large-scale Miocene movement along the Ailao Shan-Red River shear zone, China, *Earth Planet. Sci. Lett.*, 97, 65-77, 1990.
- Schärer, U., Zhang Lian-Sheng. and P. Tapponnier, Duration of strike-slip movements in large shear zones: The Red River belt, China, *Earth Planet. Sci. Lett.*, 126, 379-397. 1994.
- Shackleton, N. J. et al. 1990: An alternative astronomical calibration of the Lower Pleistocene time scale based on ODP Site 677. *Trans. of Royal Soc. of Edinburgh: Earth Sciences* 81, pp 251-261.
- Schmidt G., Zhou L., and Wasson, J. T., 1993, Iridium anomaly associated with the Australasian tektite-producing impact: masses of the impactor and of the Australasian tektites: *Geochimica et Cosmochimica Acta*, v. 57, p. 485 1-4859.
- Schneider, D. A. and D.V. Kent, Ivory coast microtektites and geomagnetic reversals. *Geophys. Res. Lett.* 17, 163-166, 1990.
- Schneider, D. A., Kent, D. V. and Mellow, G. A., 1992. A detailed chronology of the Australasian impact event, the Brunnes-Matuyama geomagnetic polarity reversal, and global climate change. *Earth Planet. Sci. Lett.*, 111, 395-405.
- Schnetzler C. C., Pinsen W. H. & Hurley, P. M. 1966. Ivory Coast tektites and Lake Bosumtwi. *Science* 151, 817.
- Schnetzler, C. C.; J.A. Philpotts and H.H. Thomas. 1967. Rare-earth and barium abundances in Ivory Coast tektites and rocks from the Bosumtwi Crater area, Ghana, *Geochim. Cosmochim. Acta* 31. 1987-1993.
- Schnetzler, C. C., 1992. Mechanism of Muong Nong-type tektite formation and Speculation on the source of Australasian tektites. *Meteoritics* 27, 154-164.
- Schnetzler, C. C.. and J. F. McHone. 1996. Source of Australasian tektites: Investigating possible impact sites in Laos, *Meteorit. Planet. Sci.*, 31,73 - 76.
- Schnetzler, C. C. Walter, L. S. and Marsh, J. G., 1988. Source of the Australasian tektite strewn field: a possible offshore impact site. *Geophys. Res. Lett.* 15, 357-360.
- Schnetzler, C. C., 1992. Mechanism of Muong Nong-type tektite formation and speculation on the source of Australasian tektites, *Meteoritics* 27 (1992) 154- 165.
- Schnetzler, C. C., et al. 1992. "Search for the 700,000-year-old source crater for the Australasian tektite strewn field." Dressler, B. O., chairperson, et al. 1992. (LPI CONTRIBUTION; Vol.790. p. 63-64)
- Schnetzler, Charles C. 1992 "Mechanism of muong nong-type tektite formation and speculation on the source of Australasian tektites." June 1992. (METEORITICS; Vol.27, No.2. p. 152-165)
- Schnetzler. C. C.. et al. 1993."Where in the world is the Australasian tektite source crater?." Oct. 26. 1993. (Eos. Transactions. American Geophysical Union ; Vol. 74. No. .13, Suppl.. p. 388-389)
- Schmidt, G., Zhou, L. and Wasson, J. T., 1993. Iridium anomaly associated with the Australasian tektite-producing impact: Masses of the impactor and the Australasian tektites. *Geochimica et Cosmochimica Acta* 57, 4851-4859.
- Schweller, W. J., Karig, D. E., and Bachman, S. B. 1983. Original setting and emplacement history of the Zambales ophiolite, Luzon, Philippines, from stratigraphic evidence. Hayes, D. E. (ed.) *The tectonic and geologic evolution of Southeast Asian seas and islands*, part 2. *Geophysical monograph* 27 (ed. D. E. Hayes) , pp. 124-38. American Geophysical Union, Washington.
- Schweller, W. J., Roth, P. H., Karig, D. E., and Bachman, S. B. 1984. Sedimentation history and biostratigraphy of ophiolite-related Tertiary sediments, Luzon, Philippines. *Geological society of America bulletin* 95,1333-42.
- Schwarz, H. P., 1962. A possible origin of Tektites by soil fusion at impact site. *Nature* 194, 8-10.
- Schindewolf, O. H., 1963. Neokastrophim? *Deutsch Geol. Gesell. Zeitschr.*, 114, 430-445.
- Schoch, R.M., Hsü, K. J. and Van Valen, L. M., 1987. Darwin's three mistakes; discussion and reply. *Geology*, 15, 175-178.
- Sclater, J. G. and Tapscott, C. 1979. The history of the Atlantic. *Scien. Am.* 240, 156-174.
- Sclater, J. G. and Christie, P. A. F. (1980). Continental stretching: an explanation of the post Mid Cretaceous subsidence of the central North Sea Basin. *Journal*

- of geophysical research 85, 3711-39.
- Sclater, J. G. and Fisher, R. L. 1974. Evolution of the East Central Indian Ocean, with emphasis on the tectonic setting of the Ninetyeast Ridge. Geological society of America Bulletin 85, 583-702.
- Scott, A. C., and Jones, T. P., 1991, Microscopical observations of recent and fossil charcoal: Microscopy and Analysis, July, p. 13-15.
- Schmidt, G., Zhou, L., and Wasson, J. T., 1993. Iridium anomaly associated with the Australasian tektite-producing impact: masses of the impactor and of the Australasian tektites: *Geochimica et cosmochimica Acta*, v. 57, p.4951-4859.
- Schneider, D. A., Kent, D. v., and Mello, G. A., 1992, A detailed chronology of the Australasian impact event: the Brunhes, & Matuyama geomagnetic polarity reversal, and global climate change: Earth and Planetary Science Letters, v. 11, p. 395-405.
- Schnetzler, C. C. et al. 1988. "Source of the Australasian tektite strewn field; a possible offshore impact site." (*Geophysical Research Letters*; 15, No. 4, 357-360.
- Schnetzler, C. C., 1992. Mechanism of Muong Nong-type tektite formation and speculation on the source of Australasian tektites: *Meteoritics and Planetary Science*, v. 27, p. 154-165.
- Scott, A. C., and Jones, T. P., 1991, Microscopical observations of recent and fossil charcoal : Microscopy and Analysis, July, p. 13 - 15.
- Sengor, A.M.C. and HsU, K. 1984: The Cimmerides of Eastern Asia; history of the eastern end of Palaeo-Tethys. *Mdm. Soc. geol. France, N.S.*, no. 147, p. 139-167.
- Sekenina, Z. 1985. .in. *Comets* (Wilkening, L.L., ed.) Univ. Arizona, Tucson, 251-287
- Seyfert, C. K. and Sirkin, L.A., 1973. *Earth History and Plate Tectonics*. Harper & Row Publishers, New York. 505 pp.
- Shackleton, N.J. and Opdyke, N. D., 1976. Oxygen-isotope and paleomagnetic stratigraphy of Pacific core V28-239, late Pliocene to latest Pleistocene. *Geol. Soc. American Mem.* 145, p. 449-464.
- Shaw, H. F. and Wasserburg, G. J., 1982. Age and provenance of the target materials tektites and possible the impactites as inferred from Sm-Nd and Rb-Sr systematics. *Earth. Planet. Sci. Lett.* 60, 155-177.
- Shen, C.; Beer, J.; Oesshger, H.; Bonani, G.; Suter, M., and Wolfli, W., 1992. 10BE in Chinese loess. *Earth. Planet. Sci. Lett.*, 109, 169-177.
- Shergold, J. Burrett, C. F., Akerman, T. and Stait, B., 1988. Late Cambrian trilobites from Tarutao island, Thailand. *New Mexico Bureau of Mineral Resources Memoir* 44, 303-320.
- Shoemaker, E. M. 1984. Large body impacts through geologic time. In H. D. Holland and A. F. Trendall (eds.), "Patterns of Change in Earth Evolution", Springer-Verlag, N.Y., 15-40.
- Shoemaker, E. M., H. R. Uhlherr, 1999. Stratigraphic relations of australites in the Port Campbell embayment, Victoria, *Meteorit. Planet. Sci.* 34, 369-384.
- Showalter, D. L. et al., 1971. A comparison between Java tektite (J2) and lunar rock 12013 (abstr.). *Meteoritics* (94), 315-316.
- Shuvalov, V. S., 2002, Radiation effects of the Chicxulub impact event, in Buffetaut, E., and Koeberl, C., eds., *Geologic and Biological Effects of Impact Events*: New York, Springer.
- Sibava, V., 1993. Quaternary Sequences in Southeast Asia and their regional correlation. In: ESCAP Atlas of Stratigraphy XII., Quaternary Stratigraphy of Asia and the Pacific, IGCP 296. Mineral Resources Development Series, No 62. United Nations. pp. 1 -29.
- Sieh, S., Herrin, J., Jicha, B., Schonwalder, D., Moore, J., Banerjee, P., Wiwegwin, W., Sihavong, V., Singer, B., Chualawanich T., and Charusiri, P., 2020. Australasian impact crater buried under the Bolaven volcanic field, Southern Laos. *Proceedings of the National Academy of Science*, 117 (3) 1346-1353; <https://doi.org/10.1073/pnas.1904368116>
- Sigurdson, H.; De'Hondt, S.; Arthur, M. A.; Bralower, T. J.; Zachos, J. C.; Fossen, M. and Channel E. T., 1991. Glass from Cretaceous/Tertiary boundary in Haiti. *Nature*, 349, 482-487.4, 6, 7 and 9.
- Silaratana, T.; Ratanasathien, B.; Tkayasu, K.; Asanachinda, P., Kantharosa, W.; and M.. Kusakabe, 2003. RCPNS 46-8th International Congress on Pacific Neogene Stratigraphy. 2-9 February 2003, Chiang Mai University, Chiangmai, Thailand.
- Silver, L. T. & Schultz, P. H. (eds.). 1982. Geological implications of impacts of large asteroids and comets on the Earth. *Geol. Soc. Amer. Spec. pap.* 190, 528pp.
- Silver, E. A., Reed, D., McCaffrey, R., and Joyodiwiryo, Y. 1983. Back arc thrusting in the eastern Sunda Arc, Indonesia: a consequence of arc-continent collision. *Journal of Geophysical Research* 88, 7429-48.
- Simonson, B. M. and Glass, B.P., 2004, Spherule layers-Records of ancient impacts: *Annual Review of Earth and Planetary Sciences*, v. 32, p. 329-361, doi: 10.1146/annurev.earth.32.101802.120458.
- Sinsakul, Sin; Niran Chaimanee, and Suwat Tiyaipairach, 2002. Quaternary Geology of Thailand. The Symp. on Geology of Thailand. 26-31 August 2002, Bangkok, Thailand. 170-180.
- Siraj, A., Loeb, A., 2021. Breakup of a long-period comet as the origin of the dinosaur extinction. *Sci Rep* 11, 3803 . <https://doi.org/10.1038/s41598-021-82320-2>
- Sithiprasana, D., 1959. Occurrences of Mae Moh fossils. . *Rept. of the Investigation 2. Roy. Dept. Mines, Bangkok*. P. 29-34, pls. 3-4, 6, 7 and 9.
- Shuvalov', V S., 2002, Radiation effects of the Chicxulub impact event, in Buffetaut, E., and Koeberl, C., eds., *Geologic and Biological Effects Compact Events*: New York, Springer.
- Soons Jane M., 1963. The glacial sequence in part of Rakaia Valley, Canterbury, New Zealand. *N. Z. geol. Geophys.* 6, 735-56.
- Soeradi, T. et al. 1985: *Geology and Stratigraphy of the Trinil Area*. In: Watanabe W. and Kadar, D. (eds.) *Quaternary Geology of the Hominid Fossil-bearing Formations in Java*, pp 49-53, Bandung.
- Somboon, J. P. R. 1988: Paleontological study of the recent marine sediments in the lower central plain, Thailand. *Journal of Southeast Asian Earth Sciences*, Vol 2, Nos 3/4, pp 201-210, Pergamon Press.
- Sompong, W., Springbeit, G. M., and Evans, P. R., 1996. Lignite mine development project (Phase III) Mae Moh coal deposit Electricity Generation Authority of Thailand, Australia AID Geological Report.
- Sonsuk, M., Hastings, P. 1984: An age from the Yasothon soil Series in the Sakhon Nakhon Basin. *J. Geol. Soc. Thailand*, Vol 7, No 1, pp 1-11.
- Spencer, L. J. 1933. Origin of tektites. *Nature* 131. 117-118, 1933.
- Smit, J. and Hertogen, J. 1980. An extraterrestrial event

- at the Cretaceous-Tertiary boundary, *Nature*, London, 235, 198-200.
- Smit, J. and Romein, J. T., 1985. A sequence of event across Cretaceous-Tertiary boundary. *Earth Planet. Sci. Lett.*, 74, 155-170.
- Smith, R. B 1987. Diachronous Neogene microcontinent collision in Buton and Eastern Sulawesi, Indonesia. *Geological Society of America Abstracts with Programs* 19, 850.
- Smit, J. and Smoluchowski, R., 1981. Hawaii as a consequence of an impact: Precipitation-driven magma convection. *Abstracts of Snowbird Houston, Lunar and Planetary Institute*, 53.
- Smit, J., van Eijden, A.J.M., and Troelstra, S.R., 1991, Analysis of the Australasian microtektite event, the Toba Lake event, and the Cretaceous/Paleogene boundary, Eastern Indian Ocean, in *Proceedings, Ocean Drilling Program, scientific results, Volume 121: College Station, Texas, Ocean Drilling Program*, p. 489-503.
- Spry, A. and Banks, M. R., eds., 1962. *The Geology of Tasmania*. *Jour. Soc. Aust.*, 9, 107-362.
- Staudacher, T., Sarda, P., and Allegre, C.J., 1990. Noble gas systematics of Reunion Island, Indian Ocean. *Chem. Geol.* 89, 1-17
- Strahler, A. N., 1973, *Environmental geoscience: interaction between natural systems and man*: New York, John Wiley
- Stauffer, P. H. 1958. Studies in the Crocker Formation, Sabah. Borneo region of Malaysia Geological Survey Bulletin 8, 1-13.
- Stauffer, P. H. 1973a. Cenozoic. *Geology of the Malay Peninsula* (ed. D. J. Gobbett and C. S. Hutchison), pp. 143-75. John Wiley-Interscience, New York.
- Stauffer, P. H. 1973b. Kenny Hill Formation. *Geology of the Malay Peninsula* (ed. D. J. Gobbett and C. S. Hutchison), pp. 87-97. John Wiley-Interscience, New York.
- Stauffer, P. H. 1974. Malaya and Southeast Asia in the pattern of continental drift. *Geological Society of Malaysia Bulletin* 7, 89-138.
- Stauffer, P. H., 1978. Anatomy of the Australasian strewn field and the probable site of its source crater. In Nutalaya, P. ed. *Proceedings of the Third Regional Conference and Mineral Resources on Southeast Asia*, pp. 285-892. Asian Institute of Technology, Bangkok
- Stauffer, P. H., 1983. Phantom tektite localities of Borneo. *Meteoritics* 18 (1), 9-13.
- Stauffer, P. H., and Lee, C. P. 1986. Late Paleozoic glacial marine facies in Southeast Asia and its implications. *Bulletin of the Geological Society of Malaysia* 20, 363-97.
- Stauffer, P. H. and Mantajit, J. 1981. Late Paleozoic tilloids of Malaya, Thailand and Burma. *Earth's pre-Pleistocene glacial record* (ed. M. J. Hambrey and W. B. Harland), pp. 331-7. Cambridge University Press, England.
- Stauffer, P. H., Nishimura, S., and Batchelor, B. C. 1980. Volcanic ash in Malaya from a catastrophic eruption of Toba, Sumatra 30,000 years ago. *Physical Geol. of Indonesian island arcs* (ed. S. Nishimura), pp. 156-64. Kyoto University Press.
- Stein, S. and Okal, E. A. 1978. Seismicity and tectonics of the Ninetyeast Ridge area: evidence for internal deformation in the Indian Plate. *Journal of Geophysical Research* 83, 2233-44.
- Stephenson, D. and Marshall, T. R. 1984. The petrology and mineralogy of Mt Popa volcano and the nature of the Late-Cainozoic Burma volcanic arc. *Journal of the Geological Society London* 141, 747-62.
- Stirton, Ruben Arthur, 1963: *Time, Life and Man*. Science Edition, John Wiley & Sons Inc., New York. 358 pp.
- Stöffler, D., 1984. Glasses formed by hypervelocity impact. *J. Non. Cryst. Solids* 67, 465-502.
- Stöffler, D., D. F. Gault, J. Wedekind, and O. Tokusaki., 1975. Experimental hypervelocity impact into quartz sand: Distribution and shock metamorphism of ejecta, *J. Geophys. Res.*, 80, 4062-4077.
- Stokes, R. A., and Loh, I. C., 1982, Streamflow and solute characteristics of a forested and deforested catchment pair in south-western Australia, in O' Loughlin, E. M., and Bren, L. J., eds., *National Forest Hydrology Symposium, I: Australia, Institution of Engineers*
- Storzer D., Wagner G. A. (1969): Correction of thermally lowered fission track ages of tektites. -*Earth and Planet Sci. Lett.* ,5, N 7, 463-468.
- Storzer- D. Wagner'G' A. (1977): Fission track dating of meteorite impacts. - *Meteoritics* 18, N 2, 368- 369.
- Storzer D. Gentner, W. 1970. Micromoldavites from the Bavarian molasse. *Meteoritics* 5, 225.
- Storzer, D. and Wagner, G. E., 1971. Fission track of North American tektites. *Earth Planet. Sci. Lett.* 10, 435-440
- Storzer, D. and Wagner, G. E., 1980. Australites older than indochinites. *Naturwissenschaften* 67, 90-91.
- Storzer, D.; Jessberger, E. K., Klay, N.; and Wagner, G. A., 1984.  $^{40}\text{Ar}/^{39}\text{Ar}$  evidence for two discrete tektite-forming events in the Australian Southeast Asian area. *Meteoritics* 19: 317.
- Storzer, D., et al. "40 Ar-30 Ar evidence for two discrete tektite-forming events in the Australian-Southeast Asian area." 1984. (*METEORITICS* ; Vol. 19, No. 4, p. 317)
- Sverdrup, H.U., Johnson, M.W. and Fleming, R.H., 1942. *The Oceans*, Prentice Hall Inc, Englewood Cliffs, NJ.
- Sublette, C., 1997, Nuclear weapons frequently asked questions (NWFAQ), in *The High Energy Weapons Archive: A Guide to Nuclear Weapons*: (URL) [http: // www. ciar. Org/~ttk/hew/hew](http://www.ciar.org/~ttk/hew/hew).
- Suess, F. E., 1900. Die herkunft der moldavite. *Jahrb. Geol. Reichsanst.* (Bundesanst.), Wien. 50: 193-382.
- Suess, H. F., and H. C. Urey, Abundances of the elements, *Rev. Mod. Phys.*, 28. 53-74, 1956.
- Suess, H. F., R. J. Hayden, and M. G. Inghram, Age of tektites. *Nature*. 168, 432-433. 1951.
- Suggate, R. P., 1965. Late Peistocene of northern part of South Island, N.Z. *Bull. of Geol. Survey N. Z.*, 77.
- Sumarso and Ismoyovati, T. 1976. Contribution to the stratigraphy of the Two Hills and their southern surroundings (central Java). *Proceedings of the fourth meeting of the petroleum association, Jakarta, June 1975* 2, 79-25.
- Suraprasit K, Jaeger JJ, Chaimanee Y, Benammi M, Chavasseau O, Yamee C, Tian P, Panha S. 2015. A complete skull of *Crocota crocuta ultima* indicates a late Middle Pleistocene age for the Khok Sung (northeastern Thailand) vertebrate fauna. *Quaternary International* 374: 34-45. doi: 10.1016/j.quaint.2014.12.062
- Suraprasit K, Jaeger J-J, Chaimanee Y, Chavasseau O, Yamee C, Tian P, Panha S. 2016. The Middle Pleistocene vertebrate fauna from Khok Sung (Nakhon

- Ratchasima, Thailand): biochronological and paleobiogeographical implications. *ZooKeys* 613: 1–157. doi: 10.3897/zookeys.613.8309
- Suraprasit, K., Bocherens, H., Chaimanee, Y., Panha, S., and Jaeger, J.-J. 2018. Late Middle Pleistocene ecology and climate in Northeastern Thailand inferred from the stable isotope analysis of Khok Sung herbivore tooth enamel and the land mammal cenogram. *Quaternary Science Reviews*, 193:24–42. <https://www.sciencedirect.com/science/article/pii/S0277379117308247>
- Suraprasit, K., Jongautcharyakul S., Yamee, C., Potichaiya, C., and Bocherens, H. 2019. New fossil and isotope evidence for the Pleistocene zoogeographic transition and hypothesized savanna corridor in Peninsular Thailand. *Quaternary Science Reviews*, 221:105861. <https://doi.org/10.1016/j.quascirev.2019.105861>
- Suteethorn, S., Leuff, J. and Suteethorn, V. 2007: Additional well preserved Vertebrate of Phuwiangosaurus sirdhornae from the Type Locality (Phu Wiang, Changwat Khon Kaen) . (GEOTHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. Geothai 2007, p. 17. 21-22 November. Imeral Hotel, Bangkok, Thailand.
- Suttirat, C.; Charusiri, P.; Farrar, E. and Clark, A., H. 1994. New <sup>40</sup>Ar/<sup>39</sup>Ar Geochronology and characteristics of some Cenozoic basalts in Thailand. Proc. of Intern. Symposium on: Stratigraphic Correlation of Southeast Asia. 15-20 Nov. 1994. Bangkok, Thailand.
- Sutthirat, Chakkaphan; Giles Droop and Michael Henderson, 2005. Thermobarometry and Speedometry of Thai Corundum-Related Basalts: Evidences for their Petrogenesis. Proc. of the Int. Conference on Geology, Geotechnology and Mineral Resources of Indochina. GEOINDO 2005, pp. 489-487. Kosa Hotel, Khon Kaen, 28-30 Nov. 2005.
- Suzuki, M. et al. 1985: Fission Track Ages of Pumice Tuff, Luff Layers and Javites of Fossil-bearing Formations in Sangiran Area, Central Java. In: Watanabe N., and Kadar, D. (eds.) Quaternary Geology of the Hominid Fossil-bearing Formations in Java, pp 309-331, Bandung.
- Takashima, 1., and Honda, S. 1989. Comparison between K-Ar and TL dating results of pyroclastic flow deposits in the Aizutajima area, Northeast Japan. *Journal of Geological Society* 95: 807-816.
- Tamara, T., 1992: Landform development and related environment changes in Chi River Basin, Northeast Thailand. The Science Series Report of the Tohoku University, 5th Series (Geography), vol. 42 (2): 107-127.
- Tapponnier, P., and P. Molnar, Slip-line field theory and large-scale continental tectonics, *Nature*, 264, 3 19-324, 1976.
- Tapponnier, P., and P. Molnar, 1977. Active faulting and tectonics in China, *J. Geophys. Res.*, 82, 2905-2930, 1977.
- Tapponnier, P., G. Peltzer, and R. Armijo, 1986. On the mechanics of the collision between India and Asia, in *Collision Tectonics*, edited by Coward, M. P., and A. C. Ries, Geological Society Special Publication, 19, pp. 115-157, Geological Society, London, 1986.
- Tapponnier, P., G. Peltzer, R. Armijo, A.-Y. Le Dam, and P. Cobbold, 1982 Propagating extrusion tectonics in Asia: new insights from simple experiments with plasticine, *Geology*, 10, 611-616, 1982.
- Tapponnier, P., R. Lacassin, P. H. Leloup, U. Schwer, Zhong Dalai, Liu Xiaohan, Ji Shaocheng, Zhang Lianshang, and Zhong Jiayou, 1990. The Ailao Shan/red River metamorphic belt: Tertiary left-lateral shear between Indochina and South China, *Nature*, 343,431-437, 1990.
- Tassy, J., P. Anupandhanant, L. Ginburg, P. Mein. B. Ratanasathien, and V. Suteethon, 1992. New *Stegolophodon* (proboscidea, Mammalia) from the early Miocene of Northern Thailand. *Geobios*, No. 25, Fasc. 4. 511-523.
- Tauxe, L., J. P. Valet, and J. Bloemendal. *Magnetostratigraphy of Leg 108 advanced hydraulic piston cores*. Proc. ODP, Sci. Results. 108: 4290-452. 1989.
- Taylor, B., and Hayes, D.E., 1983. Origin and history of the South China Basin, In: *The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands* (ed. D.E. Hayes), Am. Geophys. Union Geophys. Monogr. Ser., 27, 25-26.
- Taylor, H. P. and S.E. Epstein. Oxygen isotope studies of Ivory Coast tektites and impactite glass from the Bosumtwi crater, Ghana, *Science* 153, 173. 175. 1966.
- Taylor. H. P. and Epstein. SE., 1969. Correlations between <sup>18</sup>O/<sup>16</sup>O ratios and chemical compositions of tektites. *J. Geophys. Res.* 74: 6844-6844.
- Taylor, S. R., 1962. The chemical composition of australites. *Geochim. Cosmochim. Acta* 26, 685-722.
- Taylor S. R. 1969. Criteria for the source of australites. *Chemical Geology* 4, 451-459.
- Taylor S. R. & Kaye, M. 1969. Genetic significance of the chemical composition of tektites: A review. *Geochimica et Cosmochimica Acta* 33, 1083 - 1100.
- Taylor, S.R. & McLennan, S.M. 1985. *The Continental Crust: Its Composition and Evolution*, Blackwell, Oxford., 312 pp.
- Taylor, S. R., 1973, Tektites: A post-Apollo view: *Earth-Science Reviews*, v. 9, p. 101-123.
- Taylor, S. R. and McLennan, S. M., 1979. Chemical Relationships among irghizites, zhamanshininites, Australian tektites and Henbury impact glasses. *Geochim. Cosmochim. Acta* 43, 1551-1565.
- Taylor, S.R., and McLennan, SM., 1995, *The geochemical evolution of the continental crust: Reviews of Geophysics*, v. 33, no. 2, p. 241–265.
- Teihard de Chardin, P. & Young, C.C., 1930. Preliminary observations on the preloessic and post-Potian formations in west Shansi and Northern Shensi. *Mem. Geol. Soc. China* 8, ser A.
- Thein, J., 1987. A tektite layer in upper Eocene sediments of the New Jersey continental slope (Site 612, Leg 95). In: C. W. Poag, A. B. Watts et al. (eds.) In it. *Deep-Sea Drill. Proj.*, 95, 564-579.
- Trinh Tho and Ngugen Din Huu, (eds.). 1979. *Geology of Vietnam (North Part)*, Geol. Dept. of Geology, Research Inst. of Geol. And Miner. Resources. Soc. Republic of Vietnam, Hanoi.
- Thiramongkol, N. 1983: Reviews of Geomorphology of Thailand. In Thiramongkol, N., Pisutha Arnon V. (eds.), *Proceedings of the first Symposium on Geomorphology and Quaternary Geology of Thailand*, pp 6-23, Bangkok.
- Thiramongkol, N., 1986. Neotectonism and Rate Of Uplift in the Eastern Margin of The Lower Central Plain of Thailand. In: Thiramongkol (ed.), *Proceeding of the Workshop on Economic, Geology, Tectonics Sedimentary Processes and Environmental of the*

- Quaternary of Southeast Asia. Pp. 35-44, Bangkok.
- Thiramongkol, N. (ed.) 1989: Proceeding of the Workshop on Correlation of Quaternary successions in South, East and Southeast Asia. 269p., Chulalongkorn University, Bangkok.
- Thiramongkol, N., Pisutha Arnond, V. (eds.) 1983: Geomorphology and Quaternary Geology of Thailand. 197p., Bangkok.
- Thom, B. G. 1965. Late Quaternary coastal geology of Port Stephens-Myall Lake area. N. S. W. J. Proc. R. Soc. N. S. W. 98 26-36.
- Tindale, N. B. 1959. Ecology of primitive aboriginal man in Australia in Biogeography and Ecology in Australia. Junk, the Hague, pp. 36-51.
- Tjia, H. D., 1989: Active tectonics in the Indonesian Archipelago. In N. Thiramongkol (ed.), Proceedings of the Workshop on Correlation of Quaternary Successions in South, East and Southeast Asia, pp 165-185, Bangkok.
- Toon, O., K. Zahnle, D. Morrison, R.P. Turco, C. Covey. 1997. Environmental perturbations caused by the impacts of asteroids and comets, Rev. Geophys. 35 (1997) 41- 78.
- Toon, O. B., Pollack, J. B., Ackerman, T. P., Turco, R. P., McKay, C. P., & Liu, M. S., 1981. Evolution of an impact-generated dust cloud and its effects on the atmosphere. Geol. Soc. Am. Spec. Pap., 190, 187-201.
- Toon, O. B., Pollack, J. B., Ackerman, T. P., Turco, R. P., McKay, C. P., & Liu, M. S., 1982. Evolution of an impact-generated dust cloud and its effects on the atmosphere. Geol. Soc. Am. Spec. Pap., 190, 187-201
- Toon, O., K. Zahnle, D. Morrison, R.P. Turco, C. Covey, 1997. Environmental perturbations caused by the impacts of asteroids and comets, Rev. Geophys. 35 (1997) 4 1-78.
- Tomatsu, S. 1995. Nagasaki, 11:02. August 9, 1945: Tokyo, Schinchosha Company.
- Turcotte, D. L., 1986. Fractals and fragmentation. J. Geophys. Res. 91, 192 1-1926.
- Tweedie, M, 1977. The World of Dinosaurs. Printed in Great Britain by Morrison & Giibb Ltd., London and Edinburgh.
- Udomchoke, V., 1989. Quaternary stratigraphy of the Khorat plateau area, Northeast Thailand. Proc. of the Workshop on Correlation of Quaternary Successions in South, East and Southeast Asia, 69-94; Bangkok.
- Urey, H. C., 1971. Tektites from the Earth. Science 171, 312-314.
- Urey, H. C., 1973. Cometary collision and geological periods. Nature, London. 242, 32-33.
- Uttamo, Wutti, Gary J. Nichols and Chris F. Elde, 1999. The Tertiary Sedimentary Basins of Northern Thailand. Symposium on Mineral, Energy, and Water Resources of Thailand: Towards the year 2000. October 28-29, 1999, Bangkok, Thailand
- Uttamol, W., C. Elders, and G. Nichols. 2003. Sedimentology of Neogene sequences in the Mae Moh basin, northern Thailand. RCPNS 46 - 8th International Congress on Pacific Neogene Stratigraphy, Chiang Mai - February 2003.
- Valentin, H., 1961. The central west coast of Cape York Peninsula, Austr. Geogr. 8. 65-72.
- Van Bemmelen, R. W. 1970. The geology of Indonesia, IA; General geology of Indonesia and adjacent archipelagoes, 2; Economic geology, 1B; Portfolio and index, 2nd edn. Martinus Nijhoff, The Hague
- Van Leeuwen, T. M. 1981. The geology of southwest Sulawesi with special reference to the Biru area. The geology and tectonics of eastern Indonesia. Geological Research and Development Centre, special publication 2 (ed. A. J. Barber and S. Viryosujono), pp. 277-304. Bandung.
- Van Patter, D. M., et al. "Proton probe analysis of an irghizite and a high-magnesium Java tektite." 1981. (Geochimica Et Cosmochimica Acta; Vol.45, No.2, p.229-Ta)
- Varne, R. and Foden, J. D. 1985. Geochemical and isotopic systematics of eastern Sunda arc volcanics: implications for mantle sources and mantle mixing processes. Origins of Arcs (ed. F. C. Wezel), pp. 159-89. Elsevier, Amsterdam.
- Veevers, J. J. 1982. Western and northwestern margin of Australia. The Ocean basins and margins: The Indian Ocean, 6 (ed. A. E. M. Nairn and F. G. Stehli), pp. 513-44. Plenum Press, New York.
- Veevers, J. J. (ed.) 1982a. Phanerozoic earth history of Australia, Oxford geological sciences series 2. Clarendon Press, Oxford.
- Vening Meinesz, F. A. 1954. Indonesian archipelago geophysical study. Geological society of America bulletin 65, 143-64.
- Vannucci, R., Vannucci, S. Mazzucotelli, A. Meloni, S. and Odenne, M. 1981. Considerazioni geochimiche su alcuni terreni argillosi emarnoso-calcarei in un intervallo comprendente il limite Cretaceo-terziario della Scaglia rossa umbro-marchigiana. Rend. Scoc. Ital. Mineral. Petrogr., 38, 413.
- Veevers, J. I. 1988. Morphotectonics of Australia's northwestern margin - a review. In 'The North West Shelf, Australia, Proceedings of PESA Symposium, Perth'. 19-27, (Petroleum Exploration Society of Australia: Perth.)
- Vella, P., 1983. Tensional Faults in Tertiary Strata, Mae Moh. J. Geol. Soc. Thailand, Thailand. 6 (1), 13-14.
- Vella, P., Bunopas, S. and Kaewyana, W., 1987. A loess covered alluvial gravel terrace near Chiangmai, Northern Thailand. Geological Society of Thailand Newsletter, 1987 (3), pp 29-31.
- Verbeck, R.D.M.. 1897. Glaskogels van Billiton. Jaarb. Mijnwez. Ned Indië. 1897: 235-272.
- Verosub, K. L., 1977. Depositional and postdepositional processes in the magnetization of sediments, Rev. Geophys. Space Phys. 15, 129-143. 1977.
- Vickery, A. M. and Browning, L., 1991. Water depletion in tektites (abstract). Meteoritics 26, 403.
- Vichiansilp, J. 1997. Study on some volcanics in south-Northeastern Thailand. Satellite imagery Project National Research council of Thailand.
- Vozenin-Serra, C. and Prive-Gill, C. 1989: Bois Pliopleistocenes du gisement du Saropee, Plateau de Khorat, est de la Thaïlande. Rev. of Palaeobotany and palynology 60: 225-254.
- Vsechsvyatski (1968): The nature and genesis of comets and meteorites. - Moscow, prosvesh-chenie". p.p. (In Russian).
- Walter, L. S., et al. "Search for the Australasian tektite source crater." Prinz, Martin, chairperson. 1986. Abstracts And Program Of The 49th Annual Meeting Of The Meteoritical Society.
- Walther, L. S., 1965. Coesite discovered in tektites. Science 147, 1029-1032.
- Wampler, J.M.; D. H. Smith and A. E. Cameron, 1969. Isotopic comparison of lead in tektites with lead in

- earth materials. *Geochim. Cosmochim. Acta* 33, 1045-1055, 1969.
- Wang, J., Q. Zhao, X. Cheng, R. Wang, and P. Wang (2000). Age estimation of the mid-Pleistocene microtektite event in the South China Sea: A case showing the complexity of the sea-land correlation, *Chin. Sci. Bul.*, 45, 2277-2280.
- Wan Tianfeng, 1999. On the boundary between the Gondwanaland and Eurasia continent seen in China. *Gondwana Research* 2 (4), 611-613.
- Wang, P., J. Tian, X. Cheng, C. Liu, J. Xu, Carbon reservoir changes preceded major ice-sheet expansion at the mid-Brunhes event. *Geology* 31(2003) 239-242.
- Wang, Y.Y., Sadao, S., Teng, Z.H., Lei, X.Y. & Sun, W., 1984. Loess in China and its stratigraphic sequence. in: *The recent research of loess in China* (eds. Sadao, S. & Wang, Y.Y.), 1-31, Wako Printing, Ltd. Kyoto.
- Wang, Y. and Sasajima, Sadao, (eds.) 1985. The new development of loess studies in China. University of Shanxi, China, 208 pp. (in Chinese)
- Waring, G.A., 1965. Thermal springs of the United States and other countries of the world. A summary. USGS Prof. Paper 492, 383pp.
- Wasson, J. T., 1985. Meteorites: Their Record of Early Solar System History, Freeman, New York.
- Wasson, J. T., 1987. A multiple-impact origin of Southeast Asian Tektites (Abstr.). *Lunar Planet Sci.* 18, 1062-1063.
- Wasson, J. T., 1989. Climate and tektite origin (abstract). *Meteoritics* 24, 337-338.
- Wasson, J., T. 1991. Layered tektites: a multiple impact origin for the Australasian tektites. *Earth Planet. Sci. Lett.* 102: 95-109.
- Wasson, J. T. and Kallemeyn, G.W., 1988. Compositions of chondrites. *Phil. Trans. Roy. Soc. London A325*, 535-544.
- Wasson, J. T. and Heins, W. A., 1993. Tektites and climate. *J. Geophys. Res.* 98, 3043-3052.
- Wasson, J.T. and G.W. Kallemeyn, 1988. Composition of chondrites. *Phil. Trans. Roy. Soc. London A323*. 535-544.
- Wasson, J. T., Phitakpaivan, K., Putthapiban, P., Sulyapongse, S., Thathimthong, B., and McHorne, J. F. 1995: Field recovery of layered tektites in northeast Thailand: Evidence of a large scale melted sheet. *J. Geophys. Res.*, 100: 14385-14390.
- Watanabe, N., Kadar, D., eds. 1985. Quaternary Geology of the Hominid Fossil Bearing in Java. *Geol. Res. Devel. Centre Sp. Paper No. 4*. 378p. Bandung.
- Weissman, P. R., 1980. Physical loss of long-period comet. *Astron. Astrophys.* 85, 191-196.
- Weeks, R. J., Underwood, J. R., JR. and Giegengack, R., 1984. Libyan desert glass; In; L. D. Pye, J. A. O'Keefe and V. D. Frechette (eds.), *Natural Glasses*. North Holland Phys. Publ. Amsterdam, pp. 593-619.
- Weissel, J. K. 1980. Evidence for Eocene oceanic crust in the Celebes Basin. Hayes, D. E" (ed.) *The tectonic and geologic evolution of Southeast Asian seas and islands*, *Geophysical Monograph* 23 (ed. D. E. Hayes), pp. 3747. American Geophysical Union, Washington.
- Weissel, J. K. and Anderson, R. N. 1978. Is there a Caroline Plate? *Earth and planetary science letters* 41, 143-59.
- Wetherill, G. W. and Shoemaker, E. M., 1982. Collision of astronomically observable bodies with the Earth. In: *Geological Implications of Impacts of Large Asteroids and Comets on the Earth*. L. T. Silver and P. H. Schulz, eds., pp. 1-13. *Spec. Pap.* 190. Geol. Soc. Am., Boulder, Colo.
- Wezel, F. C., Vannucci, R., and Vannucci, S., 1981 De'couverte de divers niveaux riches en iridium dans le "Scaglia Bianca" de l'Apennine d'Ombrie-Marches (Halle). *C. R. Acad. Sci. Paris*, 293, 837.
- Wheller, G. E., Varne, R., Foden, J. D., and Abbom, M. J. (1987). *Geochemistry of Quaternary volcanism in the Sunda-Banda Arc, Indonesia, and three-component genesis of island arc basaltic magmas*. *Geochemistry of eruptive magmas and tectonic controls on petrogenesis* (ed. S. D. Weaver and R. F. Johnson), *Journal of Volcanology and geothermal research* (in press).
- Whitford-Stark, J. L., 1987. A survey of Cenozoic volcanism on mainland Asia. USGS Spec. Paper 213, 73pp.
- White, R.S., and McKenzie, O.P., 1989. Volcanism at rifts. *Scient. American*, July 1989, 62-71.
- Whitford, D. J. 1975. Strontium isotopic studies of the volcanic rocks of the Sunda arc, Indonesia, and their petrogenetic implications. *Geochimica et cosmochimica acta* 39, 1287 -302.
- Whitford, D. J. and Jezek, P. A. (1979). Origin of Late Cenozoic lavas from the Banda Sea, Indonesia: trace element and Sr isotope evidence. *Contributions to mineralogy and petrology* 68' 141-50.
- Whitford, D. J. and Nicholls, I. A. (1976). Potassium variation in lavas across the Sunda arc in Java and Bali. *Volcanism in Australasia* (ed., R. I. Johnson), pp. 53-75. Elsevier, Amsterdam.
- Whitford, D. J., Compston, W., Nicholls, I. A., and Abbott, M. J. 1977. Geochemistry of Late Cenozoic lavas from eastern Indonesia: role of subducted sediments in petrogenesis' *Geology* 5, 571-5.
- Whitmore, D. P. and Matese, J. J., 1985. Periodic comet shower x and planet X. *Nature*, London, 313, 36.
- Whitmore, T. C. 1981. 'Wallace's Line and Plate Tectonics.' (Clarendon Press: Oxford.)
- Whitmore, T. C. 1987. Introduction. In: *Biogeographical Evolution of the Malay Archipelago*. (Ed. T. C. Whitmore.) 1-2. (Clarendon Press: Oxford.)
- Willett, R. W., 1950. The New Zealand Pleistocene snow line, climate conditions and suggested climatic effects. *N.Z. J. Technol.* 32B, 18-48.
- Wood, B., 1977. The evolution of early man. Cassell Australia Ltd., 124 pp.
- Wood, B. G. M. 1985. The mechanics of progressive deformation in crustal plates-a working model for Southeast Asia. *Geological society of Malaysia bulletin* 18, 55-99.
- Wood, B. L. 1962. Wakatipu 1:250,000 geol. Map Ser. Sheet 22, *Geol. Sur. N.Z.* 64.
- Wongsomsak, S. 1985. *Geomorphology and Land Use in Northeast Thailand Case Study Changwat Buri Ram*. MS. thesis, Kyoto University. 253 pp.
- Worsley, T.R. and Nance, R.D. (1989) Carbon redox and climate control through earth history: a speculative reconstruction' *Palaeogeog. Palaeoclimat. Palaeoecol.*, v. 75, pp 259-282'
- Wu, Jiquan. et al. "Distribution of unmelted impact ejecta in the Australasian tektite strewn field." May 1993. (*Eos. Transactions, American Geophysical Union* ; Vol. 74. No. 16. Suppl., p. 193)
- Wu, N.Q., Rousseau, D.D. & Liu, T.S., 1996. Land mollusk records from the Luochuan loess sequence and their paleoenvironmental significance. *Science in China*

- 39, 494-502.
- Wyllie, P. J., 1971. *The Dynamic Earth*. Wiley Interscience Edition. John Wiley and Sons, Inc., New York. 416 pp.
- Xu Rong-Hua, Scharer, U., and Allegre, C. J. 1985. Magnetism and metamorphism in the Lhasa Block (Tibet): a geochronological study. *Journal of Geology* 93, 41-57.
- Xu, Dao-Yi; Zang, Quinwen; Sun, Yi-Yin; Yan, Zheng; Chai, Zi-Fang and He Jin-Wen, eds. 1989: *Astrogeological events in China*. Geological Publishing House, Beijing, China.
- Yabuki, H. et al. "Fission track age and chemical composition of tektite from the remain of *Homo erectus erectus*." 1981. (Scientific Papers Of The Institute Of Physical And Chemical Research; Vol. 75, No. 2, p. 102-104)
- Yabuki, H., et al. "Tektite (?) from northern Thailand." 1981. (Inst. Phys. Chem. Res. (Rikagaku Kenkyusho), Sci. Pap.; 75, 1, p.41-42)
- Yan, Z., 1960. Stratigraphic division of the Quaternary sediments in Baoji area. *Quaternaria Sinica* 3, 88-103. (in Chinese)
- Yan Zheng, 1983. Fission track age and oxygen isotope composition of tektites. *Geology & Geochemistry*, 11, 25-29.
- Yan Zheng, 1986. Oxygen isotope and fission track age in tektites and impact glass. In Zhang Qin-wen and Xu Dao-yi (eds.), "Advance in Astrogeology", Oceanic Publishing House, Beijing, 141-150.
- Yan Zheng, Yuan Bao-Yin and Ye Lian-Fang, 1979. Spaltspurendatierung an Hainan-Insel-Tektiten (Legungmo). *Scientia Geologica Sinica*, 1, 37-42.
- Yan Zheng, Yuan Bao-Yin and Ye Lian-Fang, 1983. Oxygen isotope composition of tektites (Leigongmo) and Quaternary basalts from Hainan Island. *Scientia Geologica Sinica*, 4, 387-391.
- Yan Zheng, et al. "Fission-track dating a GEOL. SINICA ; Vol. 1979, No. 1, p.37-a2)
- Yang F-C., & Grote P.J., 2018a. Variation of fluorescence intensity in Pleistocene pollen and the correlation of fluvial deposits in the Khorat Plateau, Thailand, *Palynology*, 42:1, 28-35, DOI: 10.1080/01916122.2017.1303796
- Yang F-C., & Grote P.J., 2018b. Riverine vegetation and environments of a Late Pleistocene river terrace, Khorat Plateau, Southeast Asia, *Palynology*, 42:2, 158-167, DOI: 10.1080/01916122.2017.1296044
- Yeats, R.S., Sieh, K.E., and Allen, C.R., 1997. *The Geology of earthquakes*. Oxford University Press, New York.
- Yokohama, T. and Dharma, A. 1982. Magneto-stratigraphy of the Toba Tuffs in Sumatra, Indonesia. *Palaeomagnetic research in Southeast and East Asia*, CCOP/TP 13, 201-11
- Young, D. G., 1964. Stratigraphy and petrography of north-east Otago loess. *N. Z. J. Geol. Geophys.* 7, 839-63.
- Yuan, B. 1981: Preliminary discussion on the origin of tektites (in Chinese with English abstract). *Sci. Geol. Sinica* 4: 329-336.
- Yabuki, H. et al. "Fission track age and chemical composition of tektite from the remain of *Homo erectus erectus*." 1981. (Scientific Papers Of The Institute Of Physical And Chemical Research ; Vol. 75, No. 2, p. 102-104)
- Yabuki, H., et al. "Tektite (?) from northern Thailand." 1981. (INST. PHYS. CHEMICALS. (Rikagaku Kenkyusho), Sci. Pap. ; Vol. 75, No 1, p. 41-47;)
- Zahringer, J., 1963. K-Ar measurement of tektites. In: *Radioactive Dating*. Int. Atom. Energy Agency, Vienna. 289-305.
- Zahringer, J., and W. Gentner, 1963. Radiogenic and atmospheric argon content of tektites. *Nature*, 199, 583..
- Zebera, K. (1967): Moldavite-bearing sediments between Kotoseky and Holkov in South Bohemia. *Vstik Ustrednfno ustavu geologicdkho, LII*, 327-33.
- Zeitoun, V., Forestier, H., Auetrakulvit, P., Khaokhiew, C., Rasse, M., Davtian, G., Winayalai, C., Tiamtinkrit, C., 2012 Discovery of a prehistoric site at Sao Din (Nanoi, Nan province, Northern Thailand): Stone tools and new geological insights. *Découverte d'un site préhistorique à Sao Din (Nanoi, province de Nan, Nord de la Thaïlande): études lithique et géologique préliminaires*. *Comptes Rendus Palevol* 11, 575-580. doi:10.1016/j.crpv.2012.06.007
- Zhang Hunan. 1994. Some new advances on the division of Quaternary strata along Guangdong-Fujian-Hainan coast. *Guangdong Geology*, 9, 1, p.28-37.
- Zhang Qin-Wen, et al. 1988. New approach to the origin of the tektite in China. *LPI CONTRIBUTION*, 673, p.225-226.
- Zheng, H.B., An, Z.S. & Shaw, J., 1992. contributions to Chinese Plio-Pleistocene magnetostratigraphy. *Phys. Earth Planet. Inter.* 70, 146-153..
- Zhou Xinmin and Chen Tuhua. 1981. Composition and evolution of Cenozoic basaltic rocks in southeastern coastal provinces of China, *Collected Oceanic works*, Beiing 4,99-110.
- Zhu, X.M., 1958. Discussion on the problems of red layers intercalated in the loess section. *Quaternaria Sinica* 1, 74-82. (in Chinese).
- Zoller, W. H., Barrington, J. R., and Phelan Kotra, J. M., 1983. Iridium enrichment in air-borne particle from Killaua Volcano, January 1983. *Science*, 222, 1118-1121.

# INSTRUCTIONS TO AUTHORS FOR TGJ

The Thai Geoscience Journal (TGJ) is an international (Thai and English) journal publishing original research articles dealing with the geological sciences. Two types of articles are published in the Journal: Research Articles and Reviews. Research Articles are new original articles. Review Articles are those papers that summarize the current state of knowledge on specific fields of geosciences and discuss previously published research results, rather than report new results. **Copyright of all the contents of Thai Geoscience Journal belong to the Department of Mineral Resources in Thailand.**

The important instructions for manuscript submission are as follows:

## 1. TGJ member registration

Firstly, author must register the TGJ member before manuscript submission process. Please register **TGJ members online on TGJ website at <http://www.dmr.go.th/tgjdmr>.**

## 2. Submission of manuscripts

The second, authors must completely fill up the online manuscript submission form on TGJ website. The last important submission step is sending the required electronic manuscript (word file) and Figure files to TGJ editorial team via E-mail: [tgj.2020@gmail.com](mailto:tgj.2020@gmail.com). Editorial team will send you manuscript submission confirmation, manuscript number and all publication processes by email.

## 3. Electronic file requirements for submission

1) The manuscript should be prepared both in word and pdf files. TGJ manuscript form can download from TGJ website.

2) Figure files (JPEG, PNG, TIFF, EPS, PSD or AI); Resolution of all figures must be 300 – 600 dpi. The name of the figure file should be beginning with “Fig.” and followed by the number and name of figure that related to the manuscript. *For publication, the high resolution figure file (300-600 dpi) may be sent after the accepted manuscript process via E- mail: [tgj.2020@gmail.com](mailto:tgj.2020@gmail.com)*

## 4. Manuscripts Information

The main document, containing cover sheet, main text, acknowledgement (if applicable), references, figure, table, figure and table captions. Cover sheet: Cover sheet should contain 1) title, 2) full names and affiliations and the addresses of all authors and 3) postal and e-mail addresses, phone and fax numbers of the Corresponding author who will take responsibility for the proofs.

**Title:** A title is to be brief and summarizes the major results of the paper.

**Abstract:** An abstract should be condensation and concentrate on the essential qualities of the paper. All papers, excluding short notes, are to be accompanied by an abstract not exceeding 500 words or as appropriate.

**Key words:** Select keywords (should be 4-8 words or phrases) which identify the most important subjects covered by the paper and arrange them in alphabetical order

**Main text:** The main text should contain Introduction, Methods, Results, Discussions, and Conclusions, or as appropriate.

**Figures and tables:** Each figure and table should be sequentially placed in the main text near the first time mentioned. Figure captions and mentions in text begin with the term “Fig.” in bold type, followed by the figure number, also in bold type. Each table must have a heading. Captions and headings should be explicit enough that the reader can understand the significance of the illustration or table without reference to the text.

**Acknowledgements:** Brief acknowledgement of funding sources and assistance provided.

**References:** The references should be in a common citation format as shown below.

## 5. References Format Suggestion

In-text citation: the APA 6th style requires the references in content, tables and images should specify author-date, with the researcher’s surname followed by a comma (,) and the year of publication of the document behind the quote. If there are 2-5 authors, cite all names and use & before the last name for the first time, thereafter only the first name followed by et al. but if there are 6 or more authors, cite only the surname of the first author followed by et al. and the year. For examples as followings: (Mogen, 2001), (Jones & Miler, 2008), (Mogen, Jones & Miler, 2008), (Halano et al., 2009)

In Reference Page: Citation guide in APA 6th Style. For footnote citation should be at the end of manuscript. List authors in the reference as:

- For 1 author: Author, A. A.
- For 2 authors: Author, A. A., & Author B. B.
- For 3-7 authors: Author, A. A., Author, B. B., Author, C. C., Author, D. D., Author, E. E., Author, F. F., & Author, G. G.
- For 8 or more authors (list first six authors, add a three dot, then last author): Author, A. A., Author, B. B., Author, C. C., Author, D. D., Author, E. E., Author, F. F., ... Author, Z. Z.

**Types of cited references:**

### 1. Book

Author, A. A. (Year). *Title of book*. City, Country: Publisher’s Name.

Example:

Luffman, J. M., Bulleen, C. V., Liano, A. D., McLeod, P. K., Nash, E. O., & Neuman, C. C. (2004). *Information technology resources management* (2nd ed.). Upper Saddle River, NJ: John & Sons Press.

### 2. Book Chapter

Author, A. A. (Year). Title of chapter. In A. Editor, & B. Editor (Eds.), *Title of book* (pp.xx-xx). Location: Publisher.

Example:

Ogg, J. G., Ogg, G. M., & Gradstein, F. M. (2016). 5 - Cambrian. In J. G. Ogg, G. M. Ogg, & F.M. Gradstein (Eds.), *A Concise Geologic Time Scale* (pp. 41-55). UK: Elsevier.

### 3. Journal Article

Author1, A. A., & Author2, B. B. (Year). Title of article. *Journal name*, volume number (issue number), pages - pages. doi:xx. xxxxxxxxxx

Example:

Jackson, J. J., & Samuel, T. S. (2001). The impact of climate change on sea levels. *Journal of Environmental Science*, 55(4), 233-277. doi:10. 1070/8567-6582.33.5.



If you are interested in earth sciences  
Support geoscience with a subscription to

## Thai Geoscience Journal



The **Thai Geoscience Journal** publishes original research and review articles from the international community in all fields of geological sciences such as

Engineering Geology

Petrology

Medical Geology

Paleontology

Economic Geology

Geophysics

Tectonics

Structural Geology

Geochemistry

All articles published by the **Thai Geoscience Journal** are made freely and permanently accessible online immediately upon publication, **without subscription charges or registration barriers.**



Please scan for

**VISIT US**

**ARTICLES SUBMISSION**

**SUBSCRIPTION TO TGJ  
AND MORE INFORMATION**

### Contact

Mineral Resources Research and Development Center  
Department of Mineral Resources 75/10 Rama VI Road,  
Ratchatewi, Bangkok 10400 Phone: +66 2-6219731  
Website : <https://www.dmr.go.th/tgjdmr/>  
E-mail : [tgj.2020@gmail.com](mailto:tgj.2020@gmail.com)

Published by





## CONCEPT DESIGN

This logo composes of Abbreviations of Thai Geoscience Journal

**T = THAI G = GEOSCIENCE J = JOURNAL**

Coexistence of 3 abbreviations design in a concept of modernity blend with a Thainess. Modification of G alphabet in a shape of ammonoid shows relevance to geology and infinite development of Thai Geoscience Journal.

- **SCOPE AND AIM OF THAI GEOSCIENCE JOURNAL (TGJ):** TGJ is an international (Thai and English) journal publishing original research articles dealing with the geological sciences. It focuses, mainly but not exclusively, on: Sedimentology and Geomorphology, Palaeontology, Quaternary Geology and Environmental Change, Geological Hazards, Environmental Geosciences, Geophysics, Mineral and Petroleum Geology, Tectonics and Structural Geology, Geochemistry and Geochronology, Metamorphic Geology and Volcanic and Igneous Geology. Two types of articles are published in the Journal: Research Articles and Reviews. Research Articles are new original articles, normally not exceeding 25 pages. Review Articles are those papers that summarize the current state of knowledge on specific fields or topics of geosciences. They analyze and discuss previously published research results, rather than report new results. TGJ Aim is to provide valuable geoscience knowledge and information and push more inspiration for readers and researchers to produce treasure research in the future.
- **FEEDBACK AND CONTACT:** We welcome your feedback, comments and suggestions for the development of TGJ.

**Please contact:** Dr. Apsorn Sardud (Editor-in-Chief, TGJ)  
Department of Mineral Resources  
75/10 RamaVI Road Ratchathewee Bangkok 10400, Thailand



**Phone:** +66 (0)2 6219731



**Email:** [tgj.2020@gmail.com](mailto:tgj.2020@gmail.com)



**website:** <http://www.dmr.go.th>



## TGJ Contributors

Assoc. Prof. Dr. Apichet Boonsoong  
Dr. Apsorn Sardsud  
Prof. Dr. Che Aziz bin Ali  
Prof. Dr. Clive Burrett  
Assoc. Prof. Dr. Danupon Tonnyapas  
Dr. Dhiti Tulyatid

Mr. Jitisak Premmanee  
Prof. Dr. Katsumi Ueno  
Prof. Dr. Katsuo Sashida  
Prof. Dr. Ken-Ichiro Hisada  
Assoc. Prof. Dr. Kieren Howard  
Prof. Dr. Koji Wakita  
Assoc. Prof. Dr. Lindsay Zanno  
Dr. Mallika Nillorm  
Adj. Prof. Dr. Michael Ryan King  
Prof. Dr. Montri Choowong  
Assoc. Prof. Dr. Mongkol Udchachon  
Prof. Dr. Nigel C. Hughes  
Mr. Niwat Boonnop  
Asst. Prof. Nusara Surakotra  
Asst. Prof. Dr. Passakorn Pananont  
Dr. Phumee Srisuwan  
Prof. Dr. Pitsanupong Kanjanapayont  
Dr. Pol Chaodumrong  
Dr. Pradit Nulay  
Dr. Prinya Putthapiban  
Prof. Dr. Punya Charusiri  
Asst. Prof. Dr. Rattanaporn Hanta  
Assoc. Prof. Dr. Sachiko Agematsu-Watanabe  
Dr. Sasiwimol Nawawitphisit  
Dr. Seung-bae Lee  
Dr. Siriporn Soongpankhao  
Mr. Somchai Chaisen  
Asst. Prof. Dr. Sombat Yumuang

Dr. Surin Intayos  
Mr. Sutee Chongautchariyakul  
Mr. Thananchai Mahatthanachai  
Mr. Tritip Suppasoonthornkul  
Dr. Weerachat Wiwegwin  
Assoc. Prof. Dr. Yoshihito Kamata

Chiang Mai University, Thailand  
Department of Mineral Resources, Thailand  
Universiti Kebangsaan Malaysia, Malaysia  
Mahasarakham University, Thailand  
Prince of Songkla University, Thailand  
Coordinating Committee for Geoscience Programmes in East and Southeast Asia- CCOP, Thailand  
Department of Mineral Resources, Thailand  
Fukuoka University, Japan  
Mahidol University, Kanchanaburi campus, Thailand  
University of Tsukuba, Japan  
Kingsborough Community College, City University of New York, USA  
Yamaguchi University, Japan  
North Carolina State University, USA  
Department of Mineral Resources, Thailand  
Western Colorado University, USA  
Chulalongkorn University, Thailand  
Mahasarakham University, Thailand  
University of California, Riverside, USA  
Department of Mineral Resources, Thailand  
Khon Kaen University, Thailand  
Kasetsart University, Thailand  
Department of Mineral Fuels, Thailand  
Chulalongkorn University, Thailand  
Geological Society of Thailand, Thailand  
Department of Mineral Resources, Thailand  
Mahidol University, Kanchanaburi Campus, Thailand  
Department of Mineral Resources, Thailand  
Suranaree University of Technology, Thailand  
University of Tsukuba, Japan  
Department of Mineral Resources, Thailand  
Korea Institute of Geoscience and Mineral Resources, Republic of Korea  
Department of Mineral Resources, Thailand  
Department of Mineral Resources, Thailand  
Geo-Informatics and Space technology Development Agency, Ministry of Science and Technology (GiSTDA), Thailand  
Burapha University, Chanthaburi Campus, Thailand  
Department of Mineral Resources, Thailand  
Department of Mineral Fuels, Thailand  
Department of Primary Industries and Mines, Thailand  
Department of Mineral Resources, Thailand  
University of Tsukuba, Japan.



Thai Geoscience Journal  
Vol. 2 No. 1 (Special Issue)  
January 2021  
ISSN 2730-2695

## CONTENTS

---

### Honorary Editors

Dr. Sommai Techawan  
Mr. Niwat Maneekut  
Mr. Montri Luengingkasoot

### Advisory Editors

Prof. Dr. Clive Burrett  
Prof. Dr. Katsuo Sashida  
Prof. Dr. Nigel C. Hughes  
Prof. Dr. Punya Charusiri

### Editor In Chief

Dr. Apsorn Sardud

### Associate Editors

Prof. Dr. Che Aziz bin Ali  
Prof. Dr. Clive Burrett  
Assoc. Prof. Dr. Kieren Howard  
Dr. Pol Chaodumrong  
Prof. Dr. Punya Charusiri

### Published By

Department of Mineral Resources  
Geological Society of Thailand

- 1-15 The man who made an impact – the scientific work of Dr Sangad Bunopas  
**Clive Burrett**
- 16-109 Australasian Pleistocene cometary impact and its effects in Thailand  
**Sangad Bunopas**

